



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 04:51 PM UTC

PDB ID : 4M61 / pdb_00004m61
Title : Crystal structure of unliganded anti-DNA Fab A52
Authors : Stanfield, R.L.; Eilat, D.; Wilson, I.A.
Deposited on : 2013-08-08
Resolution : 1.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6728 (1.64-1.60)
Clashscore	190562	7023 (1.64-1.60)
Ramachandran outliers	187476	6898 (1.64-1.60)
Sidechain outliers	187428	6896 (1.64-1.60)
RSRZ outliers	180081	6727 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	219	4% 90% 9% .
1	C	219	5% 89% 10%
2	B	220	4% 90% 10%
2	D	220	7% 88% 10% ..

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7686 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab A52 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	219	Total	C	N	O	S	0	11	0
			1733	1084	285	352	12			
1	C	219	Total	C	N	O	S	0	9	0
			1735	1083	289	351	12			

- Molecule 2 is a protein called Fab A52 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	220	Total	C	N	O	S	0	10	0
			1689	1067	273	338	11			
2	D	220	Total	C	N	O	S	0	4	0
			1665	1048	273	336	8			

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	189	Total O 189 189	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	207	Total 207	O 207	0	0
4	C	211	Total 211	O 211	0	0
4	D	162	Total 162	O 162	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fab A52 light chain




- Molecule 1: Fab A52 light chain



- Molecule 2: Fab A52 heavy chain



- Molecule 2: Fab A52 heavy chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	52.78Å 61.38Å 121.22Å 90.00° 96.51° 90.00°	Depositor
Resolution (Å)	21.69 – 1.62 21.69 – 1.62	Depositor EDS
% Data completeness (in resolution range)	96.1 (21.69-1.62) 96.0 (21.69-1.62)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.61 (at 1.62Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.192 , 0.232 0.193 , 0.232	Depositor DCC
R_{free} test set	9435 reflections (9.33%)	wwPDB-VP
Wilson B-factor (Å ²)	15.6	Xtriage
Anisotropy	0.573	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 45.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7686	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 28.87 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.7091e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/1805	0.77	1/2452 (0.0%)
1	C	0.42	0/1801	0.76	0/2444
2	B	0.46	0/1761	0.80	1/2398 (0.0%)
2	D	0.47	1/1719 (0.1%)	0.79	2/2342 (0.1%)
All	All	0.45	1/7086 (0.0%)	0.78	4/9636 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	157	VAL	C-O	-7.27	1.15	1.23

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	75	SER	N-CA-C	-7.70	104.78	114.56
2	D	165	SER	N-CA-C	-7.49	103.25	112.38
2	D	43	LYS	N-CA-C	5.37	122.24	110.80
1	A	206	SER	N-CA-C	5.35	117.65	109.95

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	42	GLY	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1733	0	1686	16	0
1	C	1735	0	1680	14	0
2	B	1689	0	1671	13	0
2	D	1665	0	1626	16	0
3	A	20	0	0	0	0
3	B	30	0	0	1	0
3	C	25	0	0	0	0
3	D	20	0	0	1	0
4	A	189	0	0	3	0
4	B	207	0	0	1	0
4	C	211	0	0	3	0
4	D	162	0	0	3	0
All	All	7686	0	6663	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:156:SER:HB3	4:D:548:HOH:O	1.77	0.84
2:B:30[B]:THR:HG23	2:B:54:TYR:HB2	1.65	0.78
1:C:160:ARG:NH1	1:C:190[B]:GLU:OE2	2.19	0.74
1:A:3:MET:SD	4:A:570:HOH:O	2.54	0.66
2:B:51:ILE:HD13	2:B:72:VAL:HG13	1.80	0.63
1:A:60:ARG:HH21	1:A:69[A]:THR:HG22	1.66	0.59
2:D:82:GLN:NE2	4:D:532:HOH:O	2.35	0.59
1:A:206:SER:OG	1:A:207:THR:N	2.37	0.58
1:A:60:ARG:NH2	1:A:69[A]:THR:HG22	2.19	0.57
2:B:139:THR:HG21	2:B:143:VAL:HG23	1.86	0.57
2:D:65:LYS:NZ	3:D:303:SO4:O4	2.31	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:33:ASN:OD1	2:D:50:LYS:NZ	2.38	0.55
2:D:195:TRP:CG	2:D:196:PRO:HA	2.41	0.55
2:B:42:GLY:H	2:B:43[B]:LYS:HD3	1.72	0.55
1:A:154:LYS:NZ	1:A:200:GLU:OE1	2.42	0.53
2:B:195:TRP:CG	2:B:196:PRO:HA	2.43	0.53
2:D:216:LYS:HD3	2:D:218:GLU:OE2	2.09	0.53
1:C:200:GLU:HG2	1:C:211:VAL:HG12	1.90	0.53
2:D:156:SER:O	2:D:156:SER:OG	2.23	0.52
1:A:160:ARG:NH1	4:A:583:HOH:O	2.41	0.52
1:C:43:GLN:HB2	1:C:53:LEU:HD11	1.91	0.52
2:B:65:LYS:HE2	3:B:306:SO4:O1	2.10	0.52
2:D:152:TYR:HE2	2:D:155:GLU:O	1.93	0.51
1:C:187:THR:OG1	1:C:190[A]:GLU:HG3	2.11	0.50
1:C:133:GLY:C	1:C:188:LYS:HB2	2.37	0.49
1:C:21[B]:MET:HE1	1:C:92:TYR:CB	2.43	0.49
1:A:11:LEU:HD11	1:A:19[A]:VAL:HG23	1.95	0.48
2:B:41:HIS:HA	2:B:43[B]:LYS:HZ3	1.77	0.48
1:C:188:LYS:HE3	1:C:192:GLU:HG3	1.94	0.48
1:A:145:TYR:CG	1:A:146:PRO:HA	2.50	0.47
1:A:144:PHE:CZ	1:A:180[B]:MET:HG2	2.49	0.47
1:C:51:LYS:HE3	4:C:450:HOH:O	2.16	0.46
1:C:178:TYR:OH	4:C:585:HOH:O	2.21	0.46
1:A:43:GLN:HB2	1:A:53:LEU:HD11	1.97	0.46
2:D:41:HIS:HA	2:D:43:LYS:HE2	1.98	0.46
2:B:42:GLY:O	2:B:43[B]:LYS:HB2	2.15	0.45
1:A:200:GLU:HG2	1:A:211:VAL:HG12	1.99	0.45
1:C:180:MET:HE2	1:C:180:MET:HB2	1.89	0.45
2:D:11:LEU:HB2	2:D:154:PRO:HG3	1.99	0.45
1:A:11:LEU:HD11	1:A:19[A]:VAL:CG2	2.47	0.45
2:D:199:THR:HG23	2:D:216:LYS:HE3	1.98	0.45
2:D:216:LYS:CE	4:D:495:HOH:O	2.64	0.44
2:B:53:PRO:O	2:B:74:LYS:HD3	2.18	0.44
1:C:117:ALA:HB1	1:C:210:ILE:HD11	1.99	0.44
2:D:134:GLY:HA2	2:D:220:SER:OG	2.19	0.42
2:D:164:GLY:C	2:D:166:LEU:N	2.75	0.42
1:A:1:ASN:ND2	4:A:570:HOH:O	2.51	0.42
1:A:145:TYR:CD1	1:A:146:PRO:HA	2.54	0.42
1:A:60:ARG:HG2	1:A:64:VAL:HB	2.02	0.42
2:B:100:ARG:HA	2:B:100:ARG:HD2	1.87	0.42
2:B:130:PRO:HD3	2:B:215:LYS:HE2	2.01	0.42
2:D:164:GLY:C	2:D:166:LEU:H	2.28	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:63:LYS:HE2	4:B:591:HOH:O	2.20	0.41
2:B:161:TRP:CZ3	2:B:202[A]:CYS:HB3	2.55	0.41
2:D:152:TYR:CE2	2:D:155:GLU:O	2.74	0.41
1:A:159:GLU:OE2	1:A:161:GLN:HG2	2.21	0.40
1:C:21[B]:MET:HE2	1:C:21[B]:MET:HB2	1.93	0.40
1:C:155:ILE:HD11	1:C:184:LEU:HD21	2.04	0.40
1:C:51:LYS:NZ	4:C:574:HOH:O	2.49	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	228/219 (104%)	223 (98%)	5 (2%)	0	100	100
1	C	226/219 (103%)	218 (96%)	7 (3%)	1 (0%)	30	14
2	B	228/220 (104%)	224 (98%)	2 (1%)	2 (1%)	14	3
2	D	222/220 (101%)	217 (98%)	2 (1%)	3 (1%)	9	1
All	All	904/878 (103%)	882 (98%)	16 (2%)	6 (1%)	21	6

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	43[A]	LYS
2	B	43[B]	LYS
2	D	155	GLU
1	C	204	ALA
2	D	43	LYS
2	D	156	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	205/194 (106%)	202 (98%)	3 (2%)	57	34
1	C	203/194 (105%)	199 (98%)	4 (2%)	48	24
2	B	198/188 (105%)	195 (98%)	3 (2%)	57	34
2	D	192/188 (102%)	187 (97%)	5 (3%)	40	16
All	All	798/764 (104%)	783 (98%)	15 (2%)	53	26

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	161	GLN
1	A	185[A]	THR
1	A	185[B]	THR
2	B	176	LEU
2	B	211[A]	THR
2	B	211[B]	THR
1	C	10	SER
1	C	112	LYS
1	C	160	ARG
1	C	161	GLN
2	D	19	LYS
2	D	40	SER
2	D	156	SER
2	D	166	LEU
2	D	214	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	41	HIS
1	C	48	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

19 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	304	-	4,4,4	0.20	0	6,6,6	0.07	0
3	SO4	B	302	-	4,4,4	0.25	0	6,6,6	0.12	0
3	SO4	B	304	-	4,4,4	0.23	0	6,6,6	0.13	0
3	SO4	A	302	-	4,4,4	0.20	0	6,6,6	0.11	0
3	SO4	D	303	-	4,4,4	0.25	0	6,6,6	0.11	0
3	SO4	D	302	-	4,4,4	0.26	0	6,6,6	0.13	0
3	SO4	B	301	-	4,4,4	0.24	0	6,6,6	0.19	0
3	SO4	A	301	-	4,4,4	0.24	0	6,6,6	0.20	0
3	SO4	C	302	-	4,4,4	0.23	0	6,6,6	0.26	0
3	SO4	D	304	-	4,4,4	0.26	0	6,6,6	0.12	0
3	SO4	B	305	-	4,4,4	0.22	0	6,6,6	0.12	0
3	SO4	B	306	-	4,4,4	0.23	0	6,6,6	0.15	0
3	SO4	C	301	-	4,4,4	0.26	0	6,6,6	0.07	0
3	SO4	C	303	-	4,4,4	0.20	0	6,6,6	0.16	0
3	SO4	C	304	-	4,4,4	0.24	0	6,6,6	0.12	0
3	SO4	C	305	-	4,4,4	0.24	0	6,6,6	0.12	0
3	SO4	A	303	-	4,4,4	0.25	0	6,6,6	0.05	0
3	SO4	D	301	-	4,4,4	0.20	0	6,6,6	0.14	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	B	303	-	4,4,4	0.24	0	6,6,6	0.15	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	303	SO4	1	0
3	B	306	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	219/219 (100%)	0.05	8 (3%)	45	48	6, 15, 36, 51	11 (5%)
1	C	219/219 (100%)	0.00	11 (5%)	34	36	7, 16, 35, 45	9 (4%)
2	B	220/220 (100%)	0.22	8 (3%)	46	49	6, 18, 34, 44	10 (4%)
2	D	220/220 (100%)	0.37	15 (6%)	23	25	10, 20, 38, 53	4 (1%)
All	All	878/878 (100%)	0.16	42 (4%)	35	38	6, 17, 37, 53	34 (3%)

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	219	CYS	4.0
1	C	219	CYS	3.7
1	C	131	THR	3.7
1	A	207	THR	3.7
2	B	101	LEU	3.7
2	D	166	LEU	3.6
2	B	42	GLY	3.5
2	D	217	LEU	3.3
1	C	204	ALA	3.3
1	A	161	GLN	3.1
2	D	136	GLY	3.0
2	D	168	SER	2.9
2	D	42	GLY	2.9
1	A	204	ALA	2.8
1	A	205	THR	2.8
1	C	207	THR	2.7
2	B	54	TYR	2.7
1	A	203	HIS	2.7
2	D	214	ASP	2.6
1	C	127	SER	2.6
2	D	1	GLU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	134	GLY	2.4
2	B	55	TYR	2.4
2	D	55	TYR	2.4
2	B	44	SER	2.3
2	D	207	PRO	2.3
1	A	131	THR	2.3
2	B	66	GLY	2.3
1	C	205	THR	2.3
1	C	206	SER	2.3
1	A	1	ASN	2.3
2	D	167	SER	2.3
2	B	139	THR	2.2
1	C	218	GLU	2.2
1	C	203	HIS	2.2
2	D	41	HIS	2.2
2	D	210	SER	2.1
1	C	128	GLU	2.1
2	B	56	GLY	2.1
2	D	164	GLY	2.1
1	C	130	LEU	2.1
2	D	220	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	B	306	5/5	0.75	0.11	47,51,59,62	0
3	SO4	A	304	5/5	0.77	0.11	49,54,62,70	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	B	305	5/5	0.78	0.11	49,54,58,63	0
3	SO4	B	303	5/5	0.78	0.12	42,49,54,65	0
3	SO4	C	305	5/5	0.78	0.11	38,44,55,57	0
3	SO4	C	304	5/5	0.81	0.11	38,42,45,46	0
3	SO4	B	304	5/5	0.81	0.10	36,43,51,58	0
3	SO4	D	304	5/5	0.83	0.10	47,49,53,60	0
3	SO4	D	303	5/5	0.85	0.09	33,41,52,57	0
3	SO4	C	302	5/5	0.85	0.11	31,32,43,44	0
3	SO4	C	303	5/5	0.86	0.09	36,38,50,53	0
3	SO4	B	302	5/5	0.92	0.08	29,32,41,44	0
3	SO4	B	301	5/5	0.93	0.09	27,34,42,42	0
3	SO4	A	302	5/5	0.94	0.09	29,30,37,37	0
3	SO4	D	302	5/5	0.94	0.08	34,34,39,47	0
3	SO4	A	303	5/5	0.95	0.08	30,34,36,43	0
3	SO4	D	301	5/5	0.96	0.10	26,27,32,34	0
3	SO4	C	301	5/5	0.96	0.07	27,28,32,36	0
3	SO4	A	301	5/5	0.97	0.08	23,24,26,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.