



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 14, 2026 – 05:51 PM UTC

PDB ID : 4M6D / pdb_00004m6d
Title : Crystal structure of the aptamer minF-lysozyme complex.
Authors : Malashkevich, V.N.; Padlan, F.C.; Toro, R.; Girvin, M.; Almo, S.C.; New York
Structural Genomics Research Consortium (NYSGRG)
Deposited on : 2013-08-09
Resolution : 2.68 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5070 (2.70-2.66)
Clashscore	190562	5409 (2.70-2.66)
Ramachandran outliers	187476	5324 (2.70-2.66)
Sidechain outliers	187428	5324 (2.70-2.66)
RSRZ outliers	180081	5070 (2.70-2.66)
RNA backbone	3983	1075 (2.90-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	129	55% 42% .
1	C	129	57% 39% ..
1	E	129	59% 38% ..
1	G	129	64% 33% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	I	129	 56% 38% 5% •
1	K	129	 60% 30% 9% •
2	B	45	 24% 47% 18% • 9%
2	D	45	 36% 40% 18% 7%
2	F	45	 44% 33% 13% 9%
2	H	45	 33% 53% 7% 7%
2	J	45	 40% 40% 11% 9%
2	L	45	 22% 47% 24% • •

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11432 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lysozyme C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	129	Total	C	N	O	S	0	0	0
			1001	613	193	185	10			
1	C	129	Total	C	N	O	S	0	0	0
			1001	613	193	185	10			
1	E	129	Total	C	N	O	S	0	0	0
			1001	613	193	185	10			
1	G	129	Total	C	N	O	S	0	0	0
			1001	613	193	185	10			
1	I	129	Total	C	N	O	S	1	0	0
			1001	613	193	185	10			
1	K	129	Total	C	N	O	S	0	0	0
			1001	613	193	185	10			

- Molecule 2 is a RNA chain called aptamer.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	41	Total	C	N	O	P	0	0	0
			878	392	160	285	41			
2	D	42	Total	C	N	O	P	0	0	0
			901	402	165	292	42			
2	F	41	Total	C	N	O	P	0	0	0
			878	392	160	285	41			
2	H	42	Total	C	N	O	P	0	0	0
			901	402	165	292	42			
2	J	41	Total	C	N	O	P	0	0	0
			878	392	160	285	41			
2	L	43	Total	C	N	O	P	0	0	0
			921	411	168	299	43			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	5	Total O 5 5	0	0
3	B	8	Total O 8 8	0	0
3	C	5	Total O 5 5	0	0
3	D	4	Total O 4 4	0	0
3	E	5	Total O 5 5	0	0
3	F	11	Total O 11 11	0	0
3	G	5	Total O 5 5	0	0
3	H	3	Total O 3 3	0	0
3	I	7	Total O 7 7	0	0
3	J	8	Total O 8 8	0	0
3	K	5	Total O 5 5	0	0
3	L	3	Total O 3 3	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Lysozyme C




• Molecule 1: Lysozyme C




• Molecule 1: Lysozyme C




• Molecule 1: Lysozyme C






• Molecule 1: Lysozyme C

Chain I: 56% 38% 5% •




• Molecule 1: Lysozyme C

Chain K: 60% 30% 9% •




• Molecule 2: aptamer

Chain B: 24% 47% 18% 9% •



• Molecule 2: aptamer

Chain D: 36% 40% 18% 7% •




• Molecule 2: aptamer

Chain F: 44% 33% 13% 9% •



• Molecule 2: aptamer



Chain H: 33% 53% 7% 7% •



● Molecule 2: aptamer



● Molecule 2: aptamer



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	43.91Å 132.29Å 131.54Å 118.59° 96.39° 96.27°	Depositor
Resolution (Å)	38.04 – 2.68 38.04 – 2.68	Depositor EDS
% Data completeness (in resolution range)	79.5 (38.04-2.68) 79.6 (38.04-2.68)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.20 (at 2.69Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.165 , 0.216 0.172 , 0.212	Depositor DCC
R_{free} test set	2859 reflections (3.98%)	wwPDB-VP
Wilson B-factor (Å ²)	44.3	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 67.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.390 for h,-h-k-l,k 0.390 for h,l,-h-k-l 0.018 for -h,-l,-k 0.018 for -h,h+k+l,-l 0.016 for -h,-k,h+k+l	Xtriage
Reported twinning fraction	0.351 for H, K, L 0.014 for -H, H+K+L, -L 0.015 for -H, -L, -K 0.344 for H, L, -H-K-L 0.267 for H, -H-K-L, K 0.009 for -H, -K, H+K+L	Depositor
Outliers	0 of 56402 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11432	wwPDB-VP
Average B, all atoms (Å ²)	122.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.09	0/1021	1.14	0/1379
1	C	1.02	0/1021	1.13	1/1379 (0.1%)
1	E	0.98	1/1021 (0.1%)	1.13	3/1379 (0.2%)
1	G	0.95	0/1021	1.15	2/1379 (0.1%)
1	I	1.36	5/1021 (0.5%)	1.34	8/1379 (0.6%)
1	K	1.34	0/1021	1.29	6/1379 (0.4%)
2	B	0.59	1/982 (0.1%)	0.86	1/1529 (0.1%)
2	D	0.45	0/1008	0.80	1/1570 (0.1%)
2	F	0.47	0/982	0.75	0/1529
2	H	0.53	0/1008	0.85	2/1570 (0.1%)
2	J	0.50	0/982	0.81	0/1529
2	L	0.52	0/1030	0.88	3/1604 (0.2%)
All	All	0.88	7/12118 (0.1%)	1.02	27/17605 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	K	0	1
All	All	0	2

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	I	64	CYS	C-O	6.04	1.31	1.23
1	E	40	THR	N-CA	5.83	1.53	1.46
2	B	20	A	O3'-P	-5.40	1.53	1.61
1	I	2	VAL	C-O	5.25	1.29	1.24
1	I	39	ASN	N-CA	5.14	1.53	1.46

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	L	9	G	C2'-C3'-O3'	9.12	123.18	109.50
1	I	73	ARG	N-CA-C	8.65	125.47	114.31
1	I	86	SER	N-CA-C	-8.09	103.39	113.18
1	E	73	ARG	N-CA-C	6.97	122.20	112.93
1	I	53	TYR	N-CA-C	6.06	118.83	109.07

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	100	SER	Peptide
1	K	101	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1001	0	959	50	0
1	C	1001	0	959	44	0
1	E	1001	0	959	44	0
1	G	1001	0	959	30	0
1	I	1001	0	959	48	0
1	K	1001	0	959	40	0
2	B	878	0	443	26	0
2	D	901	0	454	15	0
2	F	878	0	443	16	0
2	H	901	0	454	11	0
2	J	878	0	443	17	0
2	L	921	0	465	29	0
3	A	5	0	0	0	0
3	B	8	0	0	0	0
3	C	5	0	0	0	0
3	D	4	0	0	1	0
3	E	5	0	0	0	0
3	F	11	0	0	1	0
3	G	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	H	3	0	0	0	0
3	I	7	0	0	0	0
3	J	8	0	0	2	0
3	K	5	0	0	0	0
3	L	3	0	0	0	0
All	All	11432	0	8456	363	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 363 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:10:G:H2'	2:D:11:C:H5'	1.36	1.06
2:F:12:G:H1	2:F:48:C:H42	1.04	0.96
1:A:21:ARG:HG2	1:A:21:ARG:HH11	1.31	0.94
2:J:11:C:H2'	2:J:11:C:O2	1.64	0.94
1:K:95:ALA:HA	1:K:108:TRP:HH2	1.33	0.93

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	127/129 (98%)	108 (85%)	19 (15%)	0	100	100
1	C	127/129 (98%)	114 (90%)	12 (9%)	1 (1%)	16	34
1	E	127/129 (98%)	112 (88%)	14 (11%)	1 (1%)	16	34
1	G	127/129 (98%)	115 (91%)	11 (9%)	1 (1%)	16	34
1	I	127/129 (98%)	116 (91%)	11 (9%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	K	127/129 (98%)	114 (90%)	13 (10%)	0	100	100
All	All	762/774 (98%)	679 (89%)	80 (10%)	3 (0%)	30	51

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	109	VAL
1	C	107	ALA
1	E	109	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	105/105 (100%)	98 (93%)	7 (7%)	15	32
1	C	105/105 (100%)	98 (93%)	7 (7%)	15	32
1	E	105/105 (100%)	99 (94%)	6 (6%)	18	40
1	G	105/105 (100%)	96 (91%)	9 (9%)	10	22
1	I	105/105 (100%)	96 (91%)	9 (9%)	10	22
1	K	105/105 (100%)	91 (87%)	14 (13%)	4	8
All	All	630/630 (100%)	578 (92%)	52 (8%)	10	24

5 of 52 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	109	VAL
1	I	99	VAL
1	K	108	TRP
1	I	21	ARG
1	I	78	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	41	GLN
1	G	113	ASN
1	K	113	ASN
1	K	27	ASN
1	C	59	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	40/45 (88%)	14 (35%)	1 (2%)
2	D	41/45 (91%)	17 (41%)	2 (4%)
2	F	40/45 (88%)	11 (27%)	1 (2%)
2	H	42/45 (93%)	13 (30%)	5 (11%)
2	J	40/45 (88%)	12 (30%)	4 (10%)
2	L	43/45 (95%)	18 (41%)	5 (11%)
All	All	246/270 (91%)	85 (34%)	18 (7%)

5 of 85 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	11	C
2	B	15	U
2	B	17	A
2	B	24	C
2	B	26	G

5 of 18 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	L	36	G
2	L	40	A
2	L	38	U
2	H	40	A
2	L	9	G

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	129/129 (100%)	-0.85	0 100 100	57, 90, 131, 171	0
1	C	129/129 (100%)	-0.88	0 100 100	60, 97, 140, 160	1 (0%)
1	E	129/129 (100%)	-0.86	0 100 100	57, 103, 154, 169	0
1	G	129/129 (100%)	-0.82	0 100 100	68, 100, 146, 162	1 (0%)
1	I	129/129 (100%)	-1.06	0 100 100	44, 72, 108, 136	0
1	K	129/129 (100%)	-0.95	0 100 100	44, 71, 115, 132	1 (0%)
2	B	41/45 (91%)	-0.21	0 100 100	106, 151, 230, 278	0
2	D	42/45 (93%)	-0.27	0 100 100	134, 180, 212, 234	0
2	F	41/45 (91%)	-0.40	0 100 100	110, 154, 202, 231	0
2	H	42/45 (93%)	-0.75	0 100 100	105, 135, 186, 229	0
2	J	41/45 (91%)	-0.61	0 100 100	83, 129, 204, 207	0
2	L	43/45 (95%)	-0.64	0 100 100	106, 140, 194, 221	0
All	All	1024/1044 (98%)	-0.80	0 100 100	44, 100, 186, 278	3 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.