



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 08:42 AM UTC

PDB ID : 6M7Y / pdb_00006m7y
Title : Dehydratase, NisB, bound to a non-eliminable substrate analog
Authors : Cogan, D.P.; Nair, S.K.
Deposited on : 2018-08-21
Resolution : 2.79 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	996	73% 21%
1	B	996	72% 21%
2	C	34	24% 12% 12% 53%
2	D	34	32% 9% 6% 53%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 16392 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nisin biosynthesis protein NisB.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	952	Total	C	N	O	S	Se	0	1	0
			7947	5125	1298	1502	9	13			
1	B	948	Total	C	N	O	S	Se	0	7	0
			7944	5119	1299	1503	10	13			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP P20103
A	-1	SER	-	expression tag	UNP P20103
A	0	HIS	-	expression tag	UNP P20103
A	1	MSE	-	expression tag	UNP P20103
A	169	CYS	VAL	engineered mutation	UNP P20103
B	-2	GLY	-	expression tag	UNP P20103
B	-1	SER	-	expression tag	UNP P20103
B	0	HIS	-	expression tag	UNP P20103
B	1	MSE	-	expression tag	UNP P20103
B	169	CYS	VAL	engineered mutation	UNP P20103

- Molecule 2 is a protein called Lantibiotic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	16	Total	C	N	O	S	0	0	0
			127	81	21	24	1			
2	D	16	Total	C	N	O	S	0	0	0
			127	81	21	24	1			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-12	CYS	SER	engineered mutation	UNP Q7BB86
C	3	J9A	SER	engineered mutation	UNP Q7BB86

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	7	ALA	CYS	engineered mutation	UNP Q7BB86
C	11	ALA	CYS	engineered mutation	UNP Q7BB86
D	-12	CYS	SER	engineered mutation	UNP Q7BB86
D	3	J9A	SER	engineered mutation	UNP Q7BB86
D	7	ALA	CYS	engineered mutation	UNP Q7BB86
D	11	ALA	CYS	engineered mutation	UNP Q7BB86


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	126	Total	O	0	0
			126	126		
3	B	116	Total	O	0	0
			116	116		
3	C	2	Total	O	0	0
			2	2		
3	D	3	Total	O	0	0
			3	3		


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Nisin biosynthesis protein NisB



• Molecule 1: Nisin biosynthesis protein NisB





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	99.64Å 107.14Å 135.44Å 90.00° 109.57° 90.00°	Depositor
Resolution (Å)	29.39 – 2.79 29.39 – 2.79	Depositor EDS
% Data completeness (in resolution range)	93.1 (29.39-2.79) 93.0 (29.39-2.79)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.98 (at 2.80Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.185 , 0.270 0.192 , 0.274	Depositor DCC
R_{free} test set	3032 reflections (4.55%)	wwPDB-VP
Wilson B-factor (Å ²)	44.6	Xtriage
Anisotropy	0.127	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 47.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.019 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	16392	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: J9A

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.46	0/8095	0.68	0/10892
1	B	0.45	1/8092 (0.0%)	0.77	12/10893 (0.1%)
2	C	0.55	0/110	0.83	0/144
2	D	0.43	0/110	0.57	0/144
All	All	0.45	1/16407 (0.0%)	0.72	12/22073 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
2	C	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	132	MSE	C-O	-5.18	1.17	1.24

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	975	GLU	N-CA-CB	20.69	145.46	110.49
1	B	974	LYS	N-CA-C	-16.59	83.99	110.20
1	B	372	HIS	CA-C-N	14.01	137.36	119.84
1	B	372	HIS	C-N-CA	14.01	137.36	119.84
1	B	975	GLU	N-CA-C	-11.01	87.36	110.80

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	372	HIS	Peptide
2	C	2	THR	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7947	0	7907	147	0
1	B	7944	0	7866	180	0
2	C	127	0	113	4	0
2	D	127	0	114	6	0
3	A	126	0	0	4	0
3	B	116	0	0	3	0
3	C	2	0	0	0	0
3	D	3	0	0	1	0
All	All	16392	0	16000	326	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 326 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:977:LEU:HD23	1:B:978:ILE:HG13	1.42	1.00
1:A:628:THR:HG22	1:A:631:GLU:OE1	1.68	0.93
1:B:977:LEU:CD2	1:B:978:ILE:HG13	1.98	0.93
1:B:261[B]:GLN:HG3	1:B:262:LYS:HD3	1.51	0.92
1:A:232:LEU:HD13	1:A:510:MSE:HE1	1.50	0.91

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	947/996 (95%)	879 (93%)	55 (6%)	13 (1%)	9	30
1	B	949/996 (95%)	861 (91%)	70 (7%)	18 (2%)	6	22
2	C	11/34 (32%)	6 (54%)	2 (18%)	3 (27%)	0	0
2	D	11/34 (32%)	11 (100%)	0	0	100	100
All	All	1918/2060 (93%)	1757 (92%)	127 (7%)	34 (2%)	6	23

5 of 34 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	255	ILE
1	A	541	ASP
1	A	719	ILE
1	A	779	PRO
1	B	373	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	891/921 (97%)	871 (98%)	20 (2%)	45	78
1	B	888/921 (96%)	862 (97%)	26 (3%)	37	73
2	C	13/28 (46%)	11 (85%)	2 (15%)	2	10
2	D	13/28 (46%)	12 (92%)	1 (8%)	12	36
All	All	1805/1898 (95%)	1756 (97%)	49 (3%)	40	74

5 of 49 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	155	VAL
1	B	449	LEU
1	B	165	GLU
1	B	233	SER
1	B	583	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	410	ASN
1	B	735	ASN
1	B	503	ASN
1	B	881	ASN
1	A	307	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	J9A	D	3	2	13,14,15	1.77	2 (15%)	14,17,19	1.08	0
2	J9A	C	3	2	13,14,15	1.85	2 (15%)	14,17,19	1.25	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	J9A	D	3	2	-	3/14/16/18	-
2	J9A	C	3	2	-	8/14/16/18	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	3	J9A	C06-N07	5.54	1.46	1.33
2	D	3	J9A	C06-N07	5.13	1.45	1.33
2	D	3	J9A	O14-C06	-2.33	1.18	1.23
2	C	3	J9A	O14-C06	-2.04	1.19	1.23

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	3	J9A	C04-C05-C06	2.16	115.17	110.76

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	3	J9A	C03-C04-C05-C06
2	C	3	J9A	C03-C04-C05-N15
2	C	3	J9A	C02-C03-C04-C05
2	C	3	J9A	O01-C02-C03-C04
2	C	3	J9A	O16-C02-C03-C04

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	3	J9A	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	939/996 (94%)	-0.36	9 (0%) 79 72	18, 41, 73, 120	1 (0%)
1	B	935/996 (93%)	-0.20	16 (1%) 69 60	16, 46, 87, 128	7 (0%)
2	C	15/34 (44%)	0.73	2 (13%) 7 5	31, 47, 109, 116	0
2	D	15/34 (44%)	0.53	2 (13%) 7 5	32, 52, 73, 84	0
All	All	1904/2060 (92%)	-0.27	29 (1%) 72 63	16, 43, 81, 128	8 (0%)

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	2	THR	5.3
2	C	4	ILE	4.9
1	B	717	ALA	4.7
1	B	975	GLU	3.6
2	C	2	THR	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	J9A	D	3	15/16	0.86	0.15	33,62,82,87	0
2	J9A	C	3	15/16	0.90	0.13	41,63,112,116	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.