



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:47 AM UTC

PDB ID : 7M7V / pdb_00007m7v
Title : Crystal Structure of Mtb Pks13 Thioesterase domain in complex with Compound 6
Authors : Aggarwal, A.; Sacchettini, J.C.
Deposited on : 2021-03-29
Resolution : 2.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	13% 85% 9% 7%
1	B	286	16% 84% 12% 5%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4191 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Polyketide synthase Pks13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	267	Total	C	N	O	S	0	0	0
			2021	1287	361	368	5			
1	B	272	Total	C	N	O	S	0	0	0
			2014	1281	360	368	5			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1448	SER	-	expression tag	UNP I6X8D2
A	1449	ASN	-	expression tag	UNP I6X8D2
A	1450	ALA	-	expression tag	UNP I6X8D2
B	1448	SER	-	expression tag	UNP I6X8D2
B	1449	ASN	-	expression tag	UNP I6X8D2
B	1450	ALA	-	expression tag	UNP I6X8D2

- Molecule 2 is 5-hydroxy-2-(4-hydroxyphenyl)-N-methyl-4-[(2-oxa-6-azaspiro[3.4]octan-6-yl)methyl]-1-benzofuran-3-carboxamide (CCD ID: Z3J) (formula: C₂₃H₂₄N₂O₅) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			30	23	2	5		
2	B	1	Total	C	N	O	0	0
			30	23	2	5		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	70	Total	O	0	0
			70	70		
3	B	26	Total	O	0	0
			26	26		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Polyketide synthase Pks13



• Molecule 1: Polyketide synthase Pks13



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	89.97Å 110.28Å 57.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.53 – 2.29 48.53 – 2.29	Depositor EDS
% Data completeness (in resolution range)	97.7 (48.53-2.29) 97.9 (48.53-2.29)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.67 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.223 , 0.269 (Not available) , 0.262	Depositor DCC
R_{free} test set	1304 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	45.9	Xtriage
Anisotropy	0.450	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 52.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4191	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: Z3J

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/2068	0.58	0/2810
1	B	0.38	0/2060	0.56	0/2809
All	All	0.40	0/4128	0.57	0/5619

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2021	0	1923	13	0
1	B	2014	0	1902	22	0
2	A	30	0	0	0	0
2	B	30	0	0	0	0
3	A	70	0	0	1	0
3	B	26	0	0	0	0
All	All	4191	0	3825	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1565:GLY:HA3	1:B:1676:VAL:HG23	1.46	0.95
1:A:1468:LYS:NZ	1:A:1493:ALA:O	2.07	0.86
1:B:1534:LEU:HD12	1:B:1643:ILE:HG13	1.66	0.78
1:A:1458:THR:HG23	1:A:1498:TYR:CE1	2.35	0.61
1:B:1625:ILE:HB	1:B:1630:ILE:HD11	1.82	0.61
1:B:1547:ARG:NH1	1:B:1652:ASP:OD1	2.34	0.59
1:B:1482:VAL:HG13	1:B:1485:GLU:OE1	2.02	0.59
1:A:1544:ILE:HD13	1:A:1649:GLN:HB2	1.85	0.58
1:B:1534:LEU:CD1	1:B:1643:ILE:HG13	2.32	0.58
1:A:1690:LEU:HD23	1:B:1694:PRO:HG2	1.86	0.57
1:A:1695:ILE:HG22	1:A:1710:LYS:HD3	1.88	0.56
1:B:1534:LEU:HB2	1:B:1643:ILE:HD11	1.88	0.56
1:A:1506:SER:OG	1:A:1509:GLU:HG3	2.07	0.55
1:B:1544:ILE:HD11	1:B:1651:TYR:HB2	1.92	0.52
1:A:1533:SER:OG	1:A:1534:LEU:N	2.43	0.51
1:B:1575:ILE:HG22	1:B:1602:LEU:HD21	1.92	0.51
1:B:1703:ILE:HA	1:B:1708:ILE:HD13	1.94	0.50
1:B:1490:ARG:HG3	1:B:1709:ALA:HA	1.97	0.46
1:A:1492:PRO:HG2	1:A:1495:THR:OG1	2.16	0.46
1:A:1471:VAL:O	1:A:1497:MET:HA	2.16	0.45
1:B:1488:LEU:HD13	1:B:1497:MET:HE3	1.98	0.44
1:B:1701:GLN:HG3	1:B:1707:ILE:HD12	1.99	0.44
1:B:1547:ARG:HD2	1:B:1547:ARG:HA	1.72	0.43
1:B:1533:SER:OG	1:B:1534:LEU:N	2.49	0.43
1:A:1503:VAL:HG23	1:A:1510:ARG:HG2	2.01	0.43
1:A:1631:GLU:O	1:A:1635:THR:HG23	2.20	0.42
1:B:1507:ILE:HD13	1:B:1507:ILE:HA	1.82	0.42
1:B:1612:ARG:HA	1:B:1612:ARG:HD3	1.88	0.42
1:B:1661:ASP:OD1	1:B:1696:GLY:HA2	2.20	0.42
1:A:1576:ARG:HD2	1:A:1603:GLU:OE2	2.20	0.41
1:B:1595:PRO:HG3	1:B:1621:SER:HB3	2.03	0.41
1:B:1500:PHE:HD2	1:B:1514:TYR:CD1	2.39	0.41
3:A:1959:HOH:O	1:B:1688:SER:HB3	2.20	0.40
1:A:1473:VAL:HA	1:A:1530:VAL:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	261/286 (91%)	256 (98%)	5 (2%)	0	100	100
1	B	268/286 (94%)	263 (98%)	5 (2%)	0	100	100
All	All	529/572 (92%)	519 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	160/231 (69%)	160 (100%)	0	100	100
1	B	188/231 (81%)	188 (100%)	0	100	100
All	All	348/462 (75%)	348 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1699	HIS
1	A	1721	GLN
1	B	1513	GLN
1	B	1701	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	Z3J	B	1801	-	34,34,34	1.89	7 (20%)	39,51,51	2.25	15 (38%)
2	Z3J	A	1801	-	34,34,34	1.83	5 (14%)	39,51,51	2.20	16 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	Z3J	B	1801	-	-	6/14/33/33	0/5/5/5
2	Z3J	A	1801	-	-	2/14/33/33	0/5/5/5

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1801	Z3J	C03-N02	5.74	1.41	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1801	Z3J	C03-N02	5.54	1.41	1.33
2	B	1801	Z3J	C24-C06	5.14	1.55	1.47
2	A	1801	Z3J	C24-C06	4.53	1.54	1.47
2	B	1801	Z3J	C05-C06	3.34	1.40	1.36
2	A	1801	Z3J	C05-C06	3.22	1.40	1.36
2	A	1801	Z3J	C14-N15	2.94	1.53	1.47
2	B	1801	Z3J	C14-C13	-2.53	1.47	1.51
2	B	1801	Z3J	C22-N15	2.28	1.48	1.46
2	A	1801	Z3J	C14-C13	-2.17	1.48	1.51
2	B	1801	Z3J	C16-N15	2.16	1.51	1.47
2	B	1801	Z3J	C14-N15	2.12	1.51	1.47

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1801	Z3J	C30-C24-C25	-5.55	111.51	118.57
2	A	1801	Z3J	C30-C24-C25	-5.44	111.65	118.57
2	B	1801	Z3J	C26-C25-C24	3.96	125.03	120.80
2	B	1801	Z3J	O20-C19-C18	-3.89	89.98	91.89
2	B	1801	Z3J	C29-C30-C24	3.87	124.93	120.80
2	A	1801	Z3J	C29-C30-C24	3.82	124.89	120.80
2	A	1801	Z3J	C01-N02-C03	3.80	125.77	121.90
2	B	1801	Z3J	O07-C06-C24	3.70	118.66	114.45
2	B	1801	Z3J	C13-C14-N15	-3.69	104.39	112.55
2	A	1801	Z3J	C13-C14-N15	-3.63	104.52	112.55
2	A	1801	Z3J	C08-O07-C06	3.62	109.73	106.25
2	A	1801	Z3J	C26-C25-C24	3.62	124.67	120.80
2	B	1801	Z3J	O07-C06-C05	-3.50	108.02	111.18
2	A	1801	Z3J	O04-C03-N02	-3.42	117.07	122.84
2	A	1801	Z3J	C25-C24-C06	3.32	125.60	120.71
2	B	1801	Z3J	C08-O07-C06	3.27	109.39	106.25
2	A	1801	Z3J	O07-C06-C05	-3.22	108.27	111.18
2	B	1801	Z3J	C25-C24-C06	3.14	125.33	120.71
2	A	1801	Z3J	O20-C19-C18	-2.91	90.46	91.89
2	B	1801	Z3J	O04-C03-N02	-2.90	117.94	122.84
2	A	1801	Z3J	C14-C13-C11	-2.64	116.88	120.22
2	A	1801	Z3J	O07-C06-C24	2.57	117.37	114.45
2	A	1801	Z3J	C14-N15-C22	-2.53	109.55	112.89
2	B	1801	Z3J	C14-N15-C16	-2.52	104.03	113.67
2	B	1801	Z3J	C14-N15-C22	-2.46	109.64	112.89
2	B	1801	Z3J	O20-C21-C18	-2.40	90.71	91.89
2	B	1801	Z3J	C01-N02-C03	2.37	124.31	121.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1801	Z3J	O20-C21-C18	-2.32	90.75	91.89
2	B	1801	Z3J	C14-C13-C11	-2.22	117.41	120.22
2	A	1801	Z3J	C14-N15-C16	-2.21	105.25	113.67
2	A	1801	Z3J	C08-C23-C13	-2.09	115.91	118.92

There are no chirality outliers.


All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1801	Z3J	C05-C03-N02-C01
2	A	1801	Z3J	O04-C03-N02-C01
2	B	1801	Z3J	C05-C03-N02-C01
2	B	1801	Z3J	O04-C03-N02-C01
2	B	1801	Z3J	N02-C03-C05-C06
2	B	1801	Z3J	O04-C03-C05-C06
2	B	1801	Z3J	O04-C03-C05-C23
2	B	1801	Z3J	N02-C03-C05-C23

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	267/286 (93%)	1.09	37 (13%) 6 7	32, 47, 70, 95	0
1	B	272/286 (95%)	1.28	45 (16%) 4 5	36, 53, 77, 100	0
All	All	539/572 (94%)	1.18	82 (15%) 5 6	32, 49, 76, 100	0

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1568	ILE	4.1
1	A	1629	ILE	4.1
1	B	1524	ASP	3.9
1	B	1622	GLY	3.9
1	A	1595	PRO	3.7
1	B	1681	GLY	3.5
1	B	1593	THR	3.4
1	B	1467	GLY	3.1
1	B	1724	ALA	3.1
1	B	1647	GLN	3.1
1	B	1594	ILE	3.1
1	A	1626	PRO	3.1
1	A	1596	ALA	3.0
1	A	1591	ASN	3.0
1	A	1590	PHE	3.0
1	B	1452	ILE	3.0
1	A	1666	ASP	3.0
1	A	1625	ILE	2.9
1	B	1464	GLU	2.9
1	A	1585	PHE	2.9
1	A	1630	ILE	2.9
1	A	1615	LEU	2.8
1	B	1465	ALA	2.8
1	B	1665	ASP	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	1523	GLY	2.8
1	B	1597	ILE	2.8
1	A	1478	GLY	2.8
1	A	1619	SER	2.8
1	A	1620	GLN	2.7
1	A	1597	ILE	2.7
1	A	1617	ALA	2.7
1	B	1453	ASP	2.7
1	B	1463	PRO	2.7
1	A	1621	SER	2.7
1	A	1664	HIS	2.6
1	B	1725	ASP	2.6
1	B	1466	GLY	2.5
1	A	1704	ASP	2.5
1	B	1586	ALA	2.5
1	B	1679	PRO	2.5
1	B	1599	TYR	2.5
1	B	1694	PRO	2.5
1	A	1669	MET	2.4
1	B	1602	LEU	2.4
1	B	1525	GLY	2.4
1	A	1570	GLN	2.4
1	A	1725	ASP	2.4
1	B	1589	THR	2.4
1	B	1459	LEU	2.4
1	B	1488	LEU	2.4
1	A	1665	ASP	2.3
1	B	1618	VAL	2.3
1	B	1596	ALA	2.3
1	B	1643	ILE	2.3
1	B	1623	VAL	2.3
1	B	1587	GLU	2.3
1	B	1661	ASP	2.3
1	B	1461	ALA	2.2
1	B	1652	ASP	2.2
1	A	1618	VAL	2.2
1	A	1717	ARG	2.2
1	B	1489	GLY	2.2
1	A	1503	VAL	2.2
1	A	1661	ASP	2.2
1	B	1669	MET	2.2
1	A	1504	GLU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	1601	GLN	2.2
1	B	1686	TYR	2.1
1	A	1614	VAL	2.1
1	A	1627	ALA	2.1
1	A	1479	GLY	2.1
1	A	1508	GLU	2.1
1	A	1524	ASP	2.1
1	A	1509	GLU	2.1
1	A	1610	GLN	2.1
1	B	1516	PRO	2.1
1	B	1554	ARG	2.1
1	A	1688	SER	2.0
1	B	1635	THR	2.0
1	B	1582	TYR	2.0
1	B	1628	GLY	2.0
1	A	1576	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	Z3J	B	1801	30/30	0.83	0.15	55,68,78,79	0
2	Z3J	A	1801	30/30	0.89	0.11	39,45,52,54	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Z3J B 1801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Z3J A 1801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.