



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 18, 2026 – 06:18 AM UTC

PDB ID : 3M8L / pdb_00003m8l
Title : Crystal Structure Analysis of the Feline Calicivirus Capsid Protein
Authors : Zhou, Y.; Prasad, B.V.V.
Deposited on : 2010-03-18
Resolution : 3.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1001 (3.44-3.36)
Clashscore	190562	1022 (3.44-3.36)
Ramachandran outliers	187476	1012 (3.44-3.36)
Sidechain outliers	187428	1012 (3.44-3.36)
RSRZ outliers	180081	1001 (3.44-3.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	534	35% 22% 51% 22% 5%
1	B	534	37% 19% 52% 24% 5%
1	C	534	37% 20% 54% 21% 5%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 12360 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Capsid protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	534	Total	C	N	O	S	0	0	0
			4129	2632	687	796	14			
1	B	530	Total	C	N	O	S	0	0	0
			4101	2615	683	789	14			
1	C	534	Total	C	N	O	S	0	0	0
			4130	2632	687	797	14			


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Capsid protein





• Molecule 1: Capsid protein





• Molecule 1: Capsid protein





4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	427.08Å 450.73Å 467.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 3.40 30.00 – 3.40	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-3.40) 88.9 (30.00-3.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.28 (at 3.39Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.390 , 0.370 0.420 , 0.419	Depositor DCC
R_{free} test set	27284 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	85.6	Xtriage
Anisotropy	0.218	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 0.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.017 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.39	EDS
Total number of atoms	12360	wwPDB-VP
Average B, all atoms (Å ²)	96.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.87	7/4229 (0.2%)	1.21	42/5759 (0.7%)
1	B	1.16	15/4201 (0.4%)	1.21	40/5719 (0.7%)
1	C	0.91	11/4230 (0.3%)	1.28	51/5759 (0.9%)
All	All	0.99	33/12660 (0.3%)	1.23	133/17237 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

The worst 5 of 33 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	624	ASP	CB-CG	28.32	2.22	1.52
1	B	606	PHE	CE1-CZ	15.16	1.84	1.38
1	B	606	PHE	CE2-CZ	13.81	1.80	1.38
1	B	606	PHE	CD2-CE2	13.68	1.79	1.38
1	B	606	PHE	CD1-CE1	13.66	1.79	1.38

The worst 5 of 133 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	438	THR	CB-CA-C	18.35	146.94	110.42
1	A	439	ASN	N-CA-CB	-13.86	87.07	110.49
1	C	512	GLN	N-CA-C	11.70	123.19	108.45
1	B	360	ASP	N-CA-C	10.88	133.98	110.80
1	C	446	GLN	N-CA-C	-10.75	99.83	113.16

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	200	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4129	0	4062	636	0
1	B	4101	0	4033	645	6
1	C	4130	0	4062	598	7
All	All	12360	0	12157	1829	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 75.

The worst 5 of 1829 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:606:PHE:CE1	1:B:606:PHE:CZ	1.84	1.62
1:B:606:PHE:CE1	1:B:606:PHE:CD1	1.79	1.62
1:B:606:PHE:CE2	1:B:606:PHE:CD2	1.79	1.62
1:B:606:PHE:CZ	1:B:606:PHE:CE2	1.80	1.61
1:B:606:PHE:CZ	1:B:624:ASP:CG	1.87	1.52

The worst 5 of 7 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:299:TYR:CE1	1:C:299:TYR:OH[2_555]	1.77	0.43
1:B:299:TYR:CD1	1:C:299:TYR:CZ[2_555]	1.97	0.23
1:B:229:GLY:N	1:C:222:SER:OG[2_555]	2.06	0.14
1:B:292:TYR:OH	1:C:313:GLU:OE1[2_555]	2.07	0.13
1:B:252:PRO:N	1:C:169:PHE:CE1[2_555]	2.10	0.10

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	532/534 (100%)	310 (58%)	147 (28%)	75 (14%)	0	0
1	B	528/534 (99%)	321 (61%)	132 (25%)	75 (14%)	0	0
1	C	532/534 (100%)	332 (62%)	126 (24%)	74 (14%)	0	1
All	All	1592/1602 (99%)	963 (60%)	405 (25%)	224 (14%)	0	0

5 of 224 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	133	PRO
1	A	145	GLU
1	A	146	PRO
1	A	151	SER
1	A	279	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	461/461 (100%)	370 (80%)	91 (20%)	1	5
1	B	457/461 (99%)	350 (77%)	107 (23%)	1	2
1	C	461/461 (100%)	376 (82%)	85 (18%)	1	7
All	All	1379/1383 (100%)	1096 (80%)	283 (20%)	1	4

5 of 283 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	358	ILE
1	C	402	MET
1	C	523	SER
1	B	165	GLU
1	B	149	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 65 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	505	GLN
1	C	521	GLN
1	A	593	GLN
1	A	584	ASN
1	C	569	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	534/534 (100%)	1.79	189 (35%) 1 1	48, 97, 120, 143	0
1	B	530/534 (99%)	1.72	197 (37%) 1 1	48, 97, 112, 129	0
1	C	534/534 (100%)	1.77	195 (36%) 1 1	49, 95, 113, 123	0
All	All	1598/1602 (99%)	1.76	581 (36%) 1 1	48, 96, 115, 143	0

The worst 5 of 581 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	301	ASN	10.1
1	C	604	ASP	9.6
1	C	602	SER	9.3
1	A	637	GLY	8.5
1	C	513	ASP	7.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.