



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:43 AM UTC

PDB ID : 5M92 / pdb_00005m92
Title : PCE reductive dehalogenase from *S. multivorans* in complex with 2,4-dibromophenol
Authors : Kunze, C.; Bommer, M.; Hagen, W.R.; Uksa, M.; Dobbek, H.; Schubert, T.; Diekert, G.
Deposited on : 2016-10-31
Resolution : 1.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1365 (1.78-1.78)
Clashscore	190562	1395 (1.78-1.78)
Ramachandran outliers	187476	1382 (1.78-1.78)
Sidechain outliers	187428	1382 (1.78-1.78)
RSRZ outliers	180081	1365 (1.78-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	464	 90% 5% 5%
1	B	464	 86% 10% 4% 2%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	BVQ	A	503	X	-	-	-
3	BVQ	B	503	X	-	-	-

2 Entry composition [i](#)


There are 8 unique types of molecules in this entry. The entry contains 7904 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tetrachloroethene reductive dehalogenase catalytic subunit PceA.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	443	Total	C	N	O	S	0	5	0
			3545	2251	608	654	32			
1	B	419	Total	C	N	O	S	0	2	0
			3312	2100	565	617	30			

- Molecule 2 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe_4S_4).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	Fe	S	0	0
			8	4	4		
2	A	1	Total	Fe	S	0	0
			8	4	4		
2	B	1	Total	Fe	S	0	0
			8	4	4		
2	B	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 3 is NORPSEUDO-B12 (CCD ID: BVQ) (formula: $C_{57}H_{82}CoN_{16}O_{14}P$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total 89	C 57	Co 1	N 16	O 14	P 1	0	0
3	B	1	Total 89	C 57	Co 1	N 16	O 14	P 1	0	0

- Molecule 4 is BENZAMIDINE (CCD ID: BEN) (formula: $C_7H_8N_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	N		
4	A	1	Total 9	7	2	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			9	7	2		
4	B	1	Total	C	N	0	0
			9	7	2		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is 2,4-bis(bromanyl)phenol (CCD ID: Y8I) (formula: $C_6H_4Br_2O$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	Br	C	O	0	0
			9	2	6	1		
6	A	1	Total	Br	C	O	0	0
			9	2	6	1		
6	B	1	Total	Br	C	O	0	0
			9	2	6	1		
6	B	1	Total	Br	C	O	0	0
			9	2	6	1		

- Molecule 7 is BROMIDE ION (CCD ID: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Br	0	0
			1	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	423	Total	O	0	0
			423	423		
8	B	308	Total	O	0	0
			308	308		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tetrachloroethene reductive dehalogenase catalytic subunit PceA



- Molecule 1: Tetrachloroethene reductive dehalogenase catalytic subunit PceA



4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	73.94Å 73.94Å 185.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.38 – 1.78 47.38 – 1.78	Depositor EDS
% Data completeness (in resolution range)	99.4 (47.38-1.78) 99.5 (47.38-1.78)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.41 (at 1.78Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.149 , 0.178 0.150 , 0.180	Depositor DCC
R_{free} test set	1059 reflections (1.13%)	wwPDB-VP
Wilson B-factor (Å ²)	24.2	Xtriage
Anisotropy	0.021	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 51.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	0.065 for h,-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7904	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BEN, Y8I, BVQ, SF4, BR, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/3635	0.52	0/4931
1	B	0.32	0/3396	0.51	0/4610
All	All	0.32	0/7031	0.51	0/9541

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3545	0	3442	28	0
1	B	3312	0	3206	25	0
2	A	16	0	0	0	0
2	B	16	0	0	1	0
3	A	89	0	82	7	0
3	B	89	0	82	3	0
4	A	18	0	14	2	0
4	B	9	0	7	1	0
5	A	36	0	48	1	0
5	B	6	0	8	0	0
6	A	18	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	18	0	0	1	0
7	A	1	0	0	0	0
8	A	423	0	0	5	0
8	B	308	0	0	1	0
All	All	7904	0	6889	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:126:ALA:O	1:A:132:ARG:NH2	1.98	0.96
1:A:13:GLN:HE22	1:A:405:TRP:H	1.34	0.74
1:B:225:MET:HE2	1:B:241:THR:HG23	1.70	0.74
1:A:447:ALA:HB1	1:B:451:ARG:HG2	1.70	0.73
1:A:5:LYS:NZ	8:A:602:HOH:O	2.18	0.72
1:A:5:LYS:HA	1:A:405:TRP:CH2	2.26	0.71
1:B:225:MET:HE3	1:B:242:THR:HA	1.72	0.71
3:B:503:BVQ:H35	3:B:503:BVQ:H36A	1.74	0.69
3:A:503:BVQ:H35	3:A:503:BVQ:H36A	1.76	0.67
1:B:251:MET:HG3	1:B:255:TRP:CE3	2.33	0.64
1:A:25:ASN:HD21	1:A:27:LYS:HE2	1.61	0.64
1:B:451:ARG:HG3	8:B:711:HOH:O	1.99	0.63
3:A:503:BVQ:H53	3:A:503:BVQ:H55A	1.80	0.63
1:A:330[A]:GLU:OE1	8:A:601:HOH:O	2.16	0.62
3:B:503:BVQ:H55A	3:B:503:BVQ:H53	1.81	0.62
1:A:98:LEU:HD22	1:B:251:MET:CE	2.30	0.61
1:B:225:MET:HE3	1:B:242:THR:CA	2.32	0.60
1:A:25:ASN:OD1	1:A:27:LYS:HG2	2.02	0.60
1:A:98:LEU:HD13	1:B:251:MET:HE1	1.84	0.59
1:B:102:TYR:HB3	6:B:506:Y8I:C09	2.32	0.59
1:B:251:MET:HG3	1:B:255:TRP:CZ3	2.39	0.58
1:A:102:TYR:HB3	6:A:512:Y8I:C09	2.34	0.58
1:A:4:GLU:OE1	1:A:5:LYS:N	2.35	0.57
1:A:451:ARG:NE	8:A:603:HOH:O	2.37	0.57
1:A:5:LYS:HA	1:A:405:TRP:HH2	1.69	0.55
1:A:73:ASN:HB3	1:A:432:ILE:HD12	1.90	0.53
1:B:73:ASN:HB3	1:B:432:ILE:HD12	1.92	0.52
1:A:189:GLU:OE1	4:A:504:BEN:H4	2.11	0.51
1:A:80:LYS:NZ	8:A:604:HOH:O	2.38	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:191[B]:GLU:OE2	4:B:505:BEN:N1	2.45	0.50
1:B:225:MET:HE3	1:B:242:THR:N	2.27	0.50
1:A:442:THR:HG22	1:A:443:TYR:CD2	2.47	0.49
1:B:225:MET:CE	1:B:241:THR:HG23	2.43	0.48
1:A:98:LEU:HD22	1:B:251:MET:HE1	1.95	0.48
1:A:462:LYS:HD2	8:A:848:HOH:O	2.14	0.47
1:B:394:LYS:HA	1:B:394:LYS:HD2	1.63	0.47
1:A:261:ARG:HA	1:A:265:TYR:O	2.14	0.47
1:B:383:CYS:HB3	2:B:502:SF4:S4	2.55	0.46
1:B:287:GLY:HA2	1:B:321:ILE:HD11	1.95	0.46
1:A:447:ALA:CB	1:B:451:ARG:HG2	2.44	0.45
1:B:148:LYS:O	1:B:152:ARG:HG2	2.17	0.45
4:A:504:BEN:H4	5:A:505:GOL:O3	2.16	0.44
3:A:503:BVQ:H53	3:A:503:BVQ:C55	2.47	0.44
1:A:126:ALA:HA	1:A:132:ARG:HH22	1.84	0.43
1:B:225:MET:HE1	1:B:242:THR:OG1	2.19	0.43
3:A:503:BVQ:H26A	3:A:503:BVQ:H19	1.87	0.42
1:A:293:GLY:O	3:A:503:BVQ:H1R	2.20	0.42
3:B:503:BVQ:H35	3:B:503:BVQ:C36	2.46	0.42
1:A:98:LEU:HD22	1:B:251:MET:HE3	1.98	0.42
1:B:261:ARG:HA	1:B:265:TYR:O	2.19	0.42
1:A:38:PHE:CD2	3:A:503:BVQ:H53A	2.55	0.42
3:A:503:BVQ:H35	3:A:503:BVQ:C36	2.47	0.42
1:A:251[B]:MET:SD	1:B:251:MET:HG2	2.60	0.41
1:A:126:ALA:CA	1:A:132:ARG:HH22	2.34	0.41
1:B:173:ALA:HB2	1:B:194:ILE:HD11	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	444/464 (96%)	434 (98%)	10 (2%)	0	100	100
1	B	417/464 (90%)	408 (98%)	9 (2%)	0	100	100
All	All	861/928 (93%)	842 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	379/389 (97%)	379 (100%)	0	100	100
1	B	354/389 (91%)	354 (100%)	0	100	100
All	All	733/778 (94%)	733 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	13	GLN
1	B	123	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 21 ligands modelled in this entry, 1 is monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	BEN	A	504	-	9,9,9	1.40	1 (11%)	7,11,11	1.00	0
4	BEN	A	511	-	9,9,9	1.28	1 (11%)	7,11,11	0.83	0
5	GOL	A	509	-	5,5,5	0.38	0	5,5,5	0.30	0
6	Y8I	A	513	-	9,9,9	2.78	4 (44%)	12,12,12	2.67	4 (33%)
5	GOL	A	507	-	5,5,5	0.36	0	5,5,5	0.58	0
6	Y8I	B	506	-	9,9,9	2.01	1 (11%)	12,12,12	3.06	3 (25%)
3	BVQ	B	503	8	92,99,99	0.99	5 (5%)	140,162,162	1.35	16 (11%)
4	BEN	B	505	-	9,9,9	1.29	1 (11%)	7,11,11	0.90	0
5	GOL	A	506	-	5,5,5	0.35	0	5,5,5	0.58	0
5	GOL	A	505	-	5,5,5	0.33	0	5,5,5	0.45	0
5	GOL	B	504	-	5,5,5	0.36	0	5,5,5	0.58	0
3	BVQ	A	503	8	92,99,99	1.00	4 (4%)	140,162,162	1.38	18 (12%)
6	Y8I	B	507	-	9,9,9	2.53	4 (44%)	12,12,12	2.74	4 (33%)
2	SF4	A	502	1	0,12,12	-	-	-	-	-
2	SF4	A	501	1	0,12,12	-	-	-	-	-
2	SF4	B	502	1	0,12,12	-	-	-	-	-
2	SF4	B	501	1	0,12,12	-	-	-	-	-
5	GOL	A	510	-	5,5,5	0.29	0	5,5,5	0.71	0
5	GOL	A	508	-	5,5,5	0.36	0	5,5,5	0.30	0
6	Y8I	A	512	-	9,9,9	2.03	2 (22%)	12,12,12	3.06	4 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BEN	A	504	-	-	4/4/4/4	0/1/1/1
4	BEN	A	511	-	-	0/4/4/4	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	509	-	-	4/4/4/4	-
6	Y8I	A	513	-	-	-	0/1/1/1
5	GOL	A	507	-	-	0/4/4/4	-
6	Y8I	B	506	-	-	-	0/1/1/1
3	BVQ	B	503	8	4/4/35/37	2/54/221/221	0/3/11/11
4	BEN	B	505	-	-	0/4/4/4	0/1/1/1
5	GOL	A	506	-	-	0/4/4/4	-
5	GOL	A	505	-	-	0/4/4/4	-
5	GOL	B	504	-	-	4/4/4/4	-
3	BVQ	A	503	8	4/4/35/37	2/54/221/221	0/3/11/11
6	Y8I	B	507	-	-	-	0/1/1/1
2	SF4	A	502	1	-	-	0/6/5/5
2	SF4	A	501	1	-	-	0/6/5/5
2	SF4	B	502	1	-	-	0/6/5/5
2	SF4	B	501	1	-	-	0/6/5/5
5	GOL	A	510	-	-	2/4/4/4	-
5	GOL	A	508	-	-	0/4/4/4	-
6	Y8I	A	512	-	-	-	0/1/1/1

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	503	BVQ	C14-N23	6.22	1.43	1.35
3	B	503	BVQ	C14-N23	5.66	1.42	1.35
6	B	507	Y8I	O07-C06	5.11	1.46	1.36
6	A	513	Y8I	O07-C06	5.06	1.46	1.36
6	B	506	Y8I	O07-C06	4.83	1.46	1.36
6	A	512	Y8I	O07-C06	4.79	1.46	1.36
6	A	513	Y8I	BR1-C02	4.49	1.99	1.90
4	A	504	BEN	C1-C	-3.90	1.40	1.47
3	B	503	BVQ	C13-C14	3.64	1.59	1.52
4	B	505	BEN	C1-C	-3.47	1.40	1.47
4	A	511	BEN	C1-C	-3.42	1.40	1.47
3	A	503	BVQ	C13-C14	3.32	1.58	1.52
6	B	507	Y8I	BR1-C02	3.28	1.96	1.90
3	B	503	BVQ	C2B-N1B	-3.00	1.32	1.37
3	A	503	BVQ	C2B-N1B	-2.84	1.32	1.37
6	B	507	Y8I	C09-C08	2.70	1.43	1.38
6	A	513	Y8I	C09-C08	2.49	1.42	1.38
3	B	503	BVQ	C16-C15	2.39	1.51	1.44
6	B	507	Y8I	C09-C02	2.33	1.42	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	513	Y8I	C09-C02	2.27	1.42	1.38
6	A	512	Y8I	C09-C02	2.17	1.42	1.38
3	B	503	BVQ	C2B-N3B	2.12	1.35	1.31
3	A	503	BVQ	C2B-N3B	2.05	1.35	1.31

All (49) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	512	Y8I	C08-C06-C04	9.01	124.85	118.07
6	A	513	Y8I	C08-C06-C04	8.05	124.13	118.07
6	B	506	Y8I	C08-C06-C04	7.64	123.82	118.07
6	B	507	Y8I	C08-C06-C04	7.30	123.57	118.07
6	B	506	Y8I	BR5-C04-C06	6.37	124.26	119.38
3	B	503	BVQ	C8B-N1B-C1R	6.01	136.12	125.48
3	A	503	BVQ	C8B-N1B-C1R	5.90	135.92	125.48
3	B	503	BVQ	C8B-C9B-N3B	-4.91	105.78	110.45
6	B	507	Y8I	BR5-C04-C06	4.70	122.98	119.38
3	A	503	BVQ	C8B-C9B-N3B	-4.41	106.26	110.45
3	B	503	BVQ	C9B-N3B-C2B	4.35	106.60	103.85
3	A	503	BVQ	C9B-N3B-C2B	3.88	106.30	103.85
3	B	503	BVQ	C48-C13-C14	3.82	118.01	108.51
3	A	503	BVQ	C48-C13-C14	3.64	117.55	108.51
3	A	503	BVQ	C10-C9-N22	3.50	129.73	125.74
6	A	512	Y8I	BR5-C04-C06	3.40	121.99	119.38
3	A	503	BVQ	C9-N22-C6	3.33	109.28	105.28
3	B	503	BVQ	C1R-N1B-C2B	-3.31	119.74	127.09
6	A	512	Y8I	C09-C08-C06	-3.31	117.19	120.50
3	A	503	BVQ	C1R-N1B-C2B	-3.12	120.16	127.09
3	A	503	BVQ	C8B-N1B-C2B	-3.09	103.47	105.33
3	B	503	BVQ	C13-C12-C11	2.93	104.25	100.97
3	A	503	BVQ	C12-C11-N23	2.78	115.65	111.83
3	B	503	BVQ	C8B-N1B-C2B	-2.75	103.67	105.33
6	B	506	Y8I	C09-C08-C06	-2.75	117.75	120.50
3	B	503	BVQ	C36-C7-C8	-2.74	106.99	112.05
3	A	503	BVQ	C35-C5-C6	2.73	126.80	122.41
3	A	503	BVQ	C35-C5-C4	-2.71	111.31	116.79
3	B	503	BVQ	C5-C6-N22	-2.58	119.98	123.88
6	A	513	Y8I	C09-C08-C06	-2.54	117.96	120.50
3	B	503	BVQ	C19-N24-C16	2.53	110.06	107.29
3	A	503	BVQ	C15-C16-N24	-2.50	118.87	122.42
3	A	503	BVQ	C36-C7-C8	-2.46	107.51	112.05
3	B	503	BVQ	C9-N22-C6	2.43	108.20	105.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	507	Y8I	C03-C04-C06	-2.38	119.13	120.95
3	B	503	BVQ	C15-C16-N24	-2.34	119.08	122.42
6	A	513	Y8I	C03-C04-C06	-2.32	119.18	120.95
6	A	512	Y8I	C03-C04-C06	-2.29	119.20	120.95
3	B	503	BVQ	C12-C11-C10	-2.26	120.48	123.40
3	A	503	BVQ	C30-C3-C2	-2.26	114.00	119.00
3	B	503	BVQ	C41-C8-C9	-2.25	107.26	111.19
6	B	507	Y8I	C09-C08-C06	-2.23	118.27	120.50
6	A	513	Y8I	BR5-C04-C06	2.20	121.06	119.38
3	A	503	BVQ	C5-C6-N22	-2.13	120.65	123.88
3	A	503	BVQ	C12-C11-C10	-2.11	120.67	123.40
3	B	503	BVQ	C41-C8-C7	-2.10	108.41	114.19
3	A	503	BVQ	C53-C15-C16	-2.09	116.82	120.36
3	B	503	BVQ	C30-C3-C2	-2.07	114.43	119.00
3	A	503	BVQ	C9-C10-C11	-2.01	123.10	125.97

All (8) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	503	BVQ	N24
3	A	503	BVQ	N21
3	A	503	BVQ	N22
3	A	503	BVQ	N23
3	B	503	BVQ	N24
3	B	503	BVQ	N21
3	B	503	BVQ	N22
3	B	503	BVQ	N23

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	503	BVQ	C14-C13-C48-C49
5	A	509	GOL	O1-C1-C2-C3
5	A	509	GOL	C1-C2-C3-O3
5	B	504	GOL	O1-C1-C2-C3
5	B	504	GOL	C1-C2-C3-O3
3	A	503	BVQ	C14-C13-C48-C49
5	A	510	GOL	C1-C2-C3-O3
5	A	510	GOL	O2-C2-C3-O3
5	B	504	GOL	O1-C1-C2-O2
5	B	504	GOL	O2-C2-C3-O3
5	A	509	GOL	O1-C1-C2-O2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	A	509	GOL	O2-C2-C3-O3
3	A	503	BVQ	C38-C37-C7-C6
3	B	503	BVQ	C38-C37-C7-C6
4	A	504	BEN	N2-C-C1-C2
4	A	504	BEN	N2-C-C1-C6
4	A	504	BEN	N1-C-C1-C2
4	A	504	BEN	N1-C-C1-C6

There are no ring outliers.


8 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	504	BEN	2	0
6	B	506	Y8I	1	0
3	B	503	BVQ	3	0
4	B	505	BEN	1	0
5	A	505	GOL	1	0
3	A	503	BVQ	7	0
2	B	502	SF4	1	0
6	A	512	Y8I	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






Ligand Y8I B 506



Ligand BVQ B 503







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	443/464 (95%)	-0.45	1 (0%) 91 93	11, 26, 45, 78	5 (1%)
1	B	419/464 (90%)	-0.33	4 (0%) 79 85	13, 27, 52, 84	2 (0%)
All	All	862/928 (92%)	-0.39	5 (0%) 85 89	11, 27, 48, 84	7 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	10	ILE	3.6
1	B	7	ALA	3.0
1	A	396	ASN	2.4
1	B	461	VAL	2.2
1	B	394	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	BEN	A	511	9/9	0.82	0.15	53,55,63,65	0
5	GOL	A	509	6/6	0.82	0.17	60,60,64,65	0
5	GOL	A	508	6/6	0.84	0.14	49,53,57,59	0
4	BEN	B	505	9/9	0.85	0.14	43,44,50,57	0
4	BEN	A	504	9/9	0.87	0.17	51,57,59,61	0
5	GOL	B	504	6/6	0.88	0.14	44,55,59,63	0
5	GOL	A	505	6/6	0.89	0.14	46,52,59,60	0
5	GOL	A	507	6/6	0.91	0.12	34,39,54,55	0
5	GOL	A	510	6/6	0.93	0.09	32,38,40,40	0
7	BR	A	514	1/1	0.93	0.19	44,44,44,44	1
5	GOL	A	506	6/6	0.95	0.08	27,32,39,39	0
6	Y8I	B	507	9/9	0.96	0.14	27,37,43,52	9
6	Y8I	A	513	9/9	0.96	0.13	25,33,45,47	9
6	Y8I	A	512	9/9	0.97	0.11	23,26,32,35	9
3	BVQ	B	503	89/89	0.98	0.05	13,22,30,42	0
6	Y8I	B	506	9/9	0.98	0.10	19,24,31,33	9
2	SF4	B	501	8/8	0.98	0.05	29,30,32,33	0
2	SF4	B	502	8/8	0.98	0.04	23,24,25,26	0
2	SF4	A	502	8/8	0.99	0.03	18,21,22,22	0
3	BVQ	A	503	89/89	0.99	0.05	12,20,28,35	0
2	SF4	A	501	8/8	0.99	0.03	19,21,22,22	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Y8I B 507:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Y8I A 513:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Y8I A 512:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BVQ B 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Y8I B 506:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around BVQ A 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.