



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 08:02 AM UTC

PDB ID : 6M96 / pdb_00006m96
Title : ATP-bound conformation of the WzmWzt O antigen ABC transporter
Authors : Caffalette, C.A.; Zimmer, J.
Deposited on : 2018-08-22
Resolution : 2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	244	4% 82% 16% .
2	B	256	7% 82% 17% .

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 4982 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ABC transporter.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	238	Total	C	N	O	S	0	10	0
			1949	1267	319	356	7			


There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP O67181
A	1	GLY	-	expression tag	UNP O67181
A	167	GLN	GLU	engineered mutation	UNP O67181
A	236	LYS	-	expression tag	UNP O67181
A	237	LEU	-	expression tag	UNP O67181
A	238	HIS	-	expression tag	UNP O67181
A	239	HIS	-	expression tag	UNP O67181
A	240	HIS	-	expression tag	UNP O67181
A	241	HIS	-	expression tag	UNP O67181
A	242	HIS	-	expression tag	UNP O67181
A	243	HIS	-	expression tag	UNP O67181

- Molecule 2 is a protein called Transport permease protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	256	Total	C	N	O	S	0	27	0
			2281	1613	313	350	5			

- Molecule 3 is LAURYL DIMETHYLAMINE-N-OXIDE (CCD ID: LDA) (formula: C₁₄H₃₁NO).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C			0	0
			4	4				
3	B	1	Total	C	N	O	0	0
			13	11	1	1		
3	B	1	Total	C	N	O	0	0
			16	14	1	1		
3	B	1	Total	C	N	O	0	0
			16	14	1	1		
3	B	1	Total	C	N	O	0	0
			16	14	1	1		
3	B	1	Total	C			0	0
			8	8				
3	B	1	Total	C	N	O	0	0
			16	14	1	1		
3	B	1	Total	C	N	O	0	0
			8	6	1	1		
3	B	1	Total	C			0	0
			6	6				
3	B	1	Total	C			0	0
			6	6				
3	B	1	Total	C			0	0
			4	4				
3	B	1	Total	C			0	0
			6	6				
3	B	1	Total	C	N	O	0	0
			12	10	1	1		
3	B	1	Total	C			0	0
			10	10				

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			9	7	1	1		
3	B	1	Total	C	N	O	0	0
			16	14	1	1		
3	B	1	Total	C			0	0
			8	8				
3	B	1	Total	C			0	0
			4	4				
3	B	1	Total	C			0	0
			5	5				
3	B	1	Total	C			0	0
			3	3				
3	B	1	Total	C			0	0
			3	3				
3	B	1	Total	C			0	0
			5	5				
3	B	1	Total	C			0	0
			8	8				
3	B	1	Total	C			0	0
			9	9				
3	B	1	Total	C			0	0
			5	5				
3	B	1	Total	C			0	0
			3	3				
3	B	1	Total	C			0	0
			5	5				
3	B	1	Total	C			0	0
			3	3				
3	B	1	Total	C			0	0
			4	4				
3	B	1	Total	C	N	O	0	0
			16	14	1	1		
3	B	1	Total	C			0	0
			4	4				
3	B	1	Total	C	N	O	0	0
			5	3	1	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).




- PG4
-
- The diagram shows the chemical structure of PG4, a diphosphatidylglycerol. It consists of two identical phosphatidyl groups linked by a central glycerol backbone. Each phosphatidyl group is composed of a glycerol moiety (C1, C2, C3) and a phosphate group (O2, O3). The phosphate groups are linked to a second glycerol moiety (C4, C5, C6) which is also linked to a third glycerol moiety (C7, C8) ending in a hydroxyl group (OH, O5). The structure is shown in a zig-zag conformation, with carbon atoms labeled C1 through C8 and oxygen atoms labeled O1 through O5. The hydroxyl groups are shown in red, and the phosphate groups are shown in red. The carbon atoms are shown in green.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 13 8 5	0	0
6	A	1	Total C O 10 6 4	0	0


- 

C₄₁H₇₈NO₈P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	A	1	Total	C	N	O	P	0	0
			9	3	1	4	1		


- Molecule 8 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 9 is 1-ETHOXY-2-(2-ETHOXYETHOXY)ETHANE (CCD ID: P4G) (formula:

C₈H₁₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	C	O	0	0
			9	6	3		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	232	Total	O	0	0
			232	232		
10	B	139	Total	O	0	0
			139	139		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ABC transporter



- Molecule 2: Transport permease protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	144.75Å 144.75Å 201.47Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	24.88 – 2.05 24.88 – 2.05	Depositor EDS
% Data completeness (in resolution range)	100.0 (24.88-2.05) 99.9 (24.88-2.05)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.70 (at 2.04Å)	Xtriage
Refinement program	PHENIX (1.13_2998)	Depositor
R, R_{free}	0.164 , 0.182 0.165 , 0.183	Depositor DCC
R_{free} test set	4012 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	38.8	Xtriage
Anisotropy	0.027	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 83.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4982	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PG4, SO4, LDA, ATP, P4G, PEE, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/2015	0.52	0/2707
2	B	0.30	0/2444	0.44	0/3334
All	All	0.34	0/4459	0.48	0/6041

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1949	0	2036	37	0
2	B	2281	0	2470	45	0
3	A	4	0	7	0	0
3	B	252	0	451	15	0
4	A	36	0	46	1	0
4	B	12	0	16	0	0
5	A	31	0	12	0	0
6	A	23	0	31	2	0
7	A	9	0	7	1	0
8	B	5	0	0	0	0
9	B	9	0	13	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	A	232	0	0	11	9
10	B	139	0	0	3	6
All	All	4982	0	5089	83	10

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 83 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:13:TYR:O	10:A:401:HOH:O	1.95	0.84
2:B:109[A]:SER:OG	10:B:401:HOH:O	1.97	0.83
1:A:65:LYS:HB3	1:A:71[B]:THR:HG23	1.61	0.81
1:A:215:HIS:O	10:A:402:HOH:O	2.02	0.77
1:A:65:LYS:NZ	10:A:405:HOH:O	2.18	0.77

The worst 5 of 10 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
10:A:480:HOH:O	10:A:480:HOH:O[12_566]	1.87	0.33
10:A:608:HOH:O	10:A:608:HOH:O[10_667]	1.99	0.21
10:B:492:HOH:O	10:B:492:HOH:O[10_667]	1.99	0.21
10:A:450:HOH:O	10:B:508:HOH:O[8_567]	2.04	0.16
10:A:504:HOH:O	10:A:504:HOH:O[12_566]	2.07	0.13

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	246/244 (101%)	239 (97%)	6 (2%)	1 (0%)	30 22

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	282/256 (110%)	278 (99%)	4 (1%)	0	100	100
All	All	528/500 (106%)	517 (98%)	10 (2%)	1 (0%)	43	37

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	6	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	217/213 (102%)	217 (100%)	0	100	100
2	B	265/237 (112%)	260 (98%)	5 (2%)	50	50
All	All	482/450 (107%)	477 (99%)	5 (1%)	76	72

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	31[A]	TRP
2	B	31[B]	TRP
2	B	89	LYS
2	B	217[A]	VAL
2	B	217[B]	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

46 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	LDA	B	330	-	3,3,15	0.39	0	2,2,17	0.61	0
3	LDA	B	317	-	3,3,15	0.38	0	2,2,17	0.63	0
3	LDA	B	308	-	5,5,15	0.31	0	4,4,17	0.33	0
4	GOL	A	307	-	5,5,5	0.98	0	5,5,5	1.09	0
4	GOL	A	303	-	5,5,5	1.01	0	5,5,5	1.11	0
5	ATP	A	308	-	32,33,33	1.55	7 (21%)	48,52,52	1.71	8 (16%)
3	LDA	B	329	-	13,15,15	2.34	2 (15%)	14,17,17	0.45	0
3	LDA	B	314	-	6,8,15	3.58	2 (33%)	7,10,17	0.31	0
4	GOL	A	306	-	5,5,5	0.71	0	5,5,5	1.19	1 (20%)
3	LDA	B	307	-	5,7,15	3.99	2 (40%)	6,9,17	0.65	0
3	LDA	B	319	-	2,2,15	0.25	0	1,1,17	0.01	0
3	LDA	B	320	-	2,2,15	0.19	0	1,1,17	0.12	0
3	LDA	B	327	-	2,2,15	0.25	0	1,1,17	0.03	0
4	GOL	B	333	-	5,5,5	0.91	0	5,5,5	1.25	1 (20%)
3	LDA	B	326	-	3,3,15	0.28	0	1,1,17	0.03	0
3	LDA	B	315	-	13,15,15	2.58	2 (15%)	14,17,17	0.56	0
3	LDA	B	311	-	5,5,15	0.28	0	4,4,17	0.41	0
3	LDA	B	331	-	4,4,15	6.39	2 (50%)	6,6,17	0.26	0
3	LDA	B	303	-	13,15,15	2.36	2 (15%)	14,17,17	0.66	0
3	LDA	B	318	-	4,4,15	0.37	0	3,3,17	0.31	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	PG4	A	309	-	12,12,12	0.54	0	11,11,11	0.28	0
3	LDA	B	324	-	4,4,15	0.34	0	3,3,17	0.32	0
3	LDA	B	309	-	5,5,15	0.31	0	4,4,17	0.42	0
3	LDA	B	306	-	13,15,15	2.39	2 (15%)	14,17,17	0.49	0
6	PG4	A	310	-	9,9,12	0.51	0	8,8,11	0.28	0
3	LDA	B	304	-	13,15,15	2.28	2 (15%)	14,17,17	0.81	0
3	LDA	B	310	-	3,3,15	0.35	0	2,2,17	0.64	0
8	SO4	B	334	-	4,4,4	0.30	0	6,6,6	0.65	0
4	GOL	A	305	-	5,5,5	1.37	1 (20%)	5,5,5	1.10	1 (20%)
7	PEE	A	311	-	8,8,50	1.07	1 (12%)	9,10,55	0.48	0
3	LDA	B	301	-	10,12,15	2.71	2 (20%)	11,14,17	0.71	0
3	LDA	B	316	-	7,7,15	0.29	0	6,6,17	0.43	0
4	GOL	A	302	-	5,5,5	0.85	0	5,5,5	1.07	0
3	LDA	B	321	-	4,4,15	0.31	0	3,3,17	0.41	0
3	LDA	B	313	-	9,9,15	0.26	0	8,8,17	0.53	0
3	LDA	B	325	-	2,2,15	0.26	0	1,1,17	0.00	0
3	LDA	B	312	-	9,11,15	2.85	2 (22%)	10,13,17	0.63	0
3	LDA	B	302	-	13,15,15	2.52	2 (15%)	14,17,17	0.71	0
4	GOL	B	332	-	5,5,5	0.94	0	5,5,5	0.99	0
3	LDA	A	301	-	3,3,15	0.33	0	2,2,17	0.68	0
3	LDA	B	322	-	7,7,15	0.26	0	6,6,17	0.64	0
4	GOL	A	304	-	5,5,5	1.14	1 (20%)	5,5,5	1.01	0
3	LDA	B	323	-	8,8,15	0.28	0	7,7,17	0.49	0
3	LDA	B	305	-	7,7,15	0.29	0	6,6,17	0.47	0
9	P4G	B	335	-	8,8,10	0.56	0	7,7,9	0.26	0
3	LDA	B	328	-	3,3,15	0.38	0	2,2,17	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LDA	B	330	-	-	1/1/1/13	-
3	LDA	B	317	-	-	0/1/1/13	-
3	LDA	B	308	-	-	1/3/3/13	-
4	GOL	A	307	-	-	2/4/4/4	-
4	GOL	A	303	-	-	3/4/4/4	-
5	ATP	A	308	-	-	2/22/38/38	0/3/3/3
3	LDA	B	329	-	-	5/13/13/13	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LDA	B	314	-	-	0/6/6/13	-
4	GOL	A	306	-	-	0/4/4/4	-
3	LDA	B	307	-	-	2/5/5/13	-
4	GOL	B	333	-	-	2/4/4/4	-
3	LDA	B	315	-	-	4/13/13/13	-
3	LDA	B	311	-	-	1/3/3/13	-
3	LDA	B	303	-	-	9/13/13/13	-
3	LDA	B	318	-	-	0/2/2/13	-
6	PG4	A	309	-	-	5/10/10/10	-
3	LDA	B	324	-	-	0/2/2/13	-
3	LDA	B	309	-	-	3/3/3/13	-
3	LDA	B	306	-	-	3/13/13/13	-
6	PG4	A	310	-	-	4/7/7/10	-
3	LDA	B	304	-	-	5/13/13/13	-
3	LDA	B	310	-	-	1/1/1/13	-
4	GOL	A	305	-	-	0/4/4/4	-
7	PEE	A	311	-	-	2/8/8/54	-
3	LDA	B	301	-	-	1/10/10/13	-
3	LDA	B	316	-	-	1/5/5/13	-
4	GOL	A	302	-	-	2/4/4/4	-
3	LDA	B	321	-	-	0/2/2/13	-
3	LDA	B	313	-	-	6/7/7/13	-
3	LDA	B	312	-	-	7/9/9/13	-
3	LDA	B	302	-	-	6/13/13/13	-
4	GOL	B	332	-	-	0/4/4/4	-
3	LDA	A	301	-	-	1/1/1/13	-
3	LDA	B	322	-	-	1/5/5/13	-
4	GOL	A	304	-	-	2/4/4/4	-
3	LDA	B	323	-	-	1/6/6/13	-
3	LDA	B	305	-	-	1/5/5/13	-
9	P4G	B	335	-	-	3/6/6/8	-
3	LDA	B	328	-	-	0/1/1/13	-

The worst 5 of 32 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	331	LDA	O1-N1	-12.56	1.25	1.42
3	B	329	LDA	O1-N1	-6.95	1.25	1.42
3	B	315	LDA	O1-N1	-6.86	1.25	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	314	LDA	O1-N1	-6.82	1.25	1.42
3	B	312	LDA	O1-N1	-6.81	1.25	1.42

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	308	ATP	C5-C4-N3	-4.64	120.33	126.72
5	A	308	ATP	N3-C4-N9	4.31	134.49	127.17
5	A	308	ATP	N3-C2-N1	-4.20	122.22	128.58
5	A	308	ATP	C4-N9-C8	3.54	109.45	105.74
5	A	308	ATP	C2-N3-C4	3.41	120.17	111.83

There are no chirality outliers.

5 of 87 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	301	LDA	C2-C1-N1-CM2
3	B	303	LDA	C2-C1-N1-O1
3	B	303	LDA	C2-C1-N1-CM1
3	B	303	LDA	C2-C1-N1-CM2
3	B	312	LDA	C2-C1-N1-O1


There are no ring outliers.

14 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	314	LDA	1	0
3	B	315	LDA	4	0
3	B	311	LDA	1	0
3	B	303	LDA	1	0
6	A	309	PG4	1	0
3	B	306	LDA	1	0
6	A	310	PG4	1	0
3	B	304	LDA	5	0
4	A	305	GOL	1	0
7	A	311	PEE	1	0
3	B	313	LDA	1	0
3	B	312	LDA	1	0
3	B	302	LDA	2	0
9	B	335	P4G	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	238/244 (97%)	-0.23	10 (4%)	40 40	19, 40, 77, 133	10 (4%)
2	B	256/256 (100%)	-0.01	18 (7%)	22 22	22, 43, 82, 132	27 (10%)
All	All	494/500 (98%)	-0.11	28 (5%)	29 29	19, 42, 82, 133	37 (7%)

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	129	LEU	5.9
1	A	237	LEU	4.7
1	A	0	MET	4.2
2	B	128	THR	4.0
2	B	1	MET	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	LDA	B	318	5/16	0.55	0.24	67,79,82,83	0
3	LDA	B	321	5/16	0.60	0.20	81,83,86,86	0
3	LDA	B	331	5/16	0.66	0.32	119,123,124,124	0
3	LDA	B	307	8/16	0.67	0.24	97,117,122,122	0
3	LDA	B	329	16/16	0.73	0.27	64,88,115,118	0
3	LDA	B	325	3/16	0.73	0.22	76,76,78,78	0
3	LDA	B	330	4/16	0.76	0.24	88,90,92,92	0
3	LDA	B	324	5/16	0.76	0.19	69,76,81,81	0
3	LDA	B	326	5/16	0.78	0.24	77,81,84,84	0
3	LDA	B	312	12/16	0.78	0.25	81,101,121,123	0
3	LDA	B	306	16/16	0.79	0.18	73,80,116,117	0
3	LDA	B	309	6/16	0.80	0.20	62,75,78,79	0
3	LDA	B	314	9/16	0.81	0.23	76,94,109,116	0
3	LDA	B	319	3/16	0.82	0.23	56,56,67,74	0
3	LDA	B	305	8/16	0.82	0.20	67,72,73,76	0
7	PEE	A	311	9/51	0.83	0.21	100,105,110,110	0
4	GOL	A	307	6/6	0.85	0.19	80,90,93,100	0
3	LDA	B	322	8/16	0.85	0.17	64,68,73,76	0
3	LDA	B	317	4/16	0.86	0.20	47,61,71,80	0
3	LDA	B	313	10/16	0.86	0.20	79,85,87,88	0
3	LDA	B	301	13/16	0.86	0.17	62,68,76,77	0
3	LDA	B	316	8/16	0.86	0.17	75,80,86,87	0
3	LDA	B	328	4/16	0.86	0.18	69,70,73,78	0
9	P4G	B	335	9/11	0.86	0.22	84,88,93,94	0
3	LDA	B	308	6/16	0.87	0.15	67,71,76,79	0
3	LDA	B	323	9/16	0.87	0.15	69,74,82,82	0
3	LDA	B	311	6/16	0.87	0.17	73,76,79,80	0
4	GOL	A	306	6/6	0.88	0.20	69,73,80,82	0
3	LDA	B	315	16/16	0.88	0.20	71,84,96,98	0
3	LDA	B	327	3/16	0.89	0.23	66,66,75,77	0
3	LDA	A	301	4/16	0.90	0.26	84,86,87,89	0
6	PG4	A	309	13/13	0.90	0.17	64,83,96,96	0
6	PG4	A	310	10/13	0.90	0.15	60,73,86,87	0
3	LDA	B	302	16/16	0.90	0.13	58,78,91,94	0
3	LDA	B	320	3/16	0.90	0.20	62,62,63,74	0
4	GOL	A	303	6/6	0.91	0.14	63,69,74,74	0
4	GOL	B	333	6/6	0.91	0.14	55,57,66,66	0
3	LDA	B	303	16/16	0.91	0.15	56,71,84,88	0
4	GOL	A	305	6/6	0.92	0.21	72,79,82,85	0
3	LDA	B	310	4/16	0.92	0.14	70,79,79,81	0
4	GOL	A	304	6/6	0.93	0.18	61,82,88,97	0
4	GOL	A	302	6/6	0.94	0.12	41,71,76,88	0
3	LDA	B	304	16/16	0.95	0.12	26,46,61,66	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
8	SO4	B	334	5/5	0.96	0.10	49,50,52,64	0
4	GOL	B	332	6/6	0.96	0.07	37,42,48,51	0
5	ATP	A	308	31/31	0.99	0.04	26,28,31,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers ⓘ

There are no such residues in this entry.