



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:01 AM UTC

PDB ID : 3MBW / pdb_00003mbw
Title : Crystal structure of the human ephrin A2 LBD and CRD domains in complex with ephrin A1
Authors : Walker, J.R.; Yermekbayeva, L.; Seitova, A.; Butler-Cole, C.; Bountra, C.; Weigelt, J.; Arrowsmith, C.H.; Edwards, A.M.; Bochkarev, A.; Dhe-Paganon, S.; Structural Genomics Consortium (SGC)
Deposited on : 2010-03-26
Resolution : 2.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4591 (2.84-2.80)
Clashscore	190562	5010 (2.84-2.80)
Ramachandran outliers	187476	4916 (2.84-2.80)
Sidechain outliers	187428	4918 (2.84-2.80)
RSRZ outliers	180081	4594 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	12% 72% 14% • 13%
2	B	181	4% 52% 15% • 30%
3	C	4	75% 25%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	UNX	A	327	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3292 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ephrin type-A receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	286	Total	C	N	O	S	0	1	0
			2160	1365	360	413	22			

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	ALA	-	expression tag	UNP P29317
A	-2	PRO	-	expression tag	UNP P29317
A	-1	GLU	-	expression tag	UNP P29317
A	0	HIS	-	expression tag	UNP P29317
A	1	HIS	-	expression tag	UNP P29317
A	2	HIS	-	expression tag	UNP P29317
A	3	HIS	-	expression tag	UNP P29317
A	4	HIS	-	expression tag	UNP P29317
A	5	HIS	-	expression tag	UNP P29317
A	6	ASP	-	expression tag	UNP P29317
A	7	TYR	-	expression tag	UNP P29317
A	8	ASP	-	expression tag	UNP P29317
A	9	ILE	-	expression tag	UNP P29317
A	10	PRO	-	expression tag	UNP P29317
A	11	THR	-	expression tag	UNP P29317
A	12	THR	-	expression tag	UNP P29317
A	13	GLU	-	expression tag	UNP P29317
A	14	ASN	-	expression tag	UNP P29317
A	15	LEU	-	expression tag	UNP P29317
A	16	TYR	-	expression tag	UNP P29317
A	17	PHE	-	expression tag	UNP P29317
A	18	GLN	-	expression tag	UNP P29317
A	19	GLY	-	expression tag	UNP P29317
A	20	ALA	-	expression tag	UNP P29317
A	21	MET	-	expression tag	UNP P29317
A	22	ASP	-	expression tag	UNP P29317

- Molecule 2 is a protein called Ephrin-A1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	127	Total	C	N	O	S	0	0	0
			1055	674	182	194	5			

There are 27 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-9	ALA	-	expression tag	UNP P20827
B	-8	PRO	-	expression tag	UNP P20827
B	-7	GLU	-	expression tag	UNP P20827
B	-6	HIS	-	expression tag	UNP P20827
B	-5	HIS	-	expression tag	UNP P20827
B	-4	HIS	-	expression tag	UNP P20827
B	-3	HIS	-	expression tag	UNP P20827
B	-2	HIS	-	expression tag	UNP P20827
B	-1	HIS	-	expression tag	UNP P20827
B	0	ASP	-	expression tag	UNP P20827
B	1	TYR	-	expression tag	UNP P20827
B	2	ASP	-	expression tag	UNP P20827
B	3	ILE	-	expression tag	UNP P20827
B	4	PRO	-	expression tag	UNP P20827
B	5	THR	-	expression tag	UNP P20827
B	6	THR	-	expression tag	UNP P20827
B	7	GLU	-	expression tag	UNP P20827
B	8	ASN	-	expression tag	UNP P20827
B	9	LEU	-	expression tag	UNP P20827
B	10	TYR	-	expression tag	UNP P20827
B	11	PHE	-	expression tag	UNP P20827
B	12	GLN	-	expression tag	UNP P20827
B	13	GLY	-	expression tag	UNP P20827
B	14	ALA	-	expression tag	UNP P20827
B	15	MET	-	expression tag	UNP P20827
B	16	ASP	-	expression tag	UNP P20827
B	159	VAL	ASP	SEE REMARK 999	UNP P20827

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	4	Total	C	N	O	0	0	0
			49	28	2	19			

- Molecule 4 is UNKNOWN ATOM OR ION (CCD ID: UNX) (formula: X).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	X	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	22	Total	O	0	0
			22	22		
5	B	5	Total	O	0	0
			5	5		

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	58.27Å 215.79Å 107.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.10 – 2.81 34.10 – 2.81	Depositor EDS
% Data completeness (in resolution range)	(Not available) (34.10-2.81) 99.6 (34.10-2.81)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.33 (at 2.81Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.226 , 0.272 0.238 , 0.287	Depositor DCC
R_{free} test set	874 reflections (5.17%)	wwPDB-VP
Wilson B-factor (Å ²)	64.4	Xtriage
Anisotropy	0.327	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 79.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3292	wwPDB-VP
Average B, all atoms (Å ²)	80.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, UNX, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	0/2211	1.24	6/3004 (0.2%)
2	B	0.76	0/1086	1.25	6/1471 (0.4%)
All	All	0.78	0/3297	1.24	12/4475 (0.3%)

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	72	VAL	N-CA-C	6.06	116.22	108.82
2	B	26	ASN	CA-CB-CG	6.06	118.66	112.60
2	B	108	PHE	CA-C-N	5.64	130.44	122.09
2	B	108	PHE	C-N-CA	5.64	130.44	122.09
2	B	33	ARG	N-CA-C	5.49	119.08	112.38
1	A	148	ASP	N-CA-C	-5.33	105.92	112.90
1	A	153	SER	N-CA-C	5.30	117.06	111.28
2	B	20	ARG	N-CA-C	5.19	117.91	109.24
1	A	239	GLY	CA-C-N	5.16	124.88	119.92
1	A	239	GLY	C-N-CA	5.16	124.88	119.92
1	A	163	LEU	N-CA-C	5.07	116.91	107.99
1	A	161	VAL	N-CA-CB	5.02	117.69	111.41

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2160	0	1999	18	1
2	B	1055	0	973	17	0
3	C	49	0	40	4	0
4	A	1	0	0	0	1
5	A	22	0	0	0	0
5	B	5	0	0	0	0
All	All	3292	0	3012	32	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:53:HIS:H	3:C:1:NAG:H83	1.36	0.91
2:B:53:HIS:N	3:C:1:NAG:H83	2.07	0.69
1:A:123:TYR:HB3	1:A:142:ILE:HD11	1.78	0.65
2:B:53:HIS:O	3:C:1:NAG:H82	1.97	0.64
1:A:103:ARG:HA	1:A:151:THR:HB	1.84	0.59
2:B:22:THR:HG23	2:B:50:ILE:HD13	1.86	0.58
2:B:112:THR:HG22	2:B:119:GLU:HG3	1.86	0.57
2:B:66:GLN:HB3	2:B:133:ILE:HD12	1.90	0.54
2:B:20:ARG:HB3	2:B:46:TYR:HB2	1.90	0.53
1:A:52:TRP:CD1	1:A:81:LEU:HB2	2.44	0.52
1:A:141:LYS:NZ	1:A:144:THR:OG1	2.41	0.51
1:A:99:LYS:HB2	1:A:193:SER:HB3	1.92	0.51
2:B:53:HIS:H	3:C:1:NAG:C8	2.15	0.51
2:B:39:ILE:HG12	2:B:143:LEU:HD11	1.94	0.50
1:A:149:GLU:HB3	1:A:162:LYS:HD3	1.93	0.50
1:A:58:ILE:HD12	2:B:88:VAL:HG12	1.94	0.49
2:B:25:TRP:NE1	2:B:130:SER:OG	2.45	0.49
1:A:200:LYS:HG2	1:A:215:ILE:HA	1.95	0.48
2:B:22:THR:HG23	2:B:50:ILE:CD1	2.43	0.48
1:A:243:PRO:HG3	1:A:260:CYS:SG	2.54	0.47
1:A:103:ARG:HB3	1:A:188:CYS:HB3	2.00	0.44
1:A:80:TRP:CE2	1:A:183:GLN:HG3	2.53	0.44
1:A:294:PRO:HA	1:A:295:GLU:HA	1.77	0.43
2:B:43:LEU:HD11	2:B:110:ARG:HB2	2.01	0.42
2:B:85:LYS:C	2:B:87:GLN:H	2.28	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:127:ASP:HA	1:A:176:LYS:HB3	2.02	0.42
1:A:56:GLN:HE22	2:B:104:LEU:HA	1.84	0.42
1:A:201:CYS:HB3	1:A:253:TRP:CE2	2.55	0.41
1:A:56:GLN:HB3	2:B:90:TRP:CZ3	2.55	0.41
1:A:46:HIS:HA	1:A:47:PRO:C	2.46	0.41
1:A:103:ARG:HD2	1:A:156:PHE:HZ	1.86	0.41
2:B:112:THR:HG22	2:B:119:GLU:CG	2.51	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:143:ASP:OD1	4:A:327:UNX:UNK[3_555]	2.05	0.15

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	277/330 (84%)	260 (94%)	16 (6%)	1 (0%)	30	58
2	B	123/181 (68%)	118 (96%)	4 (3%)	1 (1%)	16	41
All	All	400/511 (78%)	378 (94%)	20 (5%)	2 (0%)	24	53

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	86	ASP
1	A	222	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/275 (82%)	208 (93%)	16 (7%)	13	38
2	B	112/165 (68%)	102 (91%)	10 (9%)	9	27
All	All	336/440 (76%)	310 (92%)	26 (8%)	11	34

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25	GLN
1	A	44	LEU
1	A	128	LEU
1	A	153	SER
1	A	157	GLU
1	A	161	VAL
1	A	168	ARG
1	A	185	ILE
1	A	204	LEU
1	A	206	GLN
1	A	213	GLU
1	A	215	ILE
1	A	236	VAL
1	A	284	GLU
1	A	293	CYS
1	A	295	GLU
2	B	35	GLU
2	B	43	LEU
2	B	55	GLU
2	B	71	LEU
2	B	91	GLN
2	B	98	LYS
2	B	104	LEU
2	B	130	SER
2	B	135	GLN
2	B	138	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	GLN
1	A	57	ASN
1	A	77	GLN
1	A	79	ASN
1	A	133	ASN
1	A	135	GLN
1	A	296	HIS
2	B	74	HIS
2	B	81	GLN
2	B	91	GLN
2	B	136	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	C	1	3,2	14,14,15	0.97	0	17,19,21	2.00	3 (17%)
3	NAG	C	2	3	14,14,15	1.33	3 (21%)	17,19,21	1.98	5 (29%)
3	BMA	C	3	3	11,11,12	2.01	4 (36%)	15,15,17	2.03	6 (40%)
3	MAN	C	4	3	10,10,12	1.82	4 (40%)	14,14,17	1.74	4 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	3,2	-	2/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
3	BMA	C	3	3	-	0/2/19/22	0/1/1/1
3	MAN	C	4	3	-	-	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	3	BMA	C2-C3	3.39	1.57	1.52
3	C	3	BMA	C4-C5	3.15	1.59	1.53
3	C	3	BMA	C4-C3	2.84	1.59	1.52
3	C	2	NAG	C4-C3	2.84	1.59	1.52
3	C	4	MAN	C4-C3	2.46	1.58	1.52
3	C	4	MAN	C1-C2	2.38	1.57	1.52
3	C	3	BMA	O2-C2	2.26	1.48	1.43
3	C	4	MAN	O2-C2	2.22	1.48	1.43
3	C	4	MAN	C4-C5	2.21	1.57	1.52
3	C	2	NAG	C7-N2	2.14	1.41	1.34
3	C	2	NAG	O7-C7	2.04	1.27	1.23

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1	NAG	C1-C2-N2	5.92	119.77	110.43
3	C	2	NAG	C1-C2-N2	4.14	116.96	110.43
3	C	1	NAG	C4-C3-C2	-3.73	105.55	111.02
3	C	2	NAG	C2-N2-C7	-3.69	117.95	122.90
3	C	3	BMA	O5-C1-C2	3.58	119.34	110.79
3	C	3	BMA	C2-C3-C4	3.53	117.08	110.86
3	C	4	MAN	O2-C2-C1	3.29	116.76	109.22
3	C	4	MAN	C3-C4-C5	3.11	114.53	109.81
3	C	2	NAG	O3-C3-C2	-3.01	103.14	109.40
3	C	2	NAG	C1-O5-C5	-2.96	108.22	112.19
3	C	4	MAN	C2-C3-C4	2.76	115.72	110.86
3	C	3	BMA	O2-C2-C1	2.75	115.52	109.22
3	C	3	BMA	O3-C3-C4	2.63	116.58	110.38
3	C	3	BMA	O2-C2-C3	2.45	115.23	110.15
3	C	1	NAG	C2-N2-C7	2.31	126.00	122.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	3	BMA	C1-C2-C3	2.21	112.86	109.64
3	C	2	NAG	C6-C5-C4	2.21	118.44	113.02
3	C	4	MAN	O3-C3-C4	2.18	115.52	110.38

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	1	NAG	C1-C2-N2-C7
3	C	1	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1	NAG	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry

Of 1 ligands modelled in this entry, 1 is unknown - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	286/330 (86%)	0.70	38 (13%) 7 5	20, 72, 137, 174	1 (0%)
2	B	127/181 (70%)	0.63	8 (6%) 26 19	53, 81, 109, 126	0
All	All	413/511 (80%)	0.68	46 (11%) 10 7	20, 78, 132, 174	1 (0%)

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	303	GLY	4.5
2	B	59	VAL	4.0
1	A	315	ARG	3.9
1	A	321	ALA	3.6
1	A	50	LYS	3.4
1	A	72	VAL	3.3
1	A	271	ASP	3.2
1	A	267	GLU	3.1
1	A	294	PRO	3.1
1	A	112	ALA	2.9
1	A	110	GLY	2.9
2	B	55	GLU	2.9
1	A	102	VAL	2.9
2	B	86	ASP	2.8
1	A	285	ALA	2.7
1	A	296	HIS	2.7
1	A	114	SER	2.7
2	B	99	HIS	2.6
1	A	152	VAL	2.5
1	A	318	GLN	2.5
1	A	113	SER	2.5
1	A	153	SER	2.5
1	A	277	SER	2.5
1	A	48	TYR	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	149	GLU	2.4
1	A	264	ALA	2.4
1	A	306	SER	2.4
1	A	310	GLU	2.4
1	A	111	GLY	2.4
1	A	160	HIS	2.3
1	A	150	ILE	2.3
1	A	137	ARG	2.3
1	A	161	VAL	2.3
1	A	104	ASP	2.3
1	A	278	PRO	2.3
1	A	164	ASN	2.2
2	B	134	HIS	2.2
1	A	238	PRO	2.2
1	A	185	ILE	2.2
2	B	139	ARG	2.2
1	A	316	ALA	2.2
2	B	73	GLU	2.2
2	B	83	GLN	2.1
1	A	77	GLN	2.1
1	A	281	PHE	2.1
1	A	148	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MAN	C	4	10/12	0.42	0.17	104,107,108,109	0
3	BMA	C	3	11/12	0.53	0.13	95,99,101,101	0
3	NAG	C	2	14/15	0.82	0.12	80,85,87,88	0
3	NAG	C	1	14/15	0.89	0.12	63,68,70,71	0

The following is a graphical depiction of the model fit to experimental electron density for oligosac-

charide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	UNX	A	327	1/1	0.53	0.60	49,49,49,49	0

6.5 Other polymers [i](#)

There are no such residues in this entry.