



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 08:51 AM UTC

PDB ID : 5MBM / pdb_00005mbm
Title : Cathepsin B in complex with DARPin 8h6
Authors : Turk, D.; Kramer, L.; Renko, M.; Turk, B.
Deposited on : 2016-11-08
Resolution : 2.76 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	256	4% 29% 49% 21% .
1	B	256	16% 32% 40% 24% .
2	C	171	9% 25% 47% 19% 5%
2	D	171	4% 31% 42% 21% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	SCH	A	29	-	-	X	-
1	SCH	B	29	-	-	X	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6499 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cathepsin B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	256	Total	C	N	O	S	14	0	0
			1971	1235	337	380	19			
1	B	255	Total	C	N	O	S	15	0	0
			1963	1229	336	379	19			

- Molecule 2 is a protein called DARPin 8h6.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	163	Total	C	N	O	0	0	0
			1211	751	217	243			
2	D	163	Total	C	N	O	29	0	0
			1211	751	217	243			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	47	Total	O	0	0
			47	47		
3	B	51	Total	O	0	0
			51	51		
3	C	22	Total	O	0	0
			22	22		
3	D	23	Total	O	0	0
			23	23		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Cathepsin B




• Molecule 1: Cathepsin B



• Molecule 2: DARPin 8h6



• Molecule 2: DARPin 8h6



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	101.45Å 201.52Å 46.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.70 – 2.76 46.70 – 2.76	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.70-2.76) 98.5 (46.70-2.76)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	0.18	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.42 (at 2.77Å)	Xtriage
Refinement program	MAIN 2016	Depositor
R, R_{free}	0.256 , 0.281 0.258 , 0.268	Depositor DCC
R_{free} test set	1257 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	56.3	Xtriage
Anisotropy	0.548	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 71.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.40$, $\langle L^2 \rangle = 0.22$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	6499	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SCH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.71	41/2021 (2.0%)	1.85	68/2744 (2.5%)
1	B	1.54	33/2013 (1.6%)	2.05	105/2733 (3.8%)
2	C	1.64	24/1232 (1.9%)	1.87	38/1682 (2.3%)
2	D	1.74	21/1232 (1.7%)	2.02	52/1682 (3.1%)
All	All	1.65	119/6498 (1.8%)	1.95	263/8841 (3.0%)

The worst 5 of 119 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	148	TYR	CA-C	11.99	1.68	1.52
2	C	63	ASP	CA-C	9.85	1.65	1.52
1	A	110	HIS	CA-C	-9.09	1.41	1.52
1	B	253	THR	CA-C	-9.04	1.40	1.52
1	A	107	PRO	CA-C	-8.87	1.42	1.52

The worst 5 of 263 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	66	MET	N-CA-C	-13.75	94.78	111.69
1	B	55	SER	CA-C-N	-10.60	104.20	120.31
1	B	55	SER	C-N-CA	-10.60	104.20	120.31
1	B	80	TRP	N-CA-C	-10.23	99.33	112.23
2	C	171	ASN	CA-CB-CG	9.89	122.50	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1971	0	1823	169	0
1	B	1963	0	1812	163	1
2	C	1211	0	1164	139	2
2	D	1211	0	1164	82	2
3	A	47	0	0	3	0
3	B	51	0	0	3	0
3	C	22	0	0	3	0
3	D	23	0	0	1	0
All	All	6499	0	5963	518	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 43.

The worst 5 of 518 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:108:CYS:SG	1:B:109:GLU:N	2.41	0.94
2:C:168:ALA:O	2:C:171:ASN:HB2	1.72	0.89
1:A:29:SCH:HA	1:A:199:HIS:CE1	2.09	0.88
2:C:110:GLN:HE21	2:C:110:GLN:HA	1.39	0.86
1:A:177:TYR:CD2	1:A:190:HIS:CE1	2.65	0.84

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:68:TYR:CD2	2:D:31:ARG:NH1[4_558]	1.52	0.68
1:B:111:HIS:CD2	1:B:141:LYS:NZ[1_554]	2.15	0.05
2:C:68:TYR:CG	2:D:31:ARG:NH1[4_558]	2.15	0.05

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	253/256 (99%)	210 (83%)	39 (15%)	4 (2%)	7	14
1	B	252/256 (98%)	195 (77%)	48 (19%)	9 (4%)	2	4
2	C	161/171 (94%)	117 (73%)	34 (21%)	10 (6%)	1	1
2	D	161/171 (94%)	136 (84%)	14 (9%)	11 (7%)	1	1
All	All	827/854 (97%)	658 (80%)	135 (16%)	34 (4%)	2	3

5 of 34 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	122	GLU
1	B	93	LEU
1	B	253	THR
2	C	41	ASN
2	C	154	ASP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	210/210 (100%)	188 (90%)	22 (10%)	6	13
1	B	209/210 (100%)	187 (90%)	22 (10%)	6	13
2	C	122/129 (95%)	106 (87%)	16 (13%)	4	7
2	D	122/129 (95%)	101 (83%)	21 (17%)	2	3
All	All	663/678 (98%)	582 (88%)	81 (12%)	5	8

5 of 81 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	142	ASP
2	D	99	LEU
2	C	166	LYS
2	D	26	GLN
2	D	111	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
2	C	36	ASN
2	D	155	ASN
2	C	101	ASN
2	D	110	GLN
2	C	88	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	SCH	A	29	1	6,7,8	0.55	0	4,7,9	0.67	0
1	SCH	B	29	1	6,7,8	0.46	0	4,7,9	1.53	1 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SCH	A	29	1	-	1/2/6/8	-
1	SCH	B	29	1	-	1/2/6/8	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	29	SCH	CB-SG-SD	2.75	110.98	103.86

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	29	SCH	CA-CB-SG-SD
1	B	29	SCH	CA-CB-SG-SD

There are no ring outliers.

2 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	29	SCH	10	0
1	B	29	SCH	6	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	254/256 (99%)	0.21	11 (4%) 40 38	13, 49, 92, 99	2 (0%)
1	B	254/256 (99%)	1.03	40 (15%) 5 4	20, 77, 98, 99	5 (1%)
2	C	163/171 (95%)	0.67	15 (9%) 14 14	28, 71, 99, 99	0
2	D	161/171 (94%)	0.34	7 (4%) 40 38	21, 55, 87, 99	3 (1%)
All	All	832/854 (97%)	0.57	73 (8%) 15 15	13, 61, 98, 99	10 (1%)

The worst 5 of 73 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	160	ILE	5.2
1	B	164	ILE	4.4
1	B	217	VAL	4.2
1	B	181	LEU	4.0
1	B	205	GLY	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SCH	A	29	8/9	0.76	0.20	54,65,70,84	0
1	SCH	B	29	8/9	0.82	0.14	56,66,70,71	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.