



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 14, 2026 – 11:00 AM UTC

PDB ID : 3MCE / pdb_00003mce
Title : Crystal structure of the NAC domain of alpha subunit of nascent polypeptide
-associated complex(NAC)
Authors : Wang, L.F.; Zhang, W.C.; Wang, L.; Zhang, X.J.C.; Li, X.M.; Rao, Z.
Deposited on : 2010-03-29
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	61	 7% 41% 28% 25% 7%
1	B	61	 59% 25% 13% . .
1	C	61	 2% 52% 33% 11% .
1	D	61	 8% 41% 28% 16% 8% 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	IOD	B	1	-	-	X	-
2	IOD	B	6	-	-	X	-
2	IOD	C	4	-	-	X	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1884 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nascent polypeptide-associated complex subunit alpha.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	A	57	445	288	74	83	0	0	0
1	B	60	469	302	78	89	0	0	0
1	C	59	459	297	76	86	0	0	0
1	D	57	450	289	75	86	0	0	0

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	73	GLY	-	expression tag	UNP Q13765
A	74	PRO	-	expression tag	UNP Q13765
A	75	LEU	-	expression tag	UNP Q13765
A	76	GLY	-	expression tag	UNP Q13765
A	77	SER	-	expression tag	UNP Q13765
A	78	PRO	-	expression tag	UNP Q13765
A	79	GLU	-	expression tag	UNP Q13765
A	80	PHE	-	expression tag	UNP Q13765
B	73	GLY	-	expression tag	UNP Q13765
B	74	PRO	-	expression tag	UNP Q13765
B	75	LEU	-	expression tag	UNP Q13765
B	76	GLY	-	expression tag	UNP Q13765
B	77	SER	-	expression tag	UNP Q13765
B	78	PRO	-	expression tag	UNP Q13765
B	79	GLU	-	expression tag	UNP Q13765
B	80	PHE	-	expression tag	UNP Q13765
C	73	GLY	-	expression tag	UNP Q13765
C	74	PRO	-	expression tag	UNP Q13765
C	75	LEU	-	expression tag	UNP Q13765
C	76	GLY	-	expression tag	UNP Q13765
C	77	SER	-	expression tag	UNP Q13765

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	78	PRO	-	expression tag	UNP Q13765
C	79	GLU	-	expression tag	UNP Q13765
C	80	PHE	-	expression tag	UNP Q13765
D	73	GLY	-	expression tag	UNP Q13765
D	74	PRO	-	expression tag	UNP Q13765
D	75	LEU	-	expression tag	UNP Q13765
D	76	GLY	-	expression tag	UNP Q13765
D	77	SER	-	expression tag	UNP Q13765
D	78	PRO	-	expression tag	UNP Q13765
D	79	GLU	-	expression tag	UNP Q13765
D	80	PHE	-	expression tag	UNP Q13765

- Molecule 2 is IODIDE ION (CCD ID: IOD) (formula: I).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total I 2 2	0	0
2	B	3	Total I 3 3	0	0
2	C	2	Total I 2 2	0	0
2	D	1	Total I 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	17	Total O 17 17	0	0
3	B	18	Total O 18 18	0	0
3	C	11	Total O 11 11	0	0
3	D	7	Total O 7 7	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nascent polypeptide-associated complex subunit alpha



- Molecule 1: Nascent polypeptide-associated complex subunit alpha



- Molecule 1: Nascent polypeptide-associated complex subunit alpha



- Molecule 1: Nascent polypeptide-associated complex subunit alpha



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	54.59Å 59.60Å 69.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.90 – 2.40 29.90 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.4 (29.90-2.40) 98.3 (29.90-2.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	10.41 (at 2.39Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.247 , 0.277 0.234 , 0.273	Depositor DCC
R_{free} test set	910 reflections (9.94%)	wwPDB-VP
Wilson B-factor (Å ²)	36.9	Xtrriage
Anisotropy	0.606	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 53.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	1884	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 61.37 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.3231e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IOD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.71	2/451 (0.4%)	1.20	1/606 (0.2%)
1	B	1.95	7/477 (1.5%)	1.29	3/642 (0.5%)
1	C	1.87	5/467 (1.1%)	1.22	3/630 (0.5%)
1	D	1.87	7/457 (1.5%)	1.43	5/615 (0.8%)
All	All	1.85	21/1852 (1.1%)	1.29	12/2493 (0.5%)

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	121	ILE	C-O	-6.39	1.17	1.24
1	B	119	THR	C-O	-6.26	1.16	1.24
1	B	111	VAL	C-O	-6.04	1.17	1.24
1	D	103	LEU	C-O	-5.99	1.17	1.24
1	D	109	PRO	C-O	-5.95	1.18	1.23

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	107	THR	N-CA-C	-9.29	93.73	109.24
1	D	85	LEU	N-CA-C	-7.41	99.55	110.52
1	A	123	PHE	N-CA-C	7.14	120.05	108.76
1	B	102	ILE	N-CA-C	6.99	118.35	108.36
1	D	131	LEU	CA-CB-CG	-6.15	94.79	116.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	445	0	469	44	0
1	B	469	0	493	20	0
1	C	459	0	485	24	0
1	D	450	0	471	71	0
2	A	2	0	0	1	0
2	B	3	0	0	7	0
2	C	2	0	0	3	0
2	D	1	0	0	1	0
3	A	17	0	0	1	0
3	B	18	0	0	2	0
3	C	11	0	0	0	0
3	D	7	0	0	1	0
All	All	1884	0	1918	147	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 39.

The worst 5 of 147 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:83:LEU:CD1	1:D:85:LEU:HD22	1.29	1.55
1:D:83:LEU:HD11	1:D:85:LEU:CD2	1.49	1.40
1:D:83:LEU:C	1:D:83:LEU:HD13	1.49	1.30
1:A:83:LEU:HB3	1:D:83:LEU:CD2	1.68	1.22
1:D:83:LEU:CD1	1:D:85:LEU:CD2	2.14	1.11

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	53/61 (87%)	47 (89%)	6 (11%)	0	100	100
1	B	58/61 (95%)	56 (97%)	2 (3%)	0	100	100
1	C	57/61 (93%)	54 (95%)	3 (5%)	0	100	100
1	D	55/61 (90%)	52 (94%)	2 (4%)	1 (2%)	6	9
All	All	223/244 (91%)	209 (94%)	13 (6%)	1 (0%)	30	43

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	108	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	51/54 (94%)	36 (71%)	15 (29%)	0	0
1	B	54/54 (100%)	45 (83%)	9 (17%)	2	3
1	C	53/54 (98%)	41 (77%)	12 (23%)	1	1
1	D	52/54 (96%)	37 (71%)	15 (29%)	0	0
All	All	210/216 (97%)	159 (76%)	51 (24%)	1	1

5 of 51 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	100	LYS
1	C	132	SER
1	D	131	LEU
1	C	103	LEU
1	C	117	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	101	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	57/61 (93%)	0.63	4 (7%) 22 19	36, 55, 80, 98	0
1	B	60/61 (98%)	-0.08	0 100 100	30, 43, 65, 80	0
1	C	59/61 (96%)	-0.01	1 (1%) 69 65	29, 41, 68, 82	0
1	D	57/61 (93%)	0.53	5 (8%) 15 12	32, 54, 84, 104	0
All	All	233/244 (95%)	0.26	10 (4%) 40 36	29, 47, 80, 104	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	83	LEU	3.6
1	D	101	ASN	3.1
1	A	80	PHE	2.8
1	A	114	SER	2.8
1	A	102	ILE	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	IOD	C	4	1/1	0.83	0.11	39,39,39,39	1
2	IOD	C	8	1/1	0.93	0.05	9,9,9,9	1
2	IOD	B	6	1/1	0.94	0.05	18,18,18,18	1
2	IOD	D	2	1/1	0.95	0.06	26,26,26,26	1
2	IOD	A	3	1/1	0.96	0.07	18,18,18,18	1
2	IOD	A	7	1/1	0.96	0.12	19,19,19,19	1
2	IOD	B	5	1/1	0.97	0.05	16,16,16,16	1
2	IOD	B	1	1/1	0.98	0.10	20,20,20,20	1

6.5 Other polymers [i](#)

There are no such residues in this entry.