



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 11:05 AM UTC

PDB ID : 4MCO / pdb_00004mco
Title : Crystal structure of a TRAP periplasmic solute binding protein from *Rhodospirillum rubrum* (Rfer_1840), target EFI-510211, with bound malonate
Authors : Vetting, M.W.; Toro, R.; Bhosle, R.; Al Obaidi, N.F.; Morisco, L.L.; Wasserman, S.R.; Sojitra, S.; Zhao, S.; Stead, M.; Washington, E.; Scott Glenn, A.; Chowdhury, S.; Evans, B.; Hammonds, J.; Hillerich, B.; Love, J.; Seidel, R.D.; Imker, H.J.; Jacobson, M.P.; Gerlt, J.A.; Almo, S.C.; Enzyme Function Initiative (EFI)
Deposited on : 2013-08-21
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	345	
1	B	345	
1	C	345	

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
 Validation Pipeline (wwPDB-VP) : 2.49

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 16308 atoms, of which 7513 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called TRAP dicarboxylate transporter-DctP subunit.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	316	Total	C	H	N	O	S	0	0	0
			4908	1550	2476	410	454	18			
1	B	317	Total	C	H	N	O	S	0	2	0
			4936	1557	2492	414	455	18			
1	C	318	Total	C	H	N	O	S	0	6	0
			4969	1567	2505	415	464	18			

There are 21 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	339	ALA	-	expression tag	UNP Q21XD7
A	340	GLU	-	expression tag	UNP Q21XD7
A	341	ASN	-	expression tag	UNP Q21XD7
A	342	LEU	-	expression tag	UNP Q21XD7
A	343	TYR	-	expression tag	UNP Q21XD7
A	344	PHE	-	expression tag	UNP Q21XD7
A	345	GLN	-	expression tag	UNP Q21XD7
B	339	ALA	-	expression tag	UNP Q21XD7
B	340	GLU	-	expression tag	UNP Q21XD7
B	341	ASN	-	expression tag	UNP Q21XD7
B	342	LEU	-	expression tag	UNP Q21XD7
B	343	TYR	-	expression tag	UNP Q21XD7
B	344	PHE	-	expression tag	UNP Q21XD7
B	345	GLN	-	expression tag	UNP Q21XD7
C	339	ALA	-	expression tag	UNP Q21XD7
C	340	GLU	-	expression tag	UNP Q21XD7
C	341	ASN	-	expression tag	UNP Q21XD7
C	342	LEU	-	expression tag	UNP Q21XD7
C	343	TYR	-	expression tag	UNP Q21XD7
C	344	PHE	-	expression tag	UNP Q21XD7
C	345	GLN	-	expression tag	UNP Q21XD7

- Molecule 2 is MALONATE ION (CCD ID: MLI) (formula: $C_3H_2O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			9	3	2	4		
2	B	1	Total	C	H	O	0	0
			9	3	2	4		
2	C	1	Total	C	H	O	0	0
			9	3	2	4		

- Molecule 3 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	S	0	0
			32	8	17	2	4	1		
3	A	1	Total	C	H	N	O	S	0	0
			32	8	17	2	4	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	483	Total	O	0	0
			483	483		
4	B	473	Total	O	0	0
			473	473		
4	C	448	Total	O	0	0
			448	448		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: TRAP dicarboxylate transporter-DctP subunit

Chain A: 



- Molecule 1: TRAP dicarboxylate transporter-DctP subunit

Chain B: 



- Molecule 1: TRAP dicarboxylate transporter-DctP subunit

Chain C: 



Q345

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	75.11Å 97.11Å 133.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	32.70 – 1.60 32.70 – 1.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (32.70-1.60) 100.0 (32.70-1.60)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.31 (at 1.60Å)	Xtriage
Refinement program	PHENIX 1.8.1_1168	Depositor
R, R_{free}	0.149 , 0.181 0.152 , 0.182	Depositor DCC
R_{free} test set	6457 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	14.2	Xtriage
Anisotropy	0.475	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	16308	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EPE, MLI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.59	0/2476	0.75	0/3328
1	B	0.57	0/2496	0.75	0/3356
1	C	0.56	0/2535	0.74	0/3406
All	All	0.58	0/7507	0.75	0/10090

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2432	2476	2475	7	0
1	B	2444	2492	2479	8	0
1	C	2464	2505	2474	11	0
2	A	7	2	2	0	0
2	B	7	2	2	0	0
2	C	7	2	2	0	0
3	A	30	34	34	3	0
4	A	483	0	0	2	0
4	B	473	0	0	3	0
4	C	448	0	0	3	0
All	All	8795	7513	7468	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:313:ARG:NH2	4:B:661:HOH:O	2.27	0.66
1:B:104:GLU:OE2	4:B:540:HOH:O	2.15	0.63
1:B:173:LYS:HG3	1:B:209:MET:HA	1.87	0.56
1:C:273:LYS:NZ	4:C:689:HOH:O	2.35	0.53
1:B:168:ASP:HA	1:C:255:LYS:HG2	1.90	0.53
1:C:183:MET:HE2	4:C:920:HOH:O	2.08	0.52
1:C:41:ILE:HG23	1:C:53:ASN:OD1	2.12	0.49
1:C:162:PRO:HB3	4:C:917:HOH:O	2.12	0.49
1:C:59[A]:GLU:HG2	1:C:67:LYS:HA	1.95	0.48
3:A:402:EPE:H92	4:A:873:HOH:O	2.14	0.48
1:B:276:LEU:HD11	1:B:280:LYS:HE3	1.97	0.47
1:A:117:VAL:HG13	1:A:123:ALA:HB2	1.99	0.45
1:B:66:MET:HE3	1:B:66:MET:HB2	1.67	0.45
1:A:161:LYS:HG2	3:A:402:EPE:S	2.57	0.45
1:C:209:MET:HE3	1:C:212:ALA:HB2	2.00	0.44
1:A:173:LYS:HD2	1:A:208:ALA:O	2.18	0.43
1:A:314:LYS:HE2	4:A:977:HOH:O	2.17	0.43
1:C:55:ALA:O	1:C:59[A]:GLU:HG3	2.19	0.42
1:C:263:MET:HE3	1:C:263:MET:HB2	1.89	0.42
1:B:141:GLU:OE2	4:B:784:HOH:O	2.22	0.42
1:C:113:MET:HE1	1:C:320:TYR:CG	2.55	0.41
1:B:117:VAL:HG13	1:B:123:ALA:HB2	2.01	0.41
1:A:161:LYS:HG2	3:A:402:EPE:O3S	2.21	0.41
1:A:67:LYS:HA	1:A:67:LYS:HD2	1.94	0.40
1:A:138:VAL:O	1:A:141:GLU:HG2	2.22	0.40
1:C:288:ALA:HB1	1:C:292:LYS:HE3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	314/345 (91%)	308 (98%)	6 (2%)	0	100	100
1	B	317/345 (92%)	314 (99%)	3 (1%)	0	100	100
1	C	322/345 (93%)	321 (100%)	1 (0%)	0	100	100
All	All	953/1035 (92%)	943 (99%)	10 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	251/273 (92%)	248 (99%)	3 (1%)	63	43
1	B	253/273 (93%)	250 (99%)	3 (1%)	63	43
1	C	258/273 (94%)	257 (100%)	1 (0%)	84	75
All	All	762/819 (93%)	755 (99%)	7 (1%)	70	55

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	161	LYS
1	A	173	LYS
1	A	195	LEU
1	B	41	ILE
1	B	60	LYS
1	B	173	LYS
1	C	255	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	35	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	260	GLN
1	B	35	GLN
1	C	35	GLN
1	C	260	GLN
1	C	335	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MLI	B	401	-	6,6,6	1.10	0	7,7,7	1.09	0
2	MLI	A	401	-	6,6,6	1.12	0	7,7,7	1.31	0
3	EPE	A	403	-	15,15,15	0.63	0	19,20,20	2.27	7 (36%)
2	MLI	C	401	-	6,6,6	1.18	0	7,7,7	1.24	1 (14%)
3	EPE	A	402	-	15,15,15	0.80	1 (6%)	19,20,20	2.26	6 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MLI	B	401	-	-	0/4/4/4	-
2	MLI	A	401	-	-	2/4/4/4	-
3	EPE	A	403	-	-	4/9/19/19	0/1/1/1
2	MLI	C	401	-	-	1/4/4/4	-
3	EPE	A	402	-	-	5/9/19/19	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	EPE	C10-S	2.68	1.81	1.77

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	403	EPE	C5-N4-C3	5.40	120.48	108.84
3	A	403	EPE	C6-N1-C2	4.91	119.41	108.84
3	A	402	EPE	C5-N4-C3	4.43	118.38	108.84
3	A	402	EPE	C6-N1-C2	4.21	117.91	108.84
3	A	402	EPE	C9-N1-C6	3.93	121.71	111.24
3	A	402	EPE	C7-N4-C5	3.58	120.78	111.24
3	A	402	EPE	C9-N1-C2	3.43	120.37	111.24
3	A	403	EPE	C7-N4-C5	3.37	120.23	111.24
3	A	403	EPE	C7-N4-C3	2.90	118.95	111.24
3	A	403	EPE	O2S-S-C10	2.75	110.89	106.73
3	A	402	EPE	O1S-S-C10	2.65	110.73	106.73
3	A	403	EPE	C10-C9-N1	-2.47	103.00	112.36
3	A	403	EPE	O3S-S-O1S	-2.31	105.62	111.40
2	C	401	MLI	O9-C3-C1	2.15	121.16	114.51

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	402	EPE	C10-C9-N1-C2
3	A	402	EPE	C9-C10-S-O2S
3	A	402	EPE	C9-C10-S-O3S
3	A	402	EPE	N4-C7-C8-O8
3	A	403	EPE	C8-C7-N4-C5
3	A	403	EPE	C10-C9-N1-C6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	402	EPE	C9-C10-S-O1S
3	A	403	EPE	N4-C7-C8-O8
3	A	403	EPE	C10-C9-N1-C2
2	A	401	MLI	C3-C1-C2-O7
2	A	401	MLI	C3-C1-C2-O6
2	C	401	MLI	C2-C1-C3-O8

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	EPE	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	316/345 (91%)	-0.72	2 (0%) 85 88	8, 16, 38, 63	0
1	B	317/345 (91%)	-0.74	0 100 100	7, 16, 36, 46	1 (0%)
1	C	318/345 (92%)	-0.68	0 100 100	6, 19, 37, 49	3 (0%)
All	All	951/1035 (91%)	-0.71	2 (0%) 91 92	6, 17, 37, 63	4 (0%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	65	ALA	2.5
1	A	345	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EPE	A	403	15/15	0.91	0.11	24,37,55,58	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	EPE	A	402	15/15	0.94	0.08	16,33,50,55	0
2	MLI	C	401	7/7	0.99	0.04	10,11,13,13	0
2	MLI	A	401	7/7	0.99	0.03	5,7,9,9	0
2	MLI	B	401	7/7	0.99	0.03	6,8,9,10	0

6.5 Other polymers [i](#)

There are no such residues in this entry.