



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 02:17 PM UTC

PDB ID : 6MD9 / pdb_00006md9
Title : NON-RECEPTOR PROTEIN TYROSINE PHOSPHATASE SHP2 IN COM-
PLEX WITH ALLOSTERIC INHIBITOR Isoxazolo-pyridinone 3
Authors : Fodor, M.; Stams, T.
Deposited on : 2018-09-04
Resolution : 2.12 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8290 (2.14-2.10)
Clashscore	190562	8817 (2.14-2.10)
Ramachandran outliers	187476	8738 (2.14-2.10)
Sidechain outliers	187428	8739 (2.14-2.10)
RSRZ outliers	180081	8294 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	526	7% 80% 9% 11%
1	B	526	6% 77% 11% 11%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8372 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein phosphatase non-receptor type 11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	468	Total	C	N	O	S	0	1	0
			3802	2401	674	709	18			
1	B	470	Total	C	N	O	S	0	4	0
			3838	2426	681	711	20			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q06124
B	0	SER	-	expression tag	UNP Q06124

- Molecule 2 is 3-(2-chlorophenyl)-6-{4-[(dimethylamino)methyl]phenyl}-5-methyl[1,2]oxazol o[4,5-c]pyridin-4(5H)-one (CCD ID: JEJ) (formula: C₂₂H₂₀ClN₃O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Cl	N	O	0	0
			28	22	1	3	2		
2	B	1	Total	C	Cl	N	O	0	0
			28	22	1	3	2		

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	333	Total	O	0	0
			333	333		
4	B	328	Total	O	0	0
			328	328		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein phosphatase non-receptor type 11



- Molecule 1: Tyrosine-protein phosphatase non-receptor type 11



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	46.34Å 213.96Å 55.83Å 90.00° 96.69° 90.00°	Depositor
Resolution (Å)	21.40 – 2.12 21.40 – 2.12	Depositor EDS
% Data completeness (in resolution range)	99.4 (21.40-2.12) 99.3 (21.40-2.12)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.69 (at 2.13Å)	Xtriage
Refinement program	BUSTER	Depositor
R, R_{free}	0.218 , 0.272 0.225 , 0.278	Depositor DCC
R_{free} test set	3045 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	28.8	Xtriage
Anisotropy	0.632	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 58.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8372	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, JEJ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	1/3881 (0.0%)	1.28	9/5230 (0.2%)
1	B	0.88	1/3925 (0.0%)	1.29	11/5286 (0.2%)
All	All	0.87	2/7806 (0.0%)	1.29	20/10516 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	306	ASN	CA-C	5.84	1.59	1.53
1	A	490	VAL	CA-C	5.17	1.58	1.52

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	64	ASP	CA-CB-CG	6.08	118.68	112.60
1	B	432	PRO	CA-C-N	5.85	126.60	119.99
1	B	432	PRO	C-N-CA	5.85	126.60	119.99
1	A	106	ASP	CA-CB-CG	5.82	118.42	112.60
1	A	396	TYR	N-CA-C	5.46	116.28	108.74
1	B	358	LYS	N-CA-C	-5.45	103.19	110.55
1	A	473	ASP	CA-CB-CG	-5.35	107.25	112.60
1	B	275	ASN	CA-CB-CG	5.16	117.76	112.60
1	A	431	ASP	CA-CB-CG	5.13	117.73	112.60
1	A	303	ASP	CA-CB-CG	5.12	117.72	112.60
1	A	432	PRO	CA-C-N	5.12	125.78	119.99
1	A	432	PRO	C-N-CA	5.12	125.78	119.99
1	B	304	TYR	N-CA-C	5.11	118.55	109.96
1	B	118	SER	CA-C-N	5.08	125.72	120.03
1	B	118	SER	C-N-CA	5.08	125.72	120.03
1	A	304	TYR	N-CA-C	5.03	117.68	109.59
1	B	16	ALA	CA-C-N	5.02	126.97	120.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	16	ALA	C-N-CA	5.02	126.97	120.44
1	B	519	GLN	CA-C-N	5.01	126.99	120.28
1	B	519	GLN	C-N-CA	5.01	126.99	120.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3802	0	3749	13	0
1	B	3838	0	3811	17	0
2	A	28	0	0	0	0
2	B	28	0	0	0	0
3	A	5	0	0	0	0
3	B	10	0	0	0	0
4	A	333	0	0	2	0
4	B	328	0	0	0	0
All	All	8372	0	7560	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:133:GLY:HA3	1:B:213:LYS:HB2	1.73	0.69
1:A:390:GLU:HG2	1:A:399:ARG:HG2	1.76	0.67
1:B:271:GLN:HA	1:B:274:LYS:HE2	1.82	0.60
1:A:256:GLN:HG3	4:A:846:HOH:O	2.04	0.58
1:A:133:GLY:HA3	1:A:213:LYS:HB2	1.84	0.57
1:A:309:ILE:HD13	1:A:328:ILE:HG12	1.91	0.53
1:B:264:SER:HB2	1:B:266:LYS:HD3	1.91	0.53
1:A:265:ARG:HH11	1:A:265:ARG:HB2	1.73	0.53
1:B:4:ARG:HD3	1:B:256:GLN:HA	1.90	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:290:VAL:HG11	1:B:344:MET:HG3	1.91	0.52
1:B:17:GLU:HG3	1:B:51:VAL:HB	1.92	0.50
1:B:129:LYS:HZ2	1:B:218:THR:HG22	1.77	0.49
1:A:125:LEU:HB3	1:A:216:LEU:HD21	1.94	0.48
1:B:440:GLU:HG3	1:B:444:HIS:CE1	2.49	0.47
1:A:41:PHE:HB2	1:A:56:ILE:HB	1.96	0.47
1:B:117:LEU:O	1:B:138:ARG:HD2	2.14	0.47
1:B:12:THR:HG22	1:B:35:LYS:HE3	1.98	0.46
1:B:129:LYS:HB2	1:B:216:LEU:HD11	1.97	0.45
1:B:222:ASN:O	1:B:225:GLU:HG2	2.17	0.45
1:A:514:ILE:O	1:A:518:VAL:HG23	2.17	0.45
1:A:4:ARG:HD3	1:A:256:GLN:HA	2.00	0.44
1:A:498:ARG:HD3	4:A:973:HOH:O	2.18	0.43
1:A:65:LEU:HD23	1:A:68:GLY:HA3	2.00	0.43
1:A:284:PRO:HG3	1:A:306:ASN:HA	2.00	0.43
1:B:126:LEU:HD22	1:B:152:ARG:HB2	2.00	0.42
1:A:107:PRO:HG3	1:A:190:LEU:HD12	2.03	0.41
1:B:525:LEU:HD12	1:B:525:LEU:HA	1.93	0.41
1:B:6:TRP:HB3	1:B:101:PRO:HB3	2.02	0.40
1:B:134:SER:HA	1:B:214:GLN:O	2.21	0.40
1:B:251:PHE:O	1:B:255:GLN:HG2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	453/526 (86%)	435 (96%)	17 (4%)	1 (0%)	43	44
1	B	458/526 (87%)	446 (97%)	10 (2%)	2 (0%)	30	28
All	All	911/1052 (87%)	881 (97%)	27 (3%)	3 (0%)	36	36

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	505	VAL
1	A	505	VAL
1	B	115	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	417/468 (89%)	401 (96%)	16 (4%)	29	30
1	B	423/468 (90%)	397 (94%)	26 (6%)	17	14
All	All	840/936 (90%)	798 (95%)	42 (5%)	23	21

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	102	LEU
1	A	151	VAL
1	A	152	ARG
1	A	165	SER
1	A	171	MET
1	A	206	LEU
1	A	221	ILE
1	A	226	ILE
1	A	265	ARG
1	A	281	ASN
1	A	408	GLN
1	A	439	LEU
1	A	447	GLU
1	A	484	VAL
1	A	485	ASP
1	A	502	SER
1	B	11	ILE
1	B	18[A]	ASN
1	B	18[B]	ASN
1	B	37	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	43	LEU
1	B	47	ARG
1	B	52	THR
1	B	70	LYS
1	B	89	LYS
1	B	96	ILE
1	B	118	SER
1	B	121	GLU
1	B	149	LEU
1	B	151	VAL
1	B	166	LYS
1	B	171	MET
1	B	249	GLU
1	B	265[A]	ARG
1	B	265[B]	ARG
1	B	275	ASN
1	B	365	SER
1	B	421	ARG
1	B	426	HIS
1	B	451	ASP
1	B	508	GLU
1	B	525	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	10	ASN
1	A	222	ASN
1	A	256	GLN
1	A	257	GLN
1	A	394	HIS
1	A	426	HIS
1	B	57	GLN
1	B	87	GLN
1	B	222	ASN
1	B	257	GLN
1	B	281	ASN
1	B	444	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	JEJ	A	601	-	31,31,31	1.46	6 (19%)	35,45,45	1.67	7 (20%)
3	PO4	B	603	-	4,4,4	1.97	1 (25%)	6,6,6	0.58	0
3	PO4	B	602	-	4,4,4	2.56	3 (75%)	6,6,6	0.60	0
3	PO4	A	602	-	4,4,4	2.60	3 (75%)	6,6,6	0.55	0
2	JEJ	B	601	-	31,31,31	1.57	6 (19%)	35,45,45	1.66	8 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	JEJ	B	601	-	-	0/12/12/12	0/4/4/4
2	JEJ	A	601	-	-	2/12/12/12	0/4/4/4

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	602	PO4	P-O1	4.29	1.60	1.50
3	B	602	PO4	P-O1	4.23	1.60	1.50
2	B	601	JEJ	C9-N2	3.51	1.46	1.39
2	A	601	JEJ	C9-N2	3.05	1.45	1.39
2	A	601	JEJ	C5-C22	3.03	1.43	1.39
2	B	601	JEJ	C5-C22	2.87	1.43	1.39
2	B	601	JEJ	C14-C13	2.47	1.43	1.39
2	B	601	JEJ	C11-C10	2.44	1.50	1.44
2	A	601	JEJ	C11-C10	2.24	1.49	1.44
2	A	601	JEJ	C15-C16	2.18	1.43	1.38
2	A	601	JEJ	C18-C13	2.14	1.42	1.39
2	B	601	JEJ	C18-C17	2.13	1.42	1.38
2	A	601	JEJ	C15-C14	2.11	1.42	1.38
2	B	601	JEJ	C15-C14	2.07	1.42	1.38
3	B	602	PO4	P-O2	2.03	1.60	1.54
3	B	603	PO4	P-O2	2.02	1.60	1.54
3	A	602	PO4	P-O3	2.01	1.60	1.54
3	B	602	PO4	P-O4	2.01	1.60	1.54
3	A	602	PO4	P-O4	2.01	1.60	1.54

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	JEJ	C11-C6-N1	-3.67	107.56	111.43
2	A	601	JEJ	C7-O1-N1	-3.65	104.33	108.38
2	A	601	JEJ	O1-N1-C6	3.64	109.79	105.91
2	B	601	JEJ	C11-C6-N1	-3.58	107.66	111.43
2	A	601	JEJ	O2-C10-N2	3.51	126.78	120.36
2	B	601	JEJ	O1-N1-C6	3.33	109.46	105.91
2	B	601	JEJ	C7-O1-N1	-3.25	104.77	108.38
2	A	601	JEJ	C5-C6-N1	3.16	122.29	118.23
2	B	601	JEJ	C16-C19-N3	3.07	118.93	113.17
2	B	601	JEJ	O2-C10-N2	3.06	125.96	120.36
2	A	601	JEJ	O2-C10-C11	-3.01	118.72	125.16
2	A	601	JEJ	C13-C9-N2	2.69	124.16	118.56
2	B	601	JEJ	O2-C10-C11	-2.63	119.53	125.16
2	B	601	JEJ	C13-C9-N2	2.55	123.87	118.56
2	B	601	JEJ	C5-C6-N1	2.30	121.18	118.23

There are no chirality outliers.


All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	601	JEJ	C16-C19-N3-C20
2	A	601	JEJ	C16-C19-N3-C21

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	468/526 (88%)	0.70	35 (7%)	20 22	17, 36, 59, 91	1 (0%)
1	B	470/526 (89%)	0.67	31 (6%)	24 26	14, 36, 58, 73	4 (0%)
All	All	938/1052 (89%)	0.68	66 (7%)	22 24	14, 36, 59, 91	5 (0%)

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	205	THR	5.0
1	B	484	VAL	4.1
1	B	190	LEU	3.9
1	B	248	TRP	3.9
1	B	301	VAL	3.9
1	A	144	PRO	3.6
1	B	144	PRO	3.4
1	A	165	SER	3.4
1	A	525	LEU	3.3
1	A	120	LYS	3.2
1	B	426	HIS	3.1
1	A	248	TRP	3.1
1	A	119	GLY	3.1
1	B	171	MET	3.1
1	A	236	LEU	3.0
1	A	203	VAL	3.0
1	A	225	GLU	3.0
1	A	437	ASP	3.0
1	A	484	VAL	2.9
1	B	246	GLY	2.9
1	B	206	LEU	2.8
1	B	522	ILE	2.8
1	A	118	SER	2.8
1	B	167	VAL	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	171	MET	2.8
1	B	96	ILE	2.8
1	A	11	ILE	2.7
1	B	177	LEU	2.7
1	B	108	THR	2.6
1	B	451	ASP	2.6
1	A	34	SER	2.6
1	B	182	GLY	2.6
1	A	38	PRO	2.6
1	A	522	ILE	2.5
1	A	66	TYR	2.5
1	A	179	TYR	2.5
1	B	224	ALA	2.5
1	A	469	PHE	2.5
1	A	52	THR	2.5
1	A	106	ASP	2.4
1	B	118	SER	2.4
1	B	256	GLN	2.3
1	B	221	ILE	2.3
1	B	488	ILE	2.3
1	A	408	GLN	2.3
1	B	213	LYS	2.3
1	A	117	LEU	2.3
1	A	206	LEU	2.3
1	B	482	LYS	2.3
1	B	275	ASN	2.3
1	A	311	MET	2.3
1	B	525	LEU	2.3
1	A	127	THR	2.2
1	A	488	ILE	2.2
1	A	170	VAL	2.2
1	B	235	LYS	2.2
1	A	482	LYS	2.2
1	B	188	ASP	2.2
1	A	523	GLU	2.1
1	A	182	GLY	2.1
1	A	302	SER	2.1
1	B	165	SER	2.1
1	B	293	HIS	2.1
1	B	139	GLU	2.0
1	A	155	ASP	2.0
1	B	124	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PO4	B	603	5/5	0.61	0.26	103,105,105,106	0
3	PO4	A	602	5/5	0.81	0.12	68,69,70,71	0
2	JEJ	B	601	28/28	0.85	0.13	29,40,56,57	0
2	JEJ	A	601	28/28	0.88	0.12	31,38,50,53	0
3	PO4	B	602	5/5	0.92	0.08	46,46,46,50	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around JEJ B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around JEJ A 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.