



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 12:04 PM UTC

PDB ID : 7MDJ / pdb_00007mdj
Title : The structure of KcsA in complex with a synthetic Fab
Authors : Rohaim, A.; Slezak, T.; Blackowicz, L.; Kossiakoff, A.; Roux, B.
Deposited on : 2021-04-05
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	229	8% 60% 28% 5% 7%
2	B	215	3% 56% 31% 5% 9%
3	C	124	10% 59% 21% 18%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4005 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	214	Total	C	N	O	S	0	0	0
			1599	1010	269	315	5			

- Molecule 2 is a protein called Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	196	Total	C	N	O	S	0	0	0
			1516	950	255	306	5			

- Molecule 3 is a protein called pH-gated potassium channel KcsA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	102	Total	C	N	O	S	0	0	0
			759	498	129	130	2			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	2	ALA	PRO	conflict	UNP P0A334
C	90	CYS	LEU	conflict	UNP P0A334

- Molecule 4 is POTASSIUM ION (CCD ID: K) (formula: K) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	4	Total	K	0	0
			4	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	59	Total 59	O 59	0	0
5	B	42	Total 42	O 42	0	0
5	C	26	Total 26	O 26	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fab heavy chain



- Molecule 2: Fab light chain



- Molecule 3: pH-gated potassium channel KcsA



4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	138.41Å 138.41Å 71.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.01 – 2.75 50.01 – 2.75	Depositor EDS
% Data completeness (in resolution range)	99.7 (50.01-2.75) 99.7 (50.01-2.75)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.31 (at 2.73Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.201 , 0.265 0.217 , 0.298	Depositor DCC
R_{free} test set	835 reflections (4.73%)	wwPDB-VP
Wilson B-factor (Å ²)	103.5	Xtriage
Anisotropy	0.174	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 101.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.029 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4005	wwPDB-VP
Average B, all atoms (Å ²)	131.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.05	1/1638 (0.1%)	1.36	4/2231 (0.2%)
2	B	1.12	0/1548	1.46	4/2099 (0.2%)
3	C	1.09	0/778	1.71	4/1070 (0.4%)
All	All	1.09	1/3964 (0.0%)	1.48	12/5400 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	42	GLN	C-O	5.02	1.30	1.23

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	58	TYR	N-CA-C	-8.37	103.05	113.18
2	B	2	ASP	CA-CB-CG	6.73	119.33	112.60
2	B	73	THR	CA-CB-OG1	-6.61	99.68	109.60
3	C	80	ASP	CA-CB-CG	6.49	119.09	112.60
1	A	41	ARG	CB-CA-C	5.47	119.38	109.37
3	C	61	THR	CA-C-N	5.32	128.84	120.97
3	C	61	THR	C-N-CA	5.32	128.84	120.97
1	A	180	SER	CA-C-N	5.24	127.30	120.28
1	A	180	SER	C-N-CA	5.24	127.30	120.28
3	C	71	GLU	N-CA-C	-5.12	105.82	111.71
2	B	121	PRO	CA-C-N	5.07	128.04	120.90
2	B	121	PRO	C-N-CA	5.07	128.04	120.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1599	0	1546	34	0
2	B	1516	0	1477	47	0
3	C	759	0	768	30	0
4	C	4	0	0	0	0
5	A	59	0	0	3	0
5	B	42	0	0	12	0
5	C	26	0	0	9	0
All	All	4005	0	3791	100	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (100) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:122:ARG:CD	5:C:301:HOH:O	1.91	1.15
3:C:122:ARG:NE	5:C:301:HOH:O	1.84	1.10
3:C:22:SER:N	3:C:25:HIS:CD2	2.22	1.07
1:A:55:ILE:HD11	3:C:49:LEU:HD13	1.65	0.78
2:B:140:PHE:CZ	5:B:325:HOH:O	2.36	0.78
3:C:122:ARG:HD3	5:C:301:HOH:O	1.66	0.75
2:B:34:ILE:HG21	2:B:72:PHE:CD1	2.24	0.72
1:A:5:VAL:HA	1:A:29:GLY:HA3	1.75	0.69
3:C:122:ARG:CZ	5:C:301:HOH:O	2.29	0.67
3:C:25:HIS:CD2	3:C:112:THR:HB	2.31	0.66
3:C:119:GLN:NE2	5:C:303:HOH:O	2.30	0.65
2:B:63:PHE:HB3	5:B:333:HOH:O	1.96	0.64
1:A:197:LEU:HD23	1:A:198:GLY:HA2	1.80	0.63
2:B:5:MET:HE3	2:B:24:CYS:SG	2.41	0.61
1:A:22:ARG:HB2	1:A:85:GLN:HE22	1.67	0.59
2:B:2:ASP:N	5:B:302:HOH:O	2.35	0.59
1:A:67:VAL:HG13	1:A:71:PHE:HB2	1.85	0.58
1:A:162:TRP:CH2	1:A:204:CYS:HB3	2.38	0.58
1:A:41:ARG:NE	1:A:51:ILE:HD11	2.19	0.58
1:A:20:SER:HA	1:A:86:MET:O	2.03	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:219:VAL:HA	5:A:317:HOH:O	2.04	0.57
2:B:114:PRO:HA	2:B:138:ASN:O	2.04	0.57
2:B:81:PRO:O	2:B:107:ILE:HD11	2.03	0.57
2:B:158:GLY:C	2:B:160:SER:N	2.63	0.57
2:B:13:SER:HA	2:B:106:GLU:O	2.05	0.57
2:B:143:ARG:NH2	2:B:164:VAL:HG21	2.20	0.56
2:B:50:LYS:HG2	2:B:54:GLU:HB2	1.88	0.56
2:B:94:ARG:HD3	3:C:58:GLN:HG3	1.88	0.56
2:B:19:ARG:HG3	5:B:303:HOH:O	2.04	0.56
2:B:144:GLU:OE2	2:B:146:LYS:NZ	2.40	0.55
2:B:36:TRP:CD1	5:B:311:HOH:O	2.59	0.55
1:A:36:TRP:CZ3	3:C:62:TYR:OH	2.57	0.55
2:B:91:GLN:HE22	2:B:94:ARG:H	1.53	0.55
3:C:25:HIS:HE1	3:C:26:TRP:CE2	2.25	0.55
2:B:94:ARG:HD2	3:C:56:GLY:O	2.07	0.54
2:B:150:LYS:HG2	2:B:155:LEU:HA	1.90	0.54
2:B:106:GLU:OE1	2:B:174:TYR:OH	2.26	0.53
1:A:35:ASP:OD2	1:A:101:ARG:NH1	2.41	0.53
2:B:48:LEU:HB3	5:B:333:HOH:O	2.07	0.53
2:B:38:GLN:NE2	5:B:301:HOH:O	2.29	0.52
1:A:131:PRO:HD3	1:A:217:LYS:HD2	1.92	0.52
1:A:162:TRP:CZ3	1:A:204:CYS:HB3	2.44	0.52
1:A:55:ILE:CD1	3:C:49:LEU:HD13	2.39	0.52
3:C:64:ARG:HD2	5:C:313:HOH:O	2.10	0.52
1:A:149:LEU:HD13	1:A:151:LYS:HD2	1.92	0.51
3:C:22:SER:N	3:C:25:HIS:NE2	2.57	0.51
1:A:36:TRP:HZ3	3:C:62:TYR:CZ	2.28	0.51
2:B:112:ALA:O	2:B:140:PHE:CD1	2.64	0.51
3:C:44:SER:OG	3:C:66:LEU:HA	2.11	0.50
3:C:82:TYR:HB2	3:C:83:PRO:HD2	1.92	0.50
1:A:189:VAL:HG11	2:B:136:LEU:HD11	1.94	0.50
1:A:90:ARG:O	1:A:119:VAL:HG11	2.12	0.49
3:C:62:TYR:N	3:C:63:PRO:CD	2.76	0.49
1:A:211:SER:O	1:A:212:ASN:C	2.55	0.49
1:A:186:LEU:HB2	5:A:316:HOH:O	2.12	0.49
2:B:38:GLN:HG3	2:B:87:TYR:CE2	2.47	0.49
2:B:152:ASP:OD2	2:B:190:HIS:HB3	2.13	0.49
2:B:140:PHE:CE1	5:B:325:HOH:O	2.61	0.49
2:B:94:ARG:CD	3:C:56:GLY:O	2.62	0.48
2:B:119:PHE:HB2	2:B:134:VAL:HB	1.96	0.48
2:B:133:VAL:HG12	2:B:149:TRP:CZ3	2.49	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:12:LEU:HD21	5:B:320:HOH:O	2.14	0.47
3:C:78:TYR:N	5:C:304:HOH:O	2.47	0.47
3:C:62:TYR:HB2	3:C:63:PRO:HD3	1.96	0.47
1:A:63:TYR:HE1	1:A:73:ILE:HG13	1.80	0.46
1:A:94:THR:HG23	1:A:118:THR:HA	1.97	0.46
2:B:126:LEU:HD11	2:B:187:TYR:HE2	1.81	0.46
1:A:209:LYS:HD3	1:A:209:LYS:HA	1.49	0.46
2:B:67:ARG:HB3	2:B:72:PHE:HA	1.97	0.46
3:C:66:LEU:HD12	3:C:69:SER:OG	2.17	0.45
2:B:140:PHE:HE2	2:B:144:GLU:O	1.99	0.45
3:C:118:GLU:HB3	5:C:314:HOH:O	2.16	0.45
1:A:38:HIS:HA	1:A:52:GLY:O	2.17	0.45
2:B:67:ARG:HD2	2:B:67:ARG:HA	1.63	0.45
1:A:214:LYS:HE3	1:A:214:LYS:HB2	1.53	0.45
3:C:64:ARG:HB2	5:C:306:HOH:O	2.16	0.45
2:B:81:PRO:HA	2:B:107:ILE:HD13	2.00	0.44
2:B:158:GLY:HA2	5:B:305:HOH:O	2.17	0.44
1:A:42:GLN:OE1	1:A:46:LYS:O	2.36	0.44
2:B:30:ILE:HD11	2:B:72:PHE:CE2	2.53	0.44
1:A:146:LEU:C	1:A:146:LEU:HD12	2.43	0.44
2:B:78:SER:N	5:B:303:HOH:O	2.46	0.44
1:A:197:LEU:HA	1:A:198:GLY:HA2	1.82	0.44
3:C:44:SER:HA	3:C:65:ALA:HB1	1.99	0.44
2:B:33:ASP:HB3	2:B:92:SER:OG	2.18	0.43
2:B:67:ARG:HE	2:B:68:SER:H	1.65	0.43
1:A:172:HIS:CD2	2:B:175:SER:OG	2.71	0.43
2:B:81:PRO:O	2:B:107:ILE:CD1	2.66	0.43
1:A:209:LYS:O	1:A:210:PRO:C	2.62	0.42
1:A:125:LYS:HD2	1:A:125:LYS:HA	1.44	0.42
1:A:186:LEU:HD12	1:A:186:LEU:H	1.83	0.42
5:B:306:HOH:O	3:C:55:PRO:HB3	2.18	0.42
2:B:94:ARG:HD2	3:C:56:GLY:C	2.44	0.41
2:B:133:VAL:CG1	2:B:149:TRP:CZ3	3.03	0.41
2:B:141:TYR:CD1	2:B:142:PRO:HA	2.55	0.41
2:B:176:LEU:HD23	2:B:176:LEU:C	2.45	0.41
1:A:152:ASP:OD1	5:A:301:HOH:O	2.21	0.41
3:C:75:THR:O	3:C:75:THR:OG1	2.38	0.41
1:A:104:GLY:HA3	3:C:62:TYR:CE1	2.56	0.41
2:B:164:VAL:HG22	2:B:176:LEU:HD12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	208/229 (91%)	186 (89%)	21 (10%)	1 (0%)	24	43
2	B	192/215 (89%)	171 (89%)	20 (10%)	1 (0%)	24	43
3	C	100/124 (81%)	94 (94%)	5 (5%)	1 (1%)	12	24
All	All	500/568 (88%)	451 (90%)	46 (9%)	3 (1%)	21	38

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	77	GLY
2	B	78	SER
1	A	134	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	175/190 (92%)	137 (78%)	38 (22%)	1	1
2	B	174/191 (91%)	144 (83%)	30 (17%)	2	3
3	C	72/91 (79%)	66 (92%)	6 (8%)	10	19
All	All	421/472 (89%)	347 (82%)	74 (18%)	2	3

All (74) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	GLN
1	A	15	VAL
1	A	16	GLN
1	A	24	SER
1	A	30	TYR
1	A	31	THR
1	A	32	PHE
1	A	33	THR
1	A	51	ILE
1	A	55	ILE
1	A	57	SER
1	A	58	TYR
1	A	65	ASP
1	A	67	VAL
1	A	68	LYS
1	A	74	SER
1	A	77	THR
1	A	78	SER
1	A	79	LYS
1	A	87	ASN
1	A	105	ASP
1	A	116	LEU
1	A	118	THR
1	A	120	SER
1	A	121	SER
1	A	125	LYS
1	A	159	THR
1	A	168	THR
1	A	186	LEU
1	A	189	VAL
1	A	190	VAL
1	A	192	VAL
1	A	194	SER
1	A	199	THR
1	A	200	GLN
1	A	209	LYS
1	A	214	LYS
1	A	218	LYS
2	B	6	THR
2	B	8	SER
2	B	16	VAL
2	B	19	ARG
2	B	21	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	23	THR
2	B	43	LYS
2	B	50	LYS
2	B	67	ARG
2	B	68	SER
2	B	74	LEU
2	B	77	SER
2	B	82	GLU
2	B	91	GLN
2	B	104	LYS
2	B	110	THR
2	B	115	SER
2	B	143	ARG
2	B	147	VAL
2	B	148	GLN
2	B	150	LYS
2	B	151	VAL
2	B	169	SER
2	B	170	LYS
2	B	184	LYS
2	B	186	ASP
2	B	190	HIS
2	B	191	LYS
2	B	192	VAL
2	B	198	THR
3	C	24	LEU
3	C	25	HIS
3	C	71	GLU
3	C	105	LEU
3	C	115	VAL
3	C	119	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	GLN
1	A	42	GLN
1	A	85	GLN
1	A	87	ASN
1	A	172	HIS
1	A	179	GLN
1	A	200	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	39	GLN
2	B	80	GLN
2	B	90	GLN
2	B	91	GLN
2	B	93	ASN
2	B	139	ASN
2	B	148	GLN
3	C	25	HIS
3	C	58	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	214/229 (93%)	0.31	18 (8%) 17 17	88, 141, 191, 217	0
2	B	196/215 (91%)	-0.10	7 (3%) 46 44	78, 124, 167, 205	0
3	C	102/124 (82%)	0.45	12 (11%) 9 7	73, 103, 203, 236	0
All	All	512/568 (90%)	0.18	37 (7%) 21 21	73, 129, 187, 236	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	61	ALA	4.8
3	C	123	GLY	4.2
1	A	77	THR	4.1
2	B	133	VAL	4.0
3	C	49	LEU	4.0
1	A	39	TRP	4.0
1	A	51	ILE	3.5
1	A	35	ASP	3.3
3	C	87	TRP	3.2
1	A	63	TYR	3.2
2	B	158	GLY	3.1
1	A	195	SER	3.1
1	A	97	TYR	3.1
2	B	193	TYR	3.1
3	C	54	ALA	3.1
2	B	198	THR	3.0
2	B	145	ALA	2.8
1	A	100	ALA	2.8
1	A	78	SER	2.7
1	A	56	PRO	2.7
1	A	171	VAL	2.6
2	B	197	VAL	2.6
3	C	113	TRP	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	23	ALA	2.4
3	C	50	ALA	2.4
3	C	22	SER	2.4
1	A	37	ILE	2.4
1	A	68	LYS	2.4
1	A	36	TRP	2.3
3	C	40	LEU	2.3
3	C	24	LEU	2.3
2	B	169	SER	2.3
1	A	52	GLY	2.2
1	A	197	LEU	2.2
3	C	103	PHE	2.2
1	A	73	ILE	2.0
3	C	104	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	K	C	204	1/1	0.93	0.23	127,127,127,127	1
4	K	C	202	1/1	0.96	0.07	143,143,143,143	1
4	K	C	203	1/1	1.00	0.02	137,137,137,137	1
4	K	C	201	1/1	1.00	0.04	109,109,109,109	1


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around K C 204:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)









6.5 Other polymers ⓘ

There are no such residues in this entry.