



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 7, 2026 – 12:21 AM UTC

PDB ID : 5MEN / pdb_00005men
Title : Human Leukocyte Antigen A02 presenting ILAKFLHWL, in complex with cognate T-Cell Receptor
Authors : Rizkallah, P.J.; Lloyd, A.; Crowther, M.; Cole, D.K.; Sewell, A.K.
Deposited on : 2016-11-16
Resolution : 2.81 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4591 (2.84-2.80)
Clashscore	190562	5010 (2.84-2.80)
Ramachandran outliers	187476	4916 (2.84-2.80)
Sidechain outliers	187428	4918 (2.84-2.80)
RSRZ outliers	180081	4594 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	 69% 26% .
2	B	100	 83% 15% .
3	C	9	 56% 44%
4	D	200	 62% 28% 10%
5	E	240	 2% 72% 20% 6% .

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 6739 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, A-2 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	276	2254	1408	410	427	9	0	0	0

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	100	837	533	141	159	4	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called ILE-LEU-ALA-LYS-PHE-LEU-HIS-TRP-LEU.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	9	82	59	13	10	0	0	0

- Molecule 4 is a protein called Protein TRAV22,Human nkt tcr alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	200	1562	978	259	318	7	0	0	0

There are 29 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	90	ASP	-	linker	UNP A0A0B4J277
D	91	SER	-	linker	UNP A0A0B4J277
D	92	ALA	-	linker	UNP A0A0B4J277

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	93	THR	-	linker	UNP A0A0B4J277
D	94	SER	-	linker	UNP A0A0B4J277
D	95	GLY	-	linker	UNP A0A0B4J277
D	96	THR	-	linker	UNP A0A0B4J277
D	97	TYR	-	linker	UNP A0A0B4J277
D	98	LYS	-	linker	UNP A0A0B4J277
D	99	TYR	-	linker	UNP A0A0B4J277
D	100	ILE	-	linker	UNP A0A0B4J277
D	101	PHE	-	linker	UNP A0A0B4J277
D	102	GLY	-	linker	UNP A0A0B4J277
D	103	THR	-	linker	UNP A0A0B4J277
D	104	GLY	-	linker	UNP A0A0B4J277
D	105	THR	-	linker	UNP A0A0B4J277
D	106	ARG	-	linker	UNP A0A0B4J277
D	107	LEU	-	linker	UNP A0A0B4J277
D	108	LYS	-	linker	UNP A0A0B4J277
D	109	VAL	-	linker	UNP A0A0B4J277
D	110	LEU	-	linker	UNP A0A0B4J277
D	111	ALA	-	linker	UNP A0A0B4J277
D	112	ASN	-	linker	UNP A0A0B4J277
D	113	ILE	ILE	linker	UNP K7N5M3
D	114	GLN	GLN	linker	UNP K7N5M3
D	115	ASN	ASN	linker	UNP K7N5M3
D	116	PRO	PRO	linker	UNP K7N5M3
D	117	ASP	ASP	linker	UNP K7N5M3
D	118	PRO	PRO	linker	UNP K7N5M3

- Molecule 5 is a protein called Protein TRBV6-5, Human nkt tcr beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	E	240	1894	1190	327	368	9	0	0	0

There are 22 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
E	96	GLN	-	linker	UNP A0A0K0K1A5
E	97	GLY	-	linker	UNP A0A0K0K1A5
E	98	THR	-	linker	UNP A0A0K0K1A5
E	99	GLU	-	linker	UNP A0A0K0K1A5
E	100	ALA	-	linker	UNP A0A0K0K1A5
E	101	PHE	-	linker	UNP A0A0K0K1A5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	102	PHE	-	linker	UNP A0A0K0K1A5
E	103	GLY	-	linker	UNP A0A0K0K1A5
E	104	GLN	-	linker	UNP A0A0K0K1A5
E	105	GLY	-	linker	UNP A0A0K0K1A5
E	106	THR	-	linker	UNP A0A0K0K1A5
E	107	ARG	-	linker	UNP A0A0K0K1A5
E	108	LEU	-	linker	UNP A0A0K0K1A5
E	109	THR	-	linker	UNP A0A0K0K1A5
E	110	VAL	-	linker	UNP A0A0K0K1A5
E	111	VAL	-	linker	UNP A0A0K0K1A5
E	112	GLU	-	linker	UNP A0A0K0K1A5
E	113	ASP	-	linker	UNP A0A0K0K1A5
E	114	LEU	-	linker	UNP A0A0K0K1A5
E	115	ASN	-	linker	UNP A0A0K0K1A5
E	116	LYS	-	linker	UNP A0A0K0K1A5
E	200	ASP	ASN	conflict	UNP K7N5M4

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf		
6	A	1	Total	C	O	0	0
			4	2	2		
			Total	C	O		
6	A	1	Total	C	O	0	0
			4	2	2		
			Total	C	O		
6	B	1	Total	C	O	0	0
			4	2	2		
			Total	C	O		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	O	S	0	0
			5	4	1		
8	A	1	Total	O	S	0	0
			5	4	1		
8	A	1	Total	O	S	0	0
			5	4	1		
8	B	1	Total	O	S	0	0
			5	4	1		
8	E	1	Total	O	S	0	0
			5	4	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	23	Total	O	0	0
			23	23		
9	B	13	Total	O	0	0
			13	13		
9	C	1	Total	O	0	0
			1	1		
9	D	11	Total	O	0	0
			11	11		
9	E	11	Total	O	0	0
			11	11		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HLA class I histocompatibility antigen, A-2 alpha chain




- Molecule 2: Beta-2-microglobulin



- Molecule 3: ILE-LEU-ALA-LYS-PHE-LEU-HIS-TRP-LEU




- Molecule 4: Protein TRAV22,Human nkt tcr alpha chain



S201

- Molecule 5: Protein TRBV6-5, Human nkt tcr beta chain

Chain E:  72% 20% 6% 2%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	93.16Å 48.69Å 118.10Å 90.00° 108.15° 90.00°	Depositor
Resolution (Å)	38.03 – 2.81 38.03 – 2.81	Depositor EDS
% Data completeness (in resolution range)	99.6 (38.03-2.81) 99.6 (38.03-2.81)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.07 (at 2.81Å)	Xtrriage
Refinement program	REFMAC 5.8.0107	Depositor
R, R_{free}	0.189 , 0.272 0.199 , 0.197	Depositor DCC
R_{free} test set	1274 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	55.3	Xtrriage
Anisotropy	0.296	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 39.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6739	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.18	5/2320 (0.2%)	1.27	15/3149 (0.5%)
2	B	1.25	1/860 (0.1%)	1.14	0/1162
3	C	1.17	0/85	1.06	0/113
4	D	1.10	0/1596	1.24	10/2171 (0.5%)
5	E	1.13	1/1945 (0.1%)	1.19	10/2649 (0.4%)
All	All	1.16	7/6806 (0.1%)	1.22	35/9244 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
4	D	0	1
5	E	0	3
All	All	0	6

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	24	ALA	CA-C	-6.67	1.44	1.52
1	A	87	GLN	CA-C	-6.07	1.45	1.52
5	E	60	VAL	CA-C	6.04	1.57	1.52
1	A	105	SER	N-CA	5.76	1.53	1.46
1	A	93	HIS	CA-C	-5.28	1.46	1.52

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	236	ALA	N-CA-C	-8.56	98.42	110.50
1	A	14	ARG	CA-C-N	6.69	128.20	119.84
1	A	14	ARG	C-N-CA	6.69	128.20	119.84
1	A	106	ASP	N-CA-C	-6.49	100.81	110.23
1	A	35	ARG	N-CA-C	6.47	118.91	109.07

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	105	SER	Peptide
1	A	175	GLY	Peptide
4	D	127	LYS	Peptide
5	E	56	ASP	Peptide
5	E	57	GLN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2254	0	2103	24	0
2	B	837	0	803	7	0
3	C	82	0	90	5	0
4	D	1562	0	1480	32	0
5	E	1894	0	1787	34	0
6	A	8	0	12	0	0
6	B	12	0	18	0	0
7	A	6	0	8	0	0
8	A	15	0	0	0	0
8	B	5	0	0	0	0
8	E	5	0	0	0	0
9	A	23	0	0	0	0
9	B	13	0	0	0	0
9	C	1	0	0	0	0
9	D	11	0	0	0	0
9	E	11	0	0	0	0
All	All	6739	0	6301	93	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 93 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:159:CYS:SG	5:E:190:ARG:NH2	2.33	1.02
2:B:4:THR:HA	2:B:86:THR:HG21	1.53	0.89
1:A:107:TRP:O	1:A:169:ARG:NH1	2.13	0.81
1:A:67:VAL:HG13	3:C:2:LEU:HD11	1.69	0.73
4:D:32:LEU:HD22	4:D:89:VAL:HG22	1.69	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/276 (99%)	250 (91%)	22 (8%)	2 (1%)	18	45
2	B	98/100 (98%)	94 (96%)	4 (4%)	0	100	100
3	C	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
4	D	198/200 (99%)	179 (90%)	18 (9%)	1 (0%)	24	53
5	E	238/240 (99%)	207 (87%)	27 (11%)	4 (2%)	7	23
All	All	815/825 (99%)	736 (90%)	72 (9%)	7 (1%)	14	38

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
5	E	57	GLN
1	A	105	SER
4	D	15	GLU
5	E	62	ASN
5	E	227	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/232 (100%)	194 (84%)	38 (16%)	2	7
2	B	95/95 (100%)	86 (90%)	9 (10%)	8	25
3	C	8/8 (100%)	8 (100%)	0	100	100
4	D	180/180 (100%)	142 (79%)	38 (21%)	1	3
5	E	206/206 (100%)	173 (84%)	33 (16%)	2	8
All	All	721/721 (100%)	603 (84%)	118 (16%)	2	7

5 of 118 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	D	58	ARG
5	E	167	VAL
4	D	137	THR
5	E	163	VAL
5	E	107	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
4	D	78	GLN
5	E	222	GLN
4	D	189	ASN
5	E	181	ASN
4	D	147	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	GOL	A	303	-	5,5,5	0.35	0	5,5,5	1.11	0
6	EDO	B	303	-	3,3,3	0.54	0	2,2,2	0.26	0
6	EDO	A	301	-	3,3,3	0.46	0	2,2,2	0.22	0
6	EDO	A	302	-	3,3,3	0.62	0	2,2,2	0.15	0
6	EDO	B	302	-	3,3,3	0.56	0	2,2,2	0.22	0
6	EDO	B	301	-	3,3,3	0.44	0	2,2,2	0.38	0
8	SO4	A	304	-	4,4,4	0.49	0	6,6,6	0.19	0
8	SO4	A	305	-	4,4,4	0.45	0	6,6,6	0.18	0
8	SO4	A	306	-	4,4,4	0.49	0	6,6,6	0.10	0
8	SO4	E	301	-	4,4,4	0.47	0	6,6,6	0.17	0
8	SO4	B	304	-	4,4,4	0.46	0	6,6,6	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	A	303	-	-	2/4/4/4	-
6	EDO	B	303	-	-	1/1/1/1	-
6	EDO	A	301	-	-	0/1/1/1	-
6	EDO	A	302	-	-	1/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	B	302	-	-	1/1/1/1	-
6	EDO	B	301	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	303	GOL	C1-C2-C3-O3
7	A	303	GOL	O2-C2-C3-O3
6	A	302	EDO	O1-C1-C2-O2
6	B	303	EDO	O1-C1-C2-O2
6	B	301	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	276/276 (100%)	-0.45	0 100 100	25, 50, 88, 110	0
2	B	100/100 (100%)	-0.62	0 100 100	25, 41, 62, 71	0
3	C	9/9 (100%)	-0.86	0 100 100	34, 37, 39, 43	0
4	D	200/200 (100%)	-0.19	0 100 100	36, 64, 94, 112	0
5	E	240/240 (100%)	-0.27	5 (2%) 63 54	37, 60, 104, 128	0
All	All	825/825 (100%)	-0.36	5 (0%) 85 80	25, 56, 92, 128	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	E	61	PRO	3.7
5	E	179	ALA	3.4
5	E	180	LEU	2.7
5	E	240	ALA	2.4
5	E	60	VAL	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	SO4	A	305	5/5	0.84	0.10	120,125,126,129	0
8	SO4	A	306	5/5	0.86	0.15	108,112,115,116	0
7	GOL	A	303	6/6	0.91	0.11	39,45,45,46	0
8	SO4	B	304	5/5	0.92	0.09	83,84,88,90	0
8	SO4	E	301	5/5	0.93	0.07	83,91,94,95	0
6	EDO	B	301	4/4	0.94	0.10	47,51,56,61	0
8	SO4	A	304	5/5	0.94	0.09	71,81,82,84	0
6	EDO	B	303	4/4	0.94	0.11	48,53,57,57	0
6	EDO	B	302	4/4	0.95	0.12	50,52,53,57	0
6	EDO	A	301	4/4	0.95	0.08	45,49,50,50	0
6	EDO	A	302	4/4	0.96	0.10	57,58,58,60	0

6.5 Other polymers [i](#)

There are no such residues in this entry.