



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 12:23 AM UTC

PDB ID : 5MES / pdb_00005mes
Title : MCL1 FAB COMPLEX IN COMPLEX WITH COMPOUND 29
Authors : Hargreaves, D.
Deposited on : 2016-11-16
Resolution : 2.24 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3416 (2.26-2.22)
Clashscore	190562	3556 (2.26-2.22)
Ramachandran outliers	187476	3500 (2.26-2.22)
Sidechain outliers	187428	3501 (2.26-2.22)
RSRZ outliers	180081	3415 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	162	4% 80% 10% 7%
2	H	230	2% 79% 11% 7%
3	L	226	% 78% 14% 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	7LT	A	401	X	-	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4679 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Induced myeloid leukemia cell differentiation protein Mcl-1 homolog, Induced myeloid leukemia cell differentiation protein Mcl-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	151	Total	C	N	O	S	0	1	0
			1219	765	226	225	3			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	166	GLY	-	expression tag	UNP P97287
A	167	PRO	-	expression tag	UNP P97287
A	168	LEU	-	expression tag	UNP P97287
A	169	GLY	-	expression tag	UNP P97287
A	170	SER	-	expression tag	UNP P97287
A	171	GLU	-	expression tag	UNP P97287


- Molecule 2 is a protein called Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	213	Total	C	N	O	S	0	0	0
			1599	1013	268	310	8			

- Molecule 3 is a protein called Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	213	Total	C	N	O	S	1	1	0
			1604	994	274	330	6			

- Molecule 4 is (5 {R},13 {S},17 {S})-5-[[4-chloranyl-3-(2-phenylethyl)phenyl]methyl]-13-[(4-chlorophenyl)methyl]-8-methyl-1,4,8,12,16-pentazatricyclo[15.8.1.0^{2,5}]{20,25}]hexacosa-20(25),21,23-triene-3,7,15,26-tetrone (CCD ID: 7LT) (formula: C₄₄H₄₉Cl₂N₅O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	Cl	N	O	0	0
			55	44	2	5	4		



- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	21	Total	O	0	0
			21	21		
5	H	101	Total	O	0	0
			101	101		
5	L	80	Total	O	0	0
			80	80		



3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Induced myeloid leukemia cell differentiation protein Mcl-1 homolog, Induced myeloid leukemia cell differentiation protein Mcl-1



- Molecule 2: Heavy Chain



- Molecule 3: Light Chain



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	168.47Å 40.33Å 102.84Å 90.00° 126.11° 90.00°	Depositor
Resolution (Å)	81.02 – 2.24 81.02 – 2.24	Depositor EDS
% Data completeness (in resolution range)	99.0 (81.02-2.24) 99.0 (81.02-2.24)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.86 (at 2.25Å)	Xtriage
Refinement program	BUSTER 2.11.6	Depositor
R, R_{free}	0.180 , 0.230 (Not available) , 0.218	Depositor DCC
R_{free} test set	1326 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	43.6	Xtriage
Anisotropy	0.023	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 56.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.014 for -h-2*k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4679	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 7LT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.89	0/1242	1.53	3/1670 (0.2%)
2	H	0.88	0/1639	1.14	5/2237 (0.2%)
3	L	0.85	0/1644	1.12	4/2244 (0.2%)
All	All	0.87	0/4525	1.25	12/6151 (0.2%)

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	60	PRO	CA-C-N	6.07	129.24	120.38
3	L	60	PRO	C-N-CA	6.07	129.24	120.38
1	A	283	GLN	CA-C-N	6.05	128.67	120.38
1	A	283	GLN	C-N-CA	6.05	128.67	120.38
2	H	193	PRO	CA-C-N	5.32	127.79	120.39
2	H	193	PRO	C-N-CA	5.32	127.79	120.39
2	H	107	ASP	CA-CB-CG	5.25	117.85	112.60
2	H	68	PHE	CA-CB-CG	5.19	118.99	113.80
1	A	172	ASP	CA-CB-CG	5.18	117.78	112.60
2	H	169	VAL	N-CA-CB	5.16	117.68	110.56
3	L	190	GLU	CA-C-N	5.05	127.99	120.31
3	L	190	GLU	C-N-CA	5.05	127.99	120.31

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1219	0	1226	7	0
2	H	1599	0	1579	10	0
3	L	1604	0	1537	15	0
4	A	55	0	0	0	0
5	A	21	0	0	0	0
5	H	101	0	0	0	0
5	L	80	0	0	0	0
All	All	4679	0	4342	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:202:GLU:C	3:L:205:GLY:N	2.44	0.75
1:A:202:ALA:HB1	1:A:206:GLY:HA3	1.70	0.74
3:L:187:ARG:HH21	3:L:191:ARG:HH22	1.49	0.59
1:A:298:LEU:HD22	1:A:306:LEU:HD11	1.86	0.56
3:L:79:LEU:HD21	3:L:107:LEU:HD21	1.87	0.56
2:H:170:HIS:CD2	3:L:141:THR:HG21	2.43	0.53
1:A:232:LEU:HD21	1:A:277:HIS:HB2	1.91	0.53
3:L:193:SER:O	3:L:213:SER:HA	2.09	0.52
2:H:125:PRO:HB3	2:H:151:TYR:HB3	1.92	0.50
3:L:173:ASN:HD21	3:L:175:LYS:HD2	1.77	0.49
2:H:176:LEU:HG	2:H:181:TYR:CE1	2.46	0.48
1:A:288:GLU:HB2	1:A:289:PRO:HD3	1.95	0.48
3:L:160:THR:HG23	3:L:161:GLN:HG2	1.95	0.47
3:L:79:LEU:HD13	3:L:109:VAL:HG22	1.97	0.47
2:H:30:SER:O	2:H:53:SER:HB2	2.15	0.46
2:H:47:TRP:CG	3:L:99:TRP:HB2	2.50	0.46
2:H:141:MET:N	2:H:191:SER:HG	2.13	0.46
2:H:20:LEU:HG	2:H:83:MET:HE2	1.97	0.45
3:L:200:THR:OG1	3:L:207:THR:HB	2.16	0.44
2:H:130:LEU:HB2	2:H:145:GLY:C	2.42	0.44
2:H:142:VAL:HG11	2:H:193:PRO:HG3	2.00	0.44
1:A:309:GLN:O	1:A:314:GLY:HA3	2.18	0.43
3:L:173:ASN:ND2	3:L:175:LYS:HD2	2.34	0.43
2:H:12:VAL:HG11	2:H:86:LEU:HD13	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:114:LYS:HG3	3:L:145:PRO:HD3	2.00	0.42
3:L:80:GLN:O	3:L:109:VAL:HG21	2.19	0.41
3:L:170:LYS:HE2	3:L:174:ASN:HA	2.01	0.41
1:A:277:HIS:CE1	1:A:281:ILE:HG21	2.56	0.41
1:A:184:ARG:HD2	1:A:210:LEU:HD11	2.03	0.40
3:L:82:GLU:H	3:L:82:GLU:CD	2.28	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	150/162 (93%)	144 (96%)	5 (3%)	1 (1%)	18	16
2	H	209/230 (91%)	206 (99%)	2 (1%)	1 (0%)	24	23
3	L	210/226 (93%)	205 (98%)	5 (2%)	0	100	100
All	All	569/618 (92%)	555 (98%)	12 (2%)	2 (0%)	30	30

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	201	GLU
2	H	191	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	131/137 (96%)	122 (93%)	9 (7%)	14	11
2	H	181/192 (94%)	168 (93%)	13 (7%)	13	10
3	L	183/192 (95%)	167 (91%)	16 (9%)	9	6
All	All	495/521 (95%)	457 (92%)	38 (8%)	12	9

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	173	ASP
1	A	184	ARG
1	A	210	LEU
1	A	229	GLN
1	A	246	LEU
1	A	281	ILE
1	A	293	SER
1	A	298	LEU
1	A	322	GLU
2	H	1	GLN
2	H	33	SER
2	H	87	ARG
2	H	111	GLN
2	H	121	LYS
2	H	130	LEU
2	H	165	LEU
2	H	169	VAL
2	H	176	LEU
2	H	183	LEU
2	H	211	LYS
2	H	214	LYS
2	H	217	VAL
3	L	9	SER
3	L	40	LEU
3	L	79	LEU
3	L	82	GLU
3	L	114	LYS
3	L	115	SER
3	L	116	SER
3	L	126	SER
3	L	136	LEU
3	L	161	GLN
3	L	163	MET
3	L	166	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	L	167	GLN
3	L	172	SER
3	L	179	SER
3	L	207	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	189	GLN
1	A	221	GLN
1	A	252	HIS
1	A	283	GLN
1	A	309	GLN
2	H	82	GLN
2	H	111	GLN
2	H	170	HIS
3	L	38	GLN
3	L	161	GLN
3	L	192	HIS
3	L	206	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	7LT	A	401	-	60,60,60	0.26	0	79,82,82	0.71	2 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	7LT	A	401	-	1/1/9/12	13/44/61/61	0/4/6/6

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	401	7LT	C4-C5-C6	-3.85	106.96	112.41
4	A	401	7LT	C16-C7-N2	2.26	113.48	108.73

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	401	7LT	N3

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	401	7LT	C2-C1-N-C
4	A	401	7LT	C37-C4-N1-C3
4	A	401	7LT	C5-C4-N1-C3
4	A	401	7LT	C37-C4-C5-C6
4	A	401	7LT	N1-C4-C5-C6
4	A	401	7LT	N-C1-C2-C3
4	A	401	7LT	C27-C29-C30-C31
4	A	401	7LT	C16-C7-N2-C6
4	A	401	7LT	C2-C3-N1-C4
4	A	401	7LT	C19-C20-C21-O3

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	401	7LT	C19-C20-C21-N
4	A	401	7LT	C2-C1-N-C21
4	A	401	7LT	C8-C7-N2-C6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
3	L	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	L	202:GLU	C	205:GLY	N	2.44

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	151/162 (93%)	0.60	7 (4%)	37 35	22, 66, 103, 131	1 (0%)
2	H	213/230 (92%)	-0.15	5 (2%)	61 62	27, 41, 81, 117	0
3	L	213/226 (94%)	-0.14	2 (0%)	81 82	18, 44, 65, 108	1 (0%)
All	All	577/618 (93%)	0.05	14 (2%)	59 61	18, 46, 86, 131	2 (0%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	275	ALA	3.2
2	H	195	PRO	3.0
2	H	193	PRO	2.6
1	A	199	LEU	2.5
1	A	321	VAL	2.4
3	L	205	GLY	2.4
2	H	221	CYS	2.4
1	A	202	ALA	2.3
1	A	227	ALA	2.2
2	H	192	SER	2.1
2	H	132	PRO	2.1
1	A	232	LEU	2.1
3	L	216	ASP	2.1
1	A	253	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	7LT	A	401	55/55	0.88	0.12	58,63,73,79	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 7LT A 401:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.