



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 07:46 AM UTC

PDB ID : 6MEP / pdb_00006mep
Title : Crystal structure of the catalytic domain of the proto-oncogene tyrosine-protein kinase MER in complex with inhibitor UNC3437
Authors : Da, C.; Zhang, D.; Stashko, M.A.; Cheng, A.; Hunter, D.; Norris-Drouin, J.; Graves, L.; Machius, M.; Miley, M.J.; DeRyckere, D.; Earp, H.S.; Graham, D.K.; Frye, S.V.; Wang, X.; Kireev, D.
Deposited on : 2018-09-06
Resolution : 2.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	313	3% 76% • 20%
1	B	313	2% 73% •• 25%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8005 atoms, of which 3996 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein kinase Mer.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	250	Total	C	H	N	O	S	0	0	0
			4023	1290	2016	332	367	18			
1	B	236	Total	C	H	N	O	S	0	0	0
			3793	1219	1896	314	346	18			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	552	MET	-	expression tag	UNP Q12866
A	553	GLY	-	expression tag	UNP Q12866
A	554	SER	-	expression tag	UNP Q12866
A	555	SER	-	expression tag	UNP Q12866
A	556	HIS	-	expression tag	UNP Q12866
A	557	HIS	-	expression tag	UNP Q12866
A	558	HIS	-	expression tag	UNP Q12866
A	559	HIS	-	expression tag	UNP Q12866
A	560	HIS	-	expression tag	UNP Q12866
A	561	HIS	-	expression tag	UNP Q12866
A	562	SER	-	expression tag	UNP Q12866
A	563	SER	-	expression tag	UNP Q12866
A	564	GLY	-	expression tag	UNP Q12866
A	565	LEU	-	expression tag	UNP Q12866
A	566	VAL	-	expression tag	UNP Q12866
A	567	PRO	-	expression tag	UNP Q12866
A	568	ARG	-	expression tag	UNP Q12866
A	569	GLY	-	expression tag	UNP Q12866
B	552	MET	-	expression tag	UNP Q12866
B	553	GLY	-	expression tag	UNP Q12866
B	554	SER	-	expression tag	UNP Q12866
B	555	SER	-	expression tag	UNP Q12866
B	556	HIS	-	expression tag	UNP Q12866
B	557	HIS	-	expression tag	UNP Q12866
B	558	HIS	-	expression tag	UNP Q12866

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	559	HIS	-	expression tag	UNP Q12866
B	560	HIS	-	expression tag	UNP Q12866
B	561	HIS	-	expression tag	UNP Q12866
B	562	SER	-	expression tag	UNP Q12866
B	563	SER	-	expression tag	UNP Q12866
B	564	GLY	-	expression tag	UNP Q12866
B	565	LEU	-	expression tag	UNP Q12866
B	566	VAL	-	expression tag	UNP Q12866
B	567	PRO	-	expression tag	UNP Q12866
B	568	ARG	-	expression tag	UNP Q12866
B	569	GLY	-	expression tag	UNP Q12866

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	3	Total Cl 3 3	0	0
2	B	2	Total Cl 2 2	0	0

- Molecule 3 is cis-4-[(2-[(4-{[4-(1,3-dioxolan-2-yl)pyridin-2-yl]ethynyl}phenyl)amino]-5-{4-[(4-methylpiperazin-1-yl)methyl]phenyl}pyrimidin-4-yl)amino]cyclohexan-1-ol (CCD ID: 9VS) (formula: C₃₈H₄₃N₇O₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	0	0
			90	38	42	7	3		
3	B	1	Total	C	H	N	O	0	0
			90	38	42	7	3		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	O	0	0
			2	2		
5	B	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tyrosine-protein kinase Mer



• Molecule 1: Tyrosine-protein kinase Mer



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.37Å 92.05Å 69.69Å 90.00° 101.67° 90.00°	Depositor
Resolution (Å)	37.07 – 2.89 37.07 – 2.89	Depositor EDS
% Data completeness (in resolution range)	88.5 (37.07-2.89) 84.7 (37.07-2.89)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.42 (at 2.90Å)	Xtriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, R_{free}	0.205 , 0.268 0.208 , 0.271	Depositor DCC
R_{free} test set	1000 reflections (7.91%)	wwPDB-VP
Wilson B-factor (Å ²)	40.3	Xtriage
Anisotropy	0.331	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 41.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	8005	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, MG, 9VS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.10	0/2049	0.25	0/2768
1	B	0.10	0/1937	0.25	0/2616
All	All	0.10	0/3986	0.25	0/5384

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2007	2016	2015	7	0
1	B	1897	1896	1896	6	0
2	A	3	0	0	1	0
2	B	2	0	0	0	0
3	A	48	42	0	0	0
3	B	48	42	0	4	0
4	B	1	0	0	0	0
5	A	2	0	0	0	0
5	B	1	0	0	0	0
All	All	4009	3996	3911	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 2.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:671:LEU:HD22	3:B:904:9VS:O41	2.02	0.59
1:A:770:GLU:OE1	1:A:844:ARG:NH2	2.37	0.52
1:A:732:ARG:NH2	2:A:902:CL:CL	2.76	0.51
1:B:652:LEU:HA	3:B:904:9VS:C42	2.42	0.49
1:B:650:ILE:HG12	3:B:904:9VS:C43	2.43	0.48
1:B:670:ILE:C	1:B:671:LEU:HD23	2.38	0.48
1:B:650:ILE:HG23	3:B:904:9VS:O44	2.14	0.46
1:A:743:GLY:O	1:A:744:LEU:HB2	2.16	0.45
1:A:580:VAL:HG11	1:A:639:ALA:HA	1.99	0.44
1:A:637:GLU:OE2	1:A:744:LEU:N	2.48	0.44
1:B:780:LYS:NZ	1:B:844:ARG:O	2.50	0.43
1:A:721:HIS:CG	1:A:742:PHE:HB3	2.56	0.41
1:A:637:GLU:HG2	1:A:641:MET:SD	2.62	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	242/313 (77%)	235 (97%)	7 (3%)	0	100	100
1	B	226/313 (72%)	219 (97%)	6 (3%)	1 (0%)	30	59
All	All	468/626 (75%)	454 (97%)	13 (3%)	1 (0%)	43	72

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	823	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/280 (80%)	224 (100%)	0	100	100
1	B	211/280 (75%)	208 (99%)	3 (1%)	59	85
All	All	435/560 (78%)	432 (99%)	3 (1%)	76	92

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	652	LEU
1	B	671	LEU
1	B	741	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	605	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	9VS	A	904	-	54,54,54	3.01	20 (37%)	73,74,74	4.39	26 (35%)
3	9VS	B	904	-	54,54,54	3.13	20 (37%)	73,74,74	3.60	21 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	9VS	A	904	-	-	2/24/52/52	0/7/7/7
3	9VS	B	904	-	-	4/24/52/52	0/7/7/7

All (40) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	904	9VS	C09-N08	9.02	1.49	1.35
3	B	904	9VS	C09-N08	8.92	1.49	1.35
3	B	904	9VS	C15-N16	-8.09	1.32	1.47
3	A	904	9VS	C15-N16	-7.95	1.32	1.47
3	A	904	9VS	C20-N19	-7.53	1.28	1.46
3	B	904	9VS	C20-N19	-7.29	1.29	1.46
3	B	904	9VS	C27-N28	6.69	1.50	1.36
3	B	904	9VS	C34-C33	6.33	1.35	1.19
3	A	904	9VS	C27-N28	6.04	1.49	1.36
3	B	904	9VS	C29-N28	5.68	1.53	1.40
3	A	904	9VS	C34-C33	5.45	1.33	1.19
3	A	904	9VS	C29-N28	5.35	1.52	1.40
3	B	904	9VS	C35-C34	5.30	1.54	1.44
3	A	904	9VS	C35-C34	4.54	1.53	1.44
3	B	904	9VS	O44-C40	-4.25	1.31	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	904	9VS	C21-N19	-4.24	1.36	1.46
3	B	904	9VS	C10-C11	4.24	1.57	1.49
3	A	904	9VS	C17-N16	-4.23	1.35	1.46
3	B	904	9VS	C17-N16	-4.13	1.35	1.46
3	A	904	9VS	O44-C40	-4.03	1.31	1.41
3	B	904	9VS	O41-C40	-3.98	1.31	1.41
3	A	904	9VS	C21-N19	-3.96	1.36	1.46
3	A	904	9VS	O41-C40	-3.79	1.32	1.41
3	A	904	9VS	C10-C11	3.70	1.56	1.49
3	B	904	9VS	C24-C23	3.33	1.44	1.38
3	B	904	9VS	C22-N16	-3.24	1.38	1.46
3	A	904	9VS	C18-N19	-3.21	1.38	1.46
3	A	904	9VS	C27-N26	-3.15	1.30	1.34
3	B	904	9VS	C18-N19	-3.10	1.39	1.46
3	A	904	9VS	C24-C23	3.10	1.43	1.38
3	B	904	9VS	O41-C42	-3.08	1.34	1.43
3	B	904	9VS	C15-C14	2.99	1.56	1.51
3	A	904	9VS	C15-C14	2.90	1.56	1.51
3	A	904	9VS	O41-C42	-2.86	1.34	1.43
3	A	904	9VS	C22-N16	-2.82	1.39	1.46
3	B	904	9VS	C43-C42	-2.81	1.40	1.49
3	B	904	9VS	O44-C43	-2.60	1.35	1.43
3	A	904	9VS	C43-C42	-2.55	1.41	1.49
3	B	904	9VS	C27-N26	-2.54	1.31	1.34
3	A	904	9VS	O44-C43	-2.40	1.36	1.43

All (47) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	904	9VS	C32-C33-C34	-22.11	110.29	176.23
3	B	904	9VS	C35-C34-C33	-18.24	129.64	175.84
3	A	904	9VS	C35-C34-C33	-17.92	130.43	175.84
3	B	904	9VS	C32-C33-C34	-14.91	131.77	176.23
3	A	904	9VS	C34-C35-N36	9.80	130.66	117.25
3	A	904	9VS	O44-C40-C39	9.25	126.67	111.95
3	B	904	9VS	O44-C40-C39	8.65	125.72	111.95
3	A	904	9VS	C45-C35-C34	-8.16	108.59	120.08
3	B	904	9VS	O41-C40-C39	6.98	123.07	111.95
3	A	904	9VS	O41-C40-C39	6.40	122.14	111.95
3	B	904	9VS	N26-C27-N48	-6.02	120.60	126.42
3	B	904	9VS	C37-N36-C35	5.78	124.31	116.51
3	A	904	9VS	C25-C10-C09	5.19	119.88	114.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	904	9VS	C38-C39-C40	4.77	129.60	120.39
3	A	904	9VS	C45-C39-C40	-4.52	112.30	119.82
3	A	904	9VS	N26-C27-N48	-4.42	122.15	126.42
3	A	904	9VS	C10-C09-N48	-4.37	117.43	122.67
3	B	904	9VS	C38-C37-N36	-4.28	118.72	123.97
3	B	904	9VS	C25-N26-C27	4.25	121.63	115.81
3	B	904	9VS	C25-C10-C09	4.16	118.85	114.67
3	A	904	9VS	C25-N26-C27	3.88	121.14	115.81
3	A	904	9VS	C10-C25-N26	-3.85	118.43	124.53
3	A	904	9VS	C46-C32-C33	-3.79	114.07	120.78
3	B	904	9VS	C29-N28-C27	-3.71	119.37	129.38
3	B	904	9VS	C10-C25-N26	-3.64	118.76	124.53
3	B	904	9VS	C10-C09-N48	-3.62	118.33	122.67
3	A	904	9VS	C37-N36-C35	3.21	120.84	116.51
3	A	904	9VS	C21-N19-C18	3.19	114.62	109.54
3	B	904	9VS	C45-C35-N36	-3.10	119.41	123.57
3	A	904	9VS	C25-C10-C11	-2.92	115.33	119.56
3	B	904	9VS	O44-C40-O41	2.85	111.20	105.76
3	A	904	9VS	C29-N28-C27	-2.84	121.72	129.38
3	B	904	9VS	C06-C07-C04	2.84	115.71	110.80
3	B	904	9VS	C05-C06-C07	2.83	114.59	111.49
3	A	904	9VS	O44-C40-O41	2.75	111.01	105.76
3	A	904	9VS	C38-C37-N36	-2.73	120.63	123.97
3	A	904	9VS	C14-C15-N16	-2.66	107.70	113.15
3	A	904	9VS	C31-C32-C33	2.60	125.38	120.78
3	B	904	9VS	C45-C35-C34	2.51	123.62	120.08
3	B	904	9VS	C06-C07-N08	-2.45	106.83	110.77
3	A	904	9VS	C22-C21-N19	2.38	114.68	110.86
3	A	904	9VS	C35-C45-C39	2.23	121.02	119.24
3	B	904	9VS	C27-N48-C09	2.22	121.83	116.36
3	B	904	9VS	C14-C15-N16	-2.19	108.67	113.15
3	A	904	9VS	C45-C35-N36	-2.10	120.75	123.57
3	A	904	9VS	C05-C06-C07	2.08	113.78	111.49
3	B	904	9VS	C09-N08-C07	-2.05	120.71	124.18

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	904	9VS	C33-C34-C35-N36
3	B	904	9VS	C14-C15-N16-C17
3	B	904	9VS	C14-C15-N16-C22

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	B	904	9VS	C47-C29-N28-C27
3	B	904	9VS	C30-C29-N28-C27
3	A	904	9VS	C33-C34-C35-C45


There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	904	9VS	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	250/313 (79%)	0.10	10 (4%) 42 34	17, 38, 83, 135	0
1	B	236/313 (75%)	0.13	7 (2%) 52 43	14, 37, 74, 94	0
All	All	486/626 (77%)	0.12	17 (3%) 47 38	14, 37, 76, 135	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	745	SER	4.0
1	A	577	LEU	4.0
1	A	598	PHE	3.7
1	B	598	PHE	3.0
1	B	777	TYR	2.9
1	B	824	ASP	2.7
1	A	576	LYS	2.4
1	B	578	GLU	2.4
1	A	824	ASP	2.4
1	B	580	VAL	2.3
1	A	633	GLU	2.3
1	A	579	ASP	2.3
1	B	689	GLU	2.3
1	A	597	GLU	2.2
1	A	635	LEU	2.1
1	B	694	HIS	2.1
1	A	596	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	9VS	A	904	48/48	0.81	0.18	29,62,93,111	0
3	9VS	B	904	48/48	0.83	0.21	23,53,93,105	0
2	CL	A	902	1/1	0.89	0.11	43,43,43,43	0
4	MG	B	903	1/1	0.92	0.17	31,31,31,31	0
2	CL	A	901	1/1	0.93	0.08	40,40,40,40	0
2	CL	B	902	1/1	0.97	0.07	44,44,44,44	0
2	CL	A	903	1/1	0.97	0.13	34,34,34,34	0
2	CL	B	901	1/1	0.98	0.03	25,25,25,25	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 9VS A 904:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 9VS B 904:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.