



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 10:26 AM UTC

PDB ID : 4MGZ / pdb_00004mgz
Title : Selective activation of Epac1 and Epac2
Authors : Rehmann, H.
Deposited on : 2013-08-29
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	694	7% 79% 11% 9%
2	R	167	28% 74% 14% 10%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6412 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rap guanine nucleotide exchange factor 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	632	Total	C	N	O	S	0	0	0
			5094	3248	880	935	31			


There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	300	GLY	-	expression tag	UNP Q9EQZ6
E	301	SER	-	expression tag	UNP Q9EQZ6
E	302	PRO	-	expression tag	UNP Q9EQZ6
E	303	GLU	-	expression tag	UNP Q9EQZ6
E	304	SER	-	expression tag	UNP Q9EQZ6
E	305	PHE	-	expression tag	UNP Q9EQZ6

- Molecule 2 is a protein called Ras-related protein Rap-1b.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	R	150	Total	C	N	O	S	0	0	0
			1186	748	196	236	6			

- Molecule 3 is (2S,4aR,6R,7R,7aS)-6-[6-amino-8-(benzylsulfanyl)-9H-purin-9-yl]-2-sulfanyltetrahydro-4H-furo[3,2-d][1,3,2]dioxaphosphinin-7-ol 2-oxide (CCD ID: HR6) (formula: C₁₇H₁₈N₅O₅PS₂).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	E	1	Total	C	N	O	P	S	0	0
			30	17	5	5	1	2		
3	E	1	Total	C	N	O	P	S	7	0
			30	17	5	5	1	2		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	R	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	E	60	Total 60	O 60	0	0
5	R	7	Total 7	O 7	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Rap guanine nucleotide exchange factor 4



- Molecule 2: Ras-related protein Rap-1b



4 Data and refinement statistics

Property	Value	Source
Space group	I 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	125.23Å 147.09Å 226.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.85 – 3.00 35.85 – 3.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (35.85-3.00) 98.2 (35.85-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.19 (at 3.00Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.238 , 0.264 0.239 , 0.263	Depositor DCC
R_{free} test set	1999 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	45.3	Xtriage
Anisotropy	0.328	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 38.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	6412	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HR6, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	E	0.43	0/5195	0.82	3/7020 (0.0%)
2	R	0.45	0/1198	0.82	0/1610
All	All	0.43	0/6393	0.82	3/8630 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	899	GLU	CA-C-N	7.23	127.82	120.38
1	E	899	GLU	C-N-CA	7.23	127.82	120.38
1	E	900	PRO	N-CA-C	6.86	119.07	110.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	5094	0	5125	44	0
2	R	1186	0	1180	17	0
3	E	60	0	36	2	0
4	R	5	0	0	0	0
5	E	60	0	0	0	0
5	R	7	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	6412	0	6341	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:905:PHE:CZ	2:R:67:MET:HE1	2.13	0.84
1:E:905:PHE:HZ	2:R:67:MET:HE1	1.45	0.79
1:E:380:LEU:HD11	1:E:428:LEU:HD13	1.70	0.71
1:E:850:MET:HG3	1:E:882:MET:HE1	1.73	0.68
1:E:365:VAL:HG22	3:E:1002:HR6:H11	1.58	0.68
1:E:774:HIS:HD2	1:E:776:LEU:H	1.41	0.68
2:R:44:VAL:HG11	2:R:159:TYR:HE2	1.61	0.66
2:R:91:ASN:O	2:R:94:GLN:HG2	1.98	0.62
1:E:846:PHE:HD2	1:E:882:MET:HE3	1.65	0.62
1:E:369:GLN:HE22	1:E:389:TYR:H	1.50	0.59
1:E:846:PHE:HD2	1:E:882:MET:CE	2.16	0.59
1:E:365:VAL:HG11	1:E:368:ASN:OD1	2.02	0.57
1:E:708:ASN:H	1:E:708:ASN:HD22	1.52	0.57
2:R:11:SER:HB2	2:R:89:THR:HG22	1.86	0.57
1:E:762:MET:HE1	1:E:808:TRP:CZ3	2.40	0.56
1:E:807:PHE:HB3	1:E:974:VAL:CG1	2.36	0.56
1:E:846:PHE:CD2	1:E:882:MET:HE3	2.43	0.53
1:E:716:LEU:HD11	1:E:722:LEU:HD13	1.90	0.52
2:R:87:GLN:H	2:R:87:GLN:CD	2.18	0.52
1:E:774:HIS:CD2	1:E:776:LEU:H	2.26	0.52
1:E:499:ARG:HD2	1:E:504:LEU:HD13	1.92	0.51
1:E:697:ASN:ND2	1:E:701:GLU:HB3	2.27	0.49
1:E:907:PRO:HB2	2:R:67:MET:HE3	1.94	0.49
1:E:333:LEU:HA	1:E:336:ILE:HD12	1.96	0.48
2:R:11:SER:O	2:R:14:VAL:HB	2.14	0.47
1:E:918:GLU:HG2	2:R:17:SER:OG	2.15	0.47
2:R:94:GLN:HG3	2:R:95:ASP:N	2.29	0.47
2:R:14:VAL:HG12	2:R:16:LYS:HG3	1.96	0.46
1:E:850:MET:CG	1:E:882:MET:HE1	2.43	0.46
1:E:766:ASP:OD1	1:E:801:ARG:NH1	2.45	0.46
1:E:451:GLU:HG3	1:E:456:VAL:HG11	1.97	0.46
1:E:313:ILE:HG22	1:E:329:ILE:HD11	1.98	0.46
2:R:144:LEU:HD21	2:R:156:GLU:HG3	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:754:SER:HB3	1:E:757:ASP:HB2	1.98	0.46
1:E:782:THR:OG1	1:E:913:MET:HG2	2.16	0.45
1:E:871:PHE:HA	1:E:874:PHE:HB2	1.99	0.45
1:E:900:PRO:O	1:E:902:LEU:N	2.50	0.45
1:E:519:HIS:HE1	1:E:570:TYR:HE1	1.65	0.45
2:R:41:ARG:HA	2:R:53:LEU:O	2.17	0.45
2:R:144:LEU:HD21	2:R:156:GLU:CG	2.48	0.44
1:E:783:PHE:O	1:E:787:ASN:ND2	2.51	0.44
1:E:907:PRO:CB	2:R:67:MET:HE3	2.48	0.44
1:E:456:VAL:HG13	1:E:489:LYS:HG2	1.99	0.44
1:E:542:GLN:CD	1:E:542:GLN:C	2.86	0.44
1:E:484:SER:HA	1:E:928:VAL:O	2.18	0.43
2:R:76:GLN:O	2:R:110:PRO:HG2	2.19	0.43
1:E:457:LEU:HB2	1:E:489:LYS:HG2	2.01	0.43
1:E:697:ASN:HD22	1:E:701:GLU:HB3	1.82	0.43
1:E:487:PRO:HB3	1:E:523:MET:HB2	2.01	0.42
1:E:357:GLU:HB3	1:E:427:PHE:HB2	2.01	0.42
1:E:365:VAL:CG2	3:E:1002:HR6:H11	2.29	0.41
1:E:350:LEU:HD11	1:E:439:LEU:HD21	2.01	0.41
1:E:935:MET:HE1	2:R:33:ASP:HA	2.02	0.41
1:E:697:ASN:HB3	1:E:699:GLY:H	1.86	0.40
1:E:311:ARG:HE	1:E:315:ARG:HH22	1.68	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	624/694 (90%)	606 (97%)	17 (3%)	1 (0%)	43	76
2	R	144/167 (86%)	138 (96%)	6 (4%)	0	100	100
All	All	768/861 (89%)	744 (97%)	23 (3%)	1 (0%)	48	80

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	900	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	567/615 (92%)	542 (96%)	25 (4%)	25	60
2	R	132/147 (90%)	121 (92%)	11 (8%)	10	37
All	All	699/762 (92%)	663 (95%)	36 (5%)	21	55

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	326	LEU
1	E	334	LEU
1	E	353	VAL
1	E	381	LYS
1	E	394	VAL
1	E	396	THR
1	E	428	LEU
1	E	456	VAL
1	E	479	LYS
1	E	505	ASN
1	E	542	GLN
1	E	584	LEU
1	E	585	GLU
1	E	596	ARG
1	E	691	LEU
1	E	694	VAL
1	E	708	ASN
1	E	769	LEU
1	E	801	ARG
1	E	821	LYS
1	E	837	LYS
1	E	852	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	855	VAL
1	E	942	THR
1	E	973	ASN
2	R	14	VAL
2	R	36	ILE
2	R	38	ASP
2	R	44	VAL
2	R	73	LYS
2	R	87	GLN
2	R	88	SER
2	R	106	THR
2	R	113	LEU
2	R	146	SER
2	R	160	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	309	HIS
1	E	369	GLN
1	E	424	ASN
1	E	426	HIS
1	E	436	ASN
1	E	535	HIS
1	E	537	HIS
1	E	539	GLN
1	E	542	GLN
1	E	644	GLN
1	E	708	ASN
2	R	116	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	HR6	E	1002	-	31,34,34	2.80	9 (29%)	46,51,51	2.94	13 (28%)
4	SO4	R	201	-	4,4,4	0.25	0	6,6,6	0.21	0
3	HR6	E	1001	-	31,34,34	2.25	8 (25%)	46,51,51	2.33	13 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HR6	E	1002	-	-	2/9/36/36	0/5/5/5
3	HR6	E	1001	-	-	2/9/36/36	0/5/5/5

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	1002	HR6	C9-S2	-8.91	1.60	1.82
3	E	1002	HR6	P-O2P	6.63	1.61	1.46
3	E	1001	HR6	P-O2P	6.49	1.60	1.46
3	E	1001	HR6	C6-N6	5.77	1.48	1.34
3	E	1002	HR6	C6-N6	5.66	1.48	1.34
3	E	1002	HR6	C5-C4	5.45	1.48	1.39
3	E	1001	HR6	C5-C4	5.38	1.48	1.39
3	E	1001	HR6	C5-C6	2.88	1.49	1.41
3	E	1002	HR6	C5-C6	2.79	1.48	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	1001	HR6	C4-N9	-2.69	1.34	1.38
3	E	1002	HR6	C1'-N9	2.69	1.50	1.45
3	E	1002	HR6	C5-N7	-2.42	1.35	1.39
3	E	1001	HR6	C5-N7	-2.30	1.35	1.39
3	E	1002	HR6	C4-N9	-2.29	1.35	1.38
3	E	1001	HR6	C1'-N9	2.27	1.49	1.45
3	E	1001	HR6	C9-S2	-2.09	1.77	1.82
3	E	1002	HR6	C11-C10	2.02	1.42	1.38

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	1002	HR6	C9-S2-C8	13.73	120.71	99.83
3	E	1001	HR6	C9-S2-C8	8.18	112.27	99.83
3	E	1002	HR6	C5-C4-N3	-6.57	117.68	126.72
3	E	1001	HR6	C5-C4-N3	-6.20	118.18	126.72
3	E	1002	HR6	N3-C4-N9	5.33	135.01	127.33
3	E	1001	HR6	N3-C4-N9	5.01	134.55	127.33
3	E	1002	HR6	C10-C9-S2	4.31	126.40	111.04
3	E	1002	HR6	C2-N3-C4	4.16	121.99	111.83
3	E	1001	HR6	C6-C5-N7	4.07	135.75	131.72
3	E	1001	HR6	C2-N3-C4	4.07	121.76	111.83
3	E	1001	HR6	O3'-C3'-C4'	-3.70	107.91	110.71
3	E	1002	HR6	C6-C5-N7	3.63	135.32	131.72
3	E	1002	HR6	O3'-C3'-C2'	3.60	119.13	115.61
3	E	1001	HR6	N1-C2-N3	-3.50	123.29	128.58
3	E	1002	HR6	N1-C2-N3	-3.47	123.32	128.58
3	E	1001	HR6	C4-C5-N7	-3.29	107.55	110.91
3	E	1001	HR6	O3'-C3'-C2'	3.04	118.58	115.61
3	E	1002	HR6	C4-C5-N7	-3.03	107.82	110.91
3	E	1002	HR6	N9-C8-N7	2.78	116.93	114.01
3	E	1002	HR6	C5-C4-N9	2.67	107.31	105.63
3	E	1001	HR6	C5-C4-N9	2.60	107.27	105.63
3	E	1002	HR6	S2-C8-N7	-2.58	119.97	125.25
3	E	1001	HR6	N9-C8-N7	2.41	116.54	114.01
3	E	1002	HR6	C2'-C1'-N9	-2.41	111.97	115.44
3	E	1001	HR6	C2'-C1'-N9	-2.31	112.10	115.44
3	E	1001	HR6	C5'-C4'-C3'	-2.08	108.66	112.56

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	E	1002	HR6	C10-C9-S2-C8
3	E	1001	HR6	C2'-C1'-N9-C4
3	E	1001	HR6	O4'-C1'-N9-C4
3	E	1002	HR6	N9-C8-S2-C9


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	1002	HR6	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	E	632/694 (91%)	0.44	50 (7%) 18 10	32, 46, 64, 76	0
2	R	150/167 (89%)	1.48	47 (31%) 1 1	44, 64, 88, 90	0
All	All	782/861 (90%)	0.64	97 (12%) 8 5	32, 48, 75, 90	0

All (97) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	R	105	ASP	6.2
2	R	51	CYS	5.3
2	R	107	ASP	5.3
2	R	125	VAL	5.0
1	E	441	ASP	4.9
2	R	142	ALA	4.9
2	R	129	GLU	4.8
1	E	900	PRO	4.8
2	R	106	THR	4.8
2	R	134	LEU	4.7
1	E	506	GLU	4.1
1	E	372	GLU	3.7
1	E	444	ALA	3.6
2	R	145	GLU	3.5
2	R	130	GLN	3.4
2	R	119	ASP	3.4
1	E	501	GLU	3.3
2	R	57	ASP	3.3
2	R	164	GLN	3.3
1	E	687	SER	3.3
2	R	108	ASP	3.1
2	R	90	PHE	3.1
1	E	309	HIS	3.0
1	E	796	ASP	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	R	113	LEU	2.9
1	E	877	GLU	2.9
2	R	94	GLN	2.8
2	R	50	GLN	2.8
1	E	440	ARG	2.8
1	E	462	VAL	2.7
1	E	963	GLN	2.7
1	E	455	ASP	2.7
1	E	616	SER	2.7
1	E	445	ASN	2.7
2	R	131	GLY	2.7
1	E	901	PRO	2.6
2	R	98	GLU	2.6
1	E	502	PRO	2.6
1	E	341	HIS	2.6
2	R	83	SER	2.6
1	E	766	ASP	2.6
1	E	701	GLU	2.6
1	E	751	GLU	2.6
2	R	162	VAL	2.6
1	E	311	ARG	2.5
1	E	686	GLY	2.5
2	R	120	LEU	2.5
1	E	952	ASN	2.5
1	E	614	GLN	2.5
1	E	603	GLU	2.5
1	E	572	ASP	2.5
2	R	132	GLN	2.5
2	R	122	ASP	2.5
2	R	147	SER	2.4
1	E	324	ASP	2.4
1	E	962	HIS	2.4
1	E	586	GLU	2.4
1	E	496	GLU	2.4
2	R	89	THR	2.4
2	R	133	ASN	2.4
2	R	143	PHE	2.4
2	R	118	CYS	2.3
1	E	312	MET	2.3
1	E	818	GLN	2.3
2	R	115	GLY	2.3
2	R	160	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	R	86	ALA	2.3
1	E	478	GLN	2.3
2	R	156	GLU	2.2
2	R	123	GLU	2.2
2	R	112	ILE	2.2
2	R	84	ILE	2.2
2	R	4	TYR	2.2
1	E	869	SER	2.2
2	R	116	ASN	2.2
1	E	453	ASP	2.2
1	E	452	HIS	2.2
2	R	117	LYS	2.2
1	E	550	ASP	2.2
1	E	344	THR	2.1
1	E	861	ALA	2.1
1	E	443	GLU	2.1
1	E	990	PRO	2.1
2	R	126	VAL	2.1
1	E	411	ASP	2.1
1	E	615	ILE	2.1
2	R	128	LYS	2.1
1	E	499	ARG	2.1
2	R	144	LEU	2.0
1	E	683	ASP	2.0
1	E	425	CYS	2.0
2	R	87	GLN	2.0
2	R	52	MET	2.0
2	R	159	TYR	2.0
1	E	757	ASP	2.0
1	E	838	GLU	2.0
2	R	161	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	HR6	E	1002	30/30	0.91	0.16	20,78,79,79	7
3	HR6	E	1001	30/30	0.95	0.11	33,34,47,47	0
4	SO4	R	201	5/5	0.98	0.06	35,35,36,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers ⓘ

There are no such residues in this entry.