



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 04:56 PM UTC

PDB ID : 7MG0 / pdb_00007mg0
Title : Crystal structure of EV-D68 2A protease
Authors : Liu, C.; Lee, M.-Y.; Liu, W.; Wang, J.
Deposited on : 2021-04-12
Resolution : 3.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1085 (3.54-3.46)
Clashscore	190562	1140 (3.54-3.46)
Ramachandran outliers	187476	1113 (3.54-3.46)
Sidechain outliers	187428	1114 (3.54-3.46)
RSRZ outliers	180081	1084 (3.54-3.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	147	78% 18% .
1	B	147	74% 21% . .
1	C	147	69% 26% 5%
1	D	147	71% 23% 6%
1	E	147	75% 22% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	147	% 76% 19% 5%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11167 atoms, of which 5169 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protease 2A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	E	142	Total	C	H	N	O	S	0	0	0
			1933	657	899	183	188	6			
1	A	141	Total	C	H	N	O	S	0	0	0
			1813	621	828	175	183	6			
1	B	141	Total	C	H	N	O	S	0	0	0
			1892	650	872	177	187	6			
1	C	139	Total	C	H	N	O	S	0	0	0
			1840	618	863	175	178	6			
1	D	138	Total	C	H	N	O	S	0	0	0
			1835	620	856	175	178	6			
1	F	140	Total	C	H	N	O	S	0	0	0
			1836	623	851	173	183	6			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	E	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		
2	C	1	Total	Zn	0	0
			1	1		
2	D	1	Total	Zn	0	0
			1	1		
2	F	1	Total	Zn	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	2	Total 2	O 2	0	0
3	A	1	Total 1	O 1	0	0
3	B	2	Total 2	O 2	0	0
3	C	1	Total 1	O 1	0	0
3	D	2	Total 2	O 2	0	0
3	F	4	Total 4	O 4	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Protease 2A




- Molecule 1: Protease 2A



- Molecule 1: Protease 2A




- Molecule 1: Protease 2A





- Molecule 1: Protease 2A





● Molecule 1: Protease 2A



4 Data and refinement statistics

Property	Value	Source
Space group	P 4	Depositor
Cell constants a, b, c, α , β , γ	118.16Å 118.16Å 81.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.88 – 3.50 47.88 – 3.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.88-3.50) 98.8 (47.88-3.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.28 (at 3.19Å)	Xtriage
Refinement program	PHENIX dev_4189	Depositor
R, R_{free}	0.238 , 0.285 0.235 , 0.283	Depositor DCC
R_{free} test set	1883 reflections (10.04%)	wwPDB-VP
Wilson B-factor (Å ²)	107.0	Xtriage
Anisotropy	0.080	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 185.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	0.468 for h,-k,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	11167	wwPDB-VP
Average B, all atoms (Å ²)	128.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.14	0/1005	0.42	0/1371
1	B	0.12	0/1046	0.39	0/1427
1	C	0.24	0/1000	0.49	0/1365
1	D	0.14	0/1002	0.35	0/1366
1	E	0.14	0/1060	0.35	0/1445
1	F	0.27	0/1007	0.50	0/1375
All	All	0.18	0/6120	0.42	0/8349

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	985	828	912	25	0
1	B	1020	872	944	34	0
1	C	977	863	893	28	0
1	D	979	856	902	33	0
1	E	1034	899	964	25	0
1	F	985	851	909	22	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	1	0	0	0	0
3	B	2	0	0	0	0
3	C	1	0	0	0	0
3	D	2	0	0	3	0
3	E	2	0	0	1	0
3	F	4	0	0	0	0
All	All	5998	5169	5524	160	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 160 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:31:VAL:HG22	1:F:31:VAL:HG22	1.51	0.91
1:A:113:CYS:HB3	1:A:118:ILE:HD11	1.50	0.90
1:B:61:TYR:OH	1:B:66:ASP:OD1	1.98	0.81
1:D:31:VAL:HG12	1:D:38:LEU:HG	1.63	0.79
1:E:82:GLU:OE1	1:E:84:ASN:ND2	2.17	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	139/147 (95%)	127 (91%)	12 (9%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	139/147 (95%)	131 (94%)	7 (5%)	1 (1%)	18	51
1	C	137/147 (93%)	120 (88%)	16 (12%)	1 (1%)	18	51
1	D	136/147 (92%)	127 (93%)	9 (7%)	0	100	100
1	E	140/147 (95%)	132 (94%)	8 (6%)	0	100	100
1	F	138/147 (94%)	130 (94%)	8 (6%)	0	100	100
All	All	829/882 (94%)	767 (92%)	60 (7%)	2 (0%)	43	74

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	140	LEU
1	C	126	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	91/115 (79%)	91 (100%)	0	100	100
1	B	97/115 (84%)	97 (100%)	0	100	100
1	C	89/115 (77%)	89 (100%)	0	100	100
1	D	91/115 (79%)	91 (100%)	0	100	100
1	E	98/115 (85%)	98 (100%)	0	100	100
1	F	92/115 (80%)	92 (100%)	0	100	100
All	All	558/690 (81%)	558 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	18	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	18	HIS
1	C	94	ASN
1	C	83	GLN
1	A	92	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	141/147 (95%)	-0.59	0	100	100	77, 132, 165, 188	0
1	B	141/147 (95%)	-0.68	0	100	100	75, 110, 161, 185	0
1	C	139/147 (94%)	-0.58	2 (1%)	73	48	80, 142, 184, 229	0
1	D	138/147 (93%)	-0.56	1 (0%)	84	63	81, 141, 175, 216	0
1	E	142/147 (96%)	-0.72	0	100	100	69, 108, 156, 189	0
1	F	140/147 (95%)	-0.60	2 (1%)	73	48	82, 134, 168, 194	0
All	All	841/882 (95%)	-0.62	5 (0%)	85	65	69, 129, 170, 229	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	123	ALA	2.8
1	C	130	ALA	2.7
1	C	66	ASP	2.1
1	D	99	ALA	2.1
1	F	89	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	E	201	1/1	1.00	0.01	99,99,99,99	0
2	ZN	A	201	1/1	1.00	0.01	148,148,148,148	0
2	ZN	B	201	1/1	1.00	0.02	58,58,58,58	0
2	ZN	C	201	1/1	1.00	0.03	138,138,138,138	0
2	ZN	D	201	1/1	1.00	0.01	99,99,99,99	0
2	ZN	F	201	1/1	1.00	0.02	122,122,122,122	0

6.5 Other polymers [i](#)

There are no such residues in this entry.