



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:01 PM UTC

PDB ID : 5MHB / pdb_00005mhb
Title : Product-Complex of E.coli 5-Amino Laevulinic Acid Dehydratase
Authors : Norton, E.; Erskine, P.T.; Shoolingin-Jordan, P.M.; Cooper, J.B.
Deposited on : 2016-11-23
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	324	 3% 75% 23%

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 2865 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Delta-aminolevulinic acid dehydratase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	323	Total	C	N	O	S	0	2	0
			2501	1564	440	478	19			

- Molecule 2 is 3-[5-(AMINOMETHYL)-4-(CARBOXYMETHYL)-1H-PYRROL-3-YL]PROPANOIC ACID (CCD ID: PBG) (formula: $C_{10}H_{14}N_2O_4$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			16	10	2	4		

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Zn	0	0
			3	3		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	327	Total	O	0	0
			327	327		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Delta-aminolevulinic acid dehydratase



4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	128.62Å 128.62Å 142.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	95.57 – 2.10 95.57 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.0 (95.57-2.10) 99.0 (95.57-2.10)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.01 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0107	Depositor
R, R_{free}	0.157 , 0.213 0.169 , 0.220	Depositor DCC
R_{free} test set	1761 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	25.1	Xtriage
Anisotropy	0.186	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 65.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2865	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GOL, PBG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.87	54/2543 (2.1%)	1.48	18/3424 (0.5%)

All (54) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	102	MET	N-CA	7.31	1.55	1.46
1	A	77	MET	CA-C	7.19	1.61	1.52
1	A	13	LYS	N-CA	7.19	1.55	1.46
1	A	67	ARG	CA-C	7.11	1.61	1.52
1	A	115	VAL	CA-CB	6.61	1.62	1.54
1	A	192	TYR	C-O	6.47	1.32	1.23
1	A	95	GLU	CA-C	-6.41	1.44	1.52
1	A	271	GLN	CA-C	6.17	1.60	1.52
1	A	229	ALA	CA-C	-6.15	1.44	1.52
1	A	196	PHE	CA-C	6.12	1.60	1.52
1	A	244	LEU	C-O	-6.09	1.15	1.23
1	A	80	GLY	C-O	6.07	1.31	1.23
1	A	64	GLU	C-O	-6.03	1.17	1.24
1	A	174	ALA	C-O	-5.99	1.17	1.24
1	A	256	VAL	N-CA	-5.92	1.40	1.46
1	A	23	GLU	C-O	-5.87	1.16	1.24
1	A	78	THR	N-CA	5.84	1.53	1.46
1	A	319	GLU	N-CA	5.84	1.53	1.46
1	A	79	PHE	CA-C	5.82	1.59	1.52
1	A	29	ASN	CA-C	5.81	1.60	1.52
1	A	201	TYR	CA-C	5.78	1.60	1.52
1	A	109	THR	C-O	-5.78	1.17	1.24
1	A	290	GLU	N-CA	-5.76	1.39	1.46
1	A	111	PRO	C-O	-5.71	1.17	1.24
1	A	252	TYR	C-O	5.68	1.32	1.23
1	A	101	ARG	N-CA	5.65	1.53	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	137	GLY	N-CA	5.65	1.51	1.45
1	A	304	ALA	CA-C	5.64	1.60	1.52
1	A	280	LYS	N-CA	5.59	1.53	1.46
1	A	181	ALA	C-O	-5.58	1.17	1.24
1	A	10	ARG	N-CA	5.47	1.52	1.46
1	A	160	ASP	C-O	5.47	1.30	1.24
1	A	69	ALA	CA-C	-5.46	1.45	1.52
1	A	66	GLU	CA-C	-5.42	1.45	1.52
1	A	202	GLY	N-CA	5.41	1.52	1.44
1	A	214	GLY	N-CA	5.40	1.51	1.44
1	A	276	TYR	C-O	-5.39	1.17	1.24
1	A	120	CYS	CA-C	5.31	1.59	1.53
1	A	179	LEU	N-CA	5.30	1.52	1.46
1	A	137	GLY	C-O	5.29	1.26	1.24
1	A	166	ALA	C-O	5.28	1.30	1.24
1	A	42	ASP	CA-C	-5.26	1.46	1.52
1	A	34	PRO	C-O	-5.26	1.18	1.23
1	A	14	SER	CA-C	-5.24	1.47	1.52
1	A	91	ASP	CA-C	5.20	1.59	1.52
1	A	83	HIS	N-CA	5.18	1.52	1.46
1	A	40	GLU	N-CA	5.18	1.54	1.46
1	A	91	ASP	N-CA	5.17	1.52	1.46
1	A	122	CYS	CA-C	5.16	1.59	1.52
1	A	281	PHE	CA-C	-5.09	1.45	1.52
1	A	291	GLU	CA-C	5.09	1.59	1.52
1	A	276	TYR	CA-C	5.04	1.59	1.52
1	A	221	MET	N-CA	5.03	1.52	1.45
1	A	16	ALA	CA-C	5.01	1.59	1.52

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	284	LEU	N-CA-C	-8.29	102.32	111.36
1	A	199	SER	N-CA-C	-7.35	104.04	113.16
1	A	302	LYS	CD-CE-NZ	6.84	133.80	111.90
1	A	77	MET	N-CA-C	-6.02	99.09	108.90
1	A	25	THR	CB-CA-C	-5.66	98.18	109.33
1	A	302	LYS	CG-CD-CE	-5.65	98.31	111.30
1	A	52	GLY	N-CA-C	-5.50	107.77	114.92
1	A	83	HIS	N-CA-C	-5.47	105.92	112.59
1	A	302	LYS	CB-CG-CD	5.47	123.88	111.30
1	A	13	LYS	N-CA-C	5.46	118.15	111.82

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	62	ALA	N-CA-C	-5.36	105.60	111.82
1	A	56	ILE	CB-CA-C	-5.33	104.88	110.53
1	A	237	GLU	CA-C-N	5.23	127.29	120.28
1	A	237	GLU	C-N-CA	5.23	127.29	120.28
1	A	297	SER	CA-CB-OG	-5.14	100.83	111.10
1	A	60	HIS	N-CA-C	5.09	119.19	112.92
1	A	153	VAL	N-CA-C	5.06	115.83	110.72
1	A	293	VAL	N-CA-C	5.04	115.76	110.62

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2501	0	2496	11	0
2	A	16	0	11	2	0
3	A	3	0	0	0	0
4	A	18	0	24	0	0
5	A	327	0	0	3	1
All	All	2865	0	2531	12	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:25:THR:HG22	5:A:617:HOH:O	2.02	0.59
1:A:190:MET:HE1	1:A:243:CYS:SG	2.46	0.55
1:A:129:HIS:CE1	1:A:212:LEU:HD11	2.49	0.48
1:A:134:CYS:SG	1:A:136:HIS:CE1	3.08	0.46
2:A:401:PBG:HHA1	2:A:401:PBG:O2A	2.17	0.44
1:A:13:LYS:NZ	5:A:504:HOH:O	2.44	0.43
1:A:122:CYS:SG	2:A:401:PBG:HHA2	2.59	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:213:LYS:HD3	1:A:213:LYS:N	2.34	0.43
1:A:247:LYS:HA	1:A:248:PRO:C	2.43	0.42
1:A:227[B]:ARG:HD3	5:A:675:HOH:O	2.18	0.42
1:A:165:SER:HB3	1:A:192:TYR:CE1	2.54	0.42
1:A:144:LEU:HD22	1:A:171:GLN:HA	2.01	0.42

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:689:HOH:O	5:A:739:HOH:O[5_555]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	323/324 (100%)	315 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	260/259 (100%)	253 (97%)	7 (3%)	39	45

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	ILE
1	A	134	CYS
1	A	135	GLU
1	A	205	ARG
1	A	213	LYS
1	A	230	ILE
1	A	302	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	108	GLN
1	A	136	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	A	406	-	5,5,5	0.29	0	5,5,5	0.85	0
4	GOL	A	407	-	5,5,5	0.64	0	5,5,5	0.75	0
4	GOL	A	405	-	5,5,5	1.01	0	5,5,5	1.34	1 (20%)
2	PBG	A	401	3	16,16,16	3.43	8 (50%)	18,21,21	2.17	8 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	406	-	-	0/4/4/4	-
4	GOL	A	407	-	-	2/4/4/4	-
4	GOL	A	405	-	-	2/4/4/4	-
2	PBG	A	401	3	-	1/9/11/11	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	PBG	C5A-C2A	-8.77	1.38	1.51
2	A	401	PBG	CHA-C1A	-7.33	1.39	1.50
2	A	401	PBG	C5A-C6A	-3.70	1.45	1.51
2	A	401	PBG	C7A-C3A	3.61	1.60	1.51
2	A	401	PBG	CHA-N1	-2.80	1.31	1.48
2	A	401	PBG	C8A-C9A	2.38	1.56	1.50
2	A	401	PBG	O2A-C6A	-2.28	1.23	1.30
2	A	401	PBG	C4A-C3A	2.14	1.41	1.38

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	PBG	C1A-CHA-N1	4.46	121.49	112.50
2	A	401	PBG	C2A-C1A-NA	4.15	114.63	109.84
2	A	401	PBG	C7A-C8A-C9A	-2.89	106.01	113.67
2	A	401	PBG	C1A-C2A-C3A	-2.75	103.55	107.52
2	A	401	PBG	C6A-C5A-C2A	2.53	120.14	113.76
2	A	401	PBG	O3A-C9A-C8A	-2.37	115.59	123.09
2	A	401	PBG	O1A-C6A-C5A	2.15	128.25	122.11
4	A	405	GOL	O2-C2-C1	-2.07	100.62	109.18
2	A	401	PBG	O4A-C9A-C8A	2.06	120.51	114.00

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	405	GOL	C1-C2-C3-O3
4	A	405	GOL	O2-C2-C3-O3
4	A	407	GOL	C1-C2-C3-O3
4	A	407	GOL	O2-C2-C3-O3
2	A	401	PBG	C2A-C5A-C6A-O1A

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	PBG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	323/324 (99%)	-0.19	11 (3%)	48 50	11, 31, 56, 93	2 (0%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	213	LYS	3.4
1	A	25	THR	2.8
1	A	133	LEU	2.7
1	A	5	ILE	2.7
1	A	212	LEU	2.7
1	A	134	CYS	2.7
1	A	45[A]	LYS	2.6
1	A	2	THR	2.5
1	A	136	HIS	2.4
1	A	3	ASP	2.2
1	A	214	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	A	407	6/6	0.83	0.25	55,80,82,83	0
4	GOL	A	405	6/6	0.85	0.20	55,60,60,63	0
4	GOL	A	406	6/6	0.87	0.17	57,62,66,70	0
2	PBG	A	401	16/16	0.96	0.07	22,33,39,47	0
3	ZN	A	403	1/1	0.99	0.03	31,31,31,31	0
3	ZN	A	404	1/1	1.00	0.05	33,33,33,33	0
3	ZN	A	402	1/1	1.00	0.03	42,42,42,42	1

6.5 Other polymers [i](#)

There are no such residues in this entry.