



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 02:45 AM UTC

PDB ID : 5MHK / pdb_00005mhh
Title : ICP4 DNA-binding domain in complex with 19mer DNA duplex from its own promoter
Authors : Tunnicliffe, R.B.; Lockhart-Cairns, M.P.; Levy, C.; Mould, P.; Jowitt, T.A.; Sito, H.; Baldock, C.; Sandri-Goldin, R.M.; Golovanov, A.P.
Deposited on : 2016-11-24
Resolution : 2.28 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	9078 (2.30-2.26)
Clashscore	190562	9802 (2.30-2.26)
Ramachandran outliers	187476	9690 (2.30-2.26)
Sidechain outliers	187428	9691 (2.30-2.26)
RSRZ outliers	180081	9085 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	19	
1	G	19	
2	F	19	
2	H	19	
3	A	231	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	B	231	
3	C	231	
3	D	231	
4	J	4	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 7513 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*GP*CP*TP*CP*CP*GP*TP*GP*TP*GP*GP*AP*CP*GP*AP*TP*CP*GP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	G	19	Total	C	N	O	P	0	0	0
			390	185	73	114	18			
1	E	15	Total	C	N	O	P	0	0	0
			313	147	60	91	15			

- Molecule 2 is a DNA chain called DNA (5'-D(*GP*CP*TP*CP*CP*GP*TP*GP*TP*GP*GP*AP*CP*GP*AP*TP*CP*GP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	18	Total	C	N	O	P	0	0	0
			364	173	70	104	17			
2	F	16	Total	C	N	O	P	0	0	0
			321	153	60	93	15			

- Molecule 3 is a protein called RS1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	186	Total	C	N	O	S	0	0	0
			1398	886	261	247	4			
3	C	195	Total	C	N	O	S	0	0	0
			1461	922	270	265	4			
3	D	197	Total	C	N	O	S	0	0	0
			1483	935	278	266	4			
3	B	197	Total	C	N	O	S	0	1	0
			1492	943	279	265	5			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	257	GLY	-	expression tag	UNP Q09I77
C	257	GLY	-	expression tag	UNP Q09I77

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	257	GLY	-	expression tag	UNP Q09I77
B	257	GLY	-	expression tag	UNP Q09I77


- Molecule 4 is a protein called ICP4 DNA BINDING DOMAIN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	J	2	Total	C	N	O	0	0	0
			18	13	2	3			

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	H	1	Total	Mg	0	0
			1	1		

- Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	C	1	Total	O	S	0	0
			5	4	1		
6	C	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	D	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	1	Total	Cl	0	0
			1	1		



- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	G	10	Total	O	0	0
			10	10		
8	H	12	Total	O	0	0
			12	12		
8	E	1	Total	O	0	0
			1	1		
8	F	9	Total	O	0	0
			9	9		
8	A	72	Total	O	0	0
			72	72		
8	C	54	Total	O	0	0
			54	54		
8	D	32	Total	O	0	0
			32	32		
8	B	45	Total	O	0	0
			45	45		
8	J	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: DNA (5'-D(*GP*CP*TP*CP*CP*GP*TP*GP*TP*GP*GP*AP*CP*GP*AP*TP*CP*GP*G)-3')




- Molecule 1: DNA (5'-D(*GP*CP*TP*CP*CP*GP*TP*GP*TP*GP*GP*AP*CP*GP*AP*TP*CP*GP*G)-3')




- Molecule 2: DNA (5'-D(*GP*CP*TP*CP*CP*GP*TP*GP*TP*GP*GP*AP*CP*GP*AP*TP*CP*GP*G)-3')




- Molecule 2: DNA (5'-D(*GP*CP*TP*CP*CP*GP*TP*GP*TP*GP*GP*AP*CP*GP*AP*TP*CP*GP*G)-3')





- Molecule 3: RS1








● Molecule 3: RS1



● Molecule 3: RS1



● Molecule 3: RS1



● Molecule 4: ICP4 DNA BINDING DOMAIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	61.51Å 100.73Å 201.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	71.30 – 2.28 71.30 – 2.28	Depositor EDS
% Data completeness (in resolution range)	99.9 (71.30-2.28) 99.9 (71.30-2.28)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.30 (at 2.27Å)	Xtriage
Refinement program	PHENIX (1.11.1 _2575: ???)	Depositor
R, R_{free}	0.211 , 0.243 0.214 , 0.246	Depositor DCC
R_{free} test set	2849 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	44.7	Xtriage
Anisotropy	0.358	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 36.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7513	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	E	0.20	0/351	0.37	0/541
1	G	0.20	0/437	0.41	0/674
2	F	0.20	0/359	0.38	0/551
2	H	0.18	0/408	0.37	0/627
3	A	0.11	0/1439	0.36	0/1960
3	B	0.14	0/1536	0.36	0/2090
3	C	0.11	0/1502	0.32	0/2047
3	D	0.11	0/1524	0.35	0/2075
4	J	0.04	0/18	0.06	0/23
All	All	0.14	0/7574	0.36	0/10588

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	313	0	169	1	0
1	G	390	0	215	2	0
2	F	321	0	180	5	0
2	H	364	0	202	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	1398	0	1359	7	0
3	B	1492	0	1454	9	0
3	C	1461	0	1413	10	0
3	D	1483	0	1441	8	0
4	J	18	0	15	0	0
5	H	1	0	0	0	0
6	A	10	0	0	0	0
6	B	10	0	0	0	0
6	C	10	0	0	0	0
6	D	5	0	0	0	0
7	C	1	0	0	0	0
8	A	72	0	0	1	0
8	B	45	0	0	0	0
8	C	54	0	0	0	0
8	D	32	0	0	0	0
8	E	1	0	0	0	0
8	F	9	0	0	0	0
8	G	10	0	0	0	0
8	H	12	0	0	0	0
8	J	1	0	0	0	0
All	All	7513	0	6448	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:346:GLU:OE2	3:D:349:ARG:NH2	2.27	0.67
2:F:10:DC:H2''	2:F:11:DA:H5''	1.76	0.66
2:F:2:DC:H2'	2:F:3:DG:C8	2.35	0.61
3:C:459:ASP:HB3	3:D:421:ILE:HD13	1.86	0.57
3:A:405:GLN:HA	3:A:408:PHE:CE2	2.39	0.56
3:A:349:ARG:NH2	3:A:365:GLU:O	2.38	0.55
3:A:459:ASP:HB3	3:B:421:ILE:HD13	1.89	0.55
3:D:397:VAL:HG11	3:B:383:PRO:HG3	1.87	0.55
3:B:306:TYR:O	3:B:307:ARG:NH1	2.36	0.54
3:A:457:ARG:NH1	3:B:420:PHE:H	2.07	0.53
3:C:382:THR:OG1	3:C:385:ALA:HB2	2.09	0.53
1:G:29:DT:H2'	1:G:30:DG:C8	2.47	0.49
3:B:349:ARG:NH2	3:B:365:GLU:O	2.46	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:1:DC:H2'	2:F:2:DC:C6	2.49	0.47
3:C:401:VAL:O	3:C:405:GLN:HG2	2.15	0.47
3:A:425:VAL:HA	3:B:457:ARG:HG3	1.96	0.47
2:F:12:DC:H2''	2:F:13:DA:C8	2.50	0.46
3:C:405:GLN:HA	3:C:408:PHE:CE2	2.51	0.46
3:A:303:TYR:OH	8:A:601:HOH:O	2.20	0.45
3:C:482:ASN:O	3:C:486:THR:OG1	2.29	0.44
1:G:26:DG:N2	2:H:14:DC:O2	2.49	0.43
1:E:38:DG:H2''	1:E:39:DG:C8	2.54	0.43
3:D:440:ARG:HD3	3:D:443:TRP:CZ2	2.54	0.42
3:D:386:GLU:CD	3:D:460:ARG:HH22	2.27	0.42
3:D:323:PRO:HB2	3:D:326:ARG:HD3	2.02	0.42
3:B:307:ARG:HD3	3:B:307:ARG:HA	1.91	0.41
3:C:378:ARG:NH1	3:C:382:THR:HG22	2.35	0.41
3:C:425:VAL:HA	3:D:457:ARG:HG3	2.02	0.41
3:B:368:ASP:OD1	3:B:368:ASP:N	2.53	0.41
3:A:330:GLY:HA2	3:A:421:ILE:O	2.20	0.41
3:C:330:GLY:HA2	3:C:421:ILE:O	2.21	0.41
3:C:341:GLY:HA2	3:C:346:GLU:OE2	2.20	0.41
3:B:475:ALA:HB3	3:B:476:PRO:HD3	2.02	0.41
3:C:293:LEU:HD12	3:C:293:LEU:HA	1.84	0.41
3:D:366:LEU:HD11	3:D:403:LEU:HD21	2.02	0.41
2:F:2:DC:H2''	2:F:3:DG:O5'	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	184/231 (80%)	184 (100%)	0	0	100	100
3	B	194/231 (84%)	188 (97%)	5 (3%)	1 (0%)	24	29

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	C	193/231 (84%)	191 (99%)	1 (0%)	1 (0%)	24	29
3	D	195/231 (84%)	193 (99%)	2 (1%)	0	100	100
All	All	766/924 (83%)	756 (99%)	8 (1%)	2 (0%)	36	44

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	412	GLY
3	B	288	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	127/155 (82%)	126 (99%)	1 (1%)	73	84
3	B	137/155 (88%)	136 (99%)	1 (1%)	76	86
3	C	134/155 (86%)	132 (98%)	2 (2%)	57	72
3	D	136/155 (88%)	136 (100%)	0	100	100
4	J	2/2 (100%)	2 (100%)	0	100	100
All	All	536/622 (86%)	532 (99%)	4 (1%)	76	86

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	396	VAL
3	C	382	THR
3	C	396	VAL
3	B	339	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	392	GLN
3	C	392	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 2 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	SO4	A	502	-	4,4,4	0.24	0	6,6,6	0.08	0
6	SO4	C	502	-	4,4,4	0.24	0	6,6,6	0.08	0
6	SO4	B	501	-	4,4,4	0.24	0	6,6,6	0.07	0
6	SO4	B	502	-	4,4,4	0.24	0	6,6,6	0.08	0
6	SO4	C	503	-	4,4,4	0.23	0	6,6,6	0.08	0
6	SO4	A	501	-	4,4,4	0.23	0	6,6,6	0.08	0
6	SO4	D	501	-	4,4,4	0.23	0	6,6,6	0.07	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å²)	Q<0.9
1	E	15/19 (78%)	0.74	0	100 100	48, 59, 110, 117	0
1	G	19/19 (100%)	0.81	2 (10%)	11 12	40, 61, 133, 135	0
2	F	16/19 (84%)	1.06	3 (18%)	3 3	31, 65, 117, 127	0
2	H	18/19 (94%)	0.62	1 (5%)	30 31	35, 62, 128, 128	0
3	A	186/231 (80%)	0.38	11 (5%)	28 29	25, 37, 57, 86	0
3	B	197/231 (85%)	1.01	32 (16%)	4 5	18, 44, 74, 113	1 (0%)
3	C	195/231 (84%)	0.81	20 (10%)	12 13	28, 43, 75, 97	0
3	D	197/231 (85%)	1.11	29 (14%)	6 6	28, 54, 79, 108	0
4	J	2/4 (50%)	2.82	2 (100%)	0 0	95, 95, 95, 100	0
All	All	845/1004 (84%)	0.84	100 (11%)	9 9	18, 45, 86, 135	1 (0%)

All (100) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	300	GLY	6.6
3	B	288	PRO	5.9
3	D	413	ALA	5.4
3	B	324	PRO	5.4
3	B	416	ASN	4.8
3	B	291	VAL	4.4
3	B	414	ALA	4.3
3	D	418	SER	4.2
3	D	417	SER	4.1
3	D	415	ARG	3.9
3	D	485	LEU	3.9
3	C	385	ALA	3.8
3	D	324	PRO	3.7
3	D	291	VAL	3.6
3	D	301	ALA	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	B	292	GLU	3.6
3	D	414	ALA	3.6
3	B	340	TRP	3.5
3	B	325	GLY	3.5
3	D	352	PHE	3.5
3	C	294	ASP	3.5
3	D	416	ASN	3.5
3	D	412	GLY	3.4
1	G	21	DG	3.4
3	C	448	ALA	3.4
3	B	339	LEU	3.3
3	B	341	GLY	3.3
3	B	323	PRO	3.3
3	B	321	PRO	3.2
3	A	420	PHE	3.2
3	B	413	ALA	3.2
3	D	408	PHE	3.2
3	C	292	GLU	3.1
3	A	302	PHE	3.1
3	B	408	PHE	3.1
3	A	301	ALA	3.1
3	A	408	PHE	3.1
3	C	293	LEU	3.0
2	F	16	DG	3.0
4	J	284	THR	3.0
3	C	460	ARG	2.9
3	C	350	ARG	2.9
3	C	486	THR	2.9
3	D	289	ARG	2.9
3	B	287	GLN	2.9
3	B	290	ARG	2.8
3	D	339	LEU	2.8
3	D	304	ALA	2.8
3	B	312	SER	2.7
3	C	485	LEU	2.7
3	B	415	ARG	2.7
3	C	405	GLN	2.7
3	A	485	LEU	2.7
3	B	420	PHE	2.7
4	J	283	PHE	2.7
3	D	311	VAL	2.7
3	C	346	GLU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	A	417	SER	2.6
3	D	329	TYR	2.6
3	B	367	GLY	2.6
3	D	420	PHE	2.6
3	C	298	THR	2.5
3	C	386	GLU	2.5
3	B	338	GLY	2.5
3	D	290	ARG	2.5
3	D	484	ALA	2.5
3	C	341	GLY	2.4
3	D	309	GLY	2.4
3	D	328	LEU	2.4
3	C	396	VAL	2.4
2	H	18	DG	2.4
3	B	289	ARG	2.4
3	B	296	ASP	2.3
3	C	459	ASP	2.3
3	C	354	ALA	2.3
3	A	314	GLU	2.3
2	F	13	DA	2.2
1	G	29	DT	2.2
3	B	412	GLY	2.2
3	D	361	VAL	2.2
3	B	349	ARG	2.2
3	B	484	ALA	2.2
3	C	297	ALA	2.2
2	F	1	DC	2.1
3	A	396	VAL	2.1
3	D	400	ASP	2.1
3	B	314	GLU	2.1
3	D	341	GLY	2.1
3	C	395	ARG	2.1
3	B	329	TYR	2.1
3	D	334	ASP	2.1
3	D	343	PRO	2.1
3	A	415	ARG	2.1
3	C	480	ARG	2.1
3	B	485	LEU	2.1
3	A	303	TYR	2.1
3	B	409	ARG	2.1
3	B	482	ASN	2.0
3	B	480	ARG	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	D	438	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	SO4	B	502	5/5	0.68	0.12	64,75,93,102	0
6	SO4	C	503	5/5	0.76	0.17	55,60,77,88	0
6	SO4	A	502	5/5	0.85	0.20	44,45,70,89	0
6	SO4	D	501	5/5	0.88	0.12	60,65,68,76	0
6	SO4	B	501	5/5	0.89	0.12	55,67,69,70	0
6	SO4	C	502	5/5	0.89	0.09	72,74,84,84	0
6	SO4	A	501	5/5	0.90	0.12	52,65,68,72	0
5	MG	H	101	1/1	0.94	0.15	50,50,50,50	0
7	CL	C	501	1/1	0.96	0.12	56,56,56,56	0

6.5 Other polymers [i](#)

There are no such residues in this entry.