



wwPDB NMR Structure Validation Summary Report ⓘ

Mar 5, 2026 – 06:21 PM UTC

PDB ID : 2MJZ / pdb_00002mjz
BMRB ID : 19747
Title : Capsid model of M13 bacteriophage virus from Magic-angle spinning NMR and Rosetta modeling
Authors : Morag, O.; Sgourakis, N.G.; Baker, D.; Goldbourn, A.
Deposited on : 2014-01-22

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLID-STATE NMR

The overall completeness of chemical shifts assignment is 1%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	50	86% 14%
1	B	50	86% 14%
1	C	50	86% 14%
1	D	50	86% 14%
1	E	50	86% 14%
1	F	50	86% 14%
1	G	50	86% 14%
1	H	50	86% 14%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain	
1	I	50		86%14%
1	J	50		86%14%
1	K	50		86%14%
1	L	50		86%14%
1	M	50		86%14%
1	N	50		86%14%
1	O	50		86%14%
1	P	50		86%14%
1	Q	50		86%14%
1	R	50		86%14%
1	S	50		86%14%
1	T	50		86%14%
1	U	50		86%14%
1	V	50		86%14%
1	W	50		86%14%
1	X	50		86%14%
1	Y	50		86%14%
1	Z	50		86%14%
1	a	50		86%14%
1	b	50		86%14%
1	c	50		86%14%
1	d	50		86%14%
1	e	50		86%14%
1	f	50		86%14%
1	g	50		86%14%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain	
1	h	50		
1	i	50		

2 Ensemble composition and analysis

This entry contains 3 models. Model 3 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:5-A:47, B:5-B:47, C:5-C:47, D:5-D:47, E:5-E:47, F:5-F:47, G:5-G:47, H:5-H:47, I:5-I:47, J:5-J:47, K:5-K:47, L:5-L:47, M:5-M:47, N:5-N:47, O:5-O:47, P:5-P:47, Q:5-Q:47, R:5-R:47, S:5-S:47, T:5-T:47, U:5-U:47, V:5-V:47, W:5-W:47, X:5-X:47, Y:5-Y:47, Z:5-Z:47, a:5-a:47, b:5-b:47, c:5-c:47, d:5-d:47, e:5-e:47, f:5-f:47, g:5-g:47, h:5-h:47, i:5-i:47 (1505)	0.63	3

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

NmrClust was unable to cluster the ensemble.

Error message: No clusters in NmrClust output

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 26145 atoms, of which 13195 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Capsid protein G8P.

Mol	Chain	Residues	Atoms						Trace
1	A	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	B	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	C	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	D	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	E	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	F	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	G	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	H	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	I	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	J	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	K	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	L	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	M	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	N	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	O	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	P	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	Q	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	R	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	S	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms						Trace
1	T	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	U	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	V	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	W	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	X	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	Y	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	Z	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	a	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	b	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	c	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	d	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	e	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	f	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	g	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	h	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	
1	i	50	Total	C	H	N	O	S	0
			747	241	377	58	70	1	

4 Residue-property plots [i](#)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: Capsid protein G8P




- Molecule 1: Capsid protein G8P




- Molecule 1: Capsid protein G8P



- Molecule 1: Capsid protein G8P



- Molecule 1: Capsid protein G8P



- Molecule 1: Capsid protein G8P

Chain F:  86% 14%



- Molecule 1: Capsid protein G8P

Chain G:  86% 14%



- Molecule 1: Capsid protein G8P

Chain H:  86% 14%



- Molecule 1: Capsid protein G8P

Chain I:  86% 14%



- Molecule 1: Capsid protein G8P

Chain J:  86% 14%



- Molecule 1: Capsid protein G8P

Chain K:  86% 14%



- Molecule 1: Capsid protein G8P

Chain L:  86% 14%



- Molecule 1: Capsid protein G8P

Chain M:  86% 14%



- Molecule 1: Capsid protein G8P

Chain N: 86% 14%



- Molecule 1: Capsid protein G8P

Chain O: 86% 14%



- Molecule 1: Capsid protein G8P

Chain P: 86% 14%



- Molecule 1: Capsid protein G8P

Chain Q: 86% 14%



- Molecule 1: Capsid protein G8P

Chain R: 86% 14%



- Molecule 1: Capsid protein G8P

Chain S: 86% 14%



- Molecule 1: Capsid protein G8P

Chain T: 86% 14%



- Molecule 1: Capsid protein G8P

Chain U:  86% 14%



- Molecule 1: Capsid protein G8P

Chain V:  86% 14%



- Molecule 1: Capsid protein G8P

Chain W:  86% 14%



- Molecule 1: Capsid protein G8P

Chain X:  86% 14%



- Molecule 1: Capsid protein G8P

Chain Y:  86% 14%



- Molecule 1: Capsid protein G8P

Chain Z:  86% 14%



- Molecule 1: Capsid protein G8P

Chain a:  86% 14%



- Molecule 1: Capsid protein G8P

Chain b:  86% 14%



- Molecule 1: Capsid protein G8P

Chain c:  86% 14%



- Molecule 1: Capsid protein G8P

Chain d:  86% 14%



- Molecule 1: Capsid protein G8P

Chain e:  86% 14%



- Molecule 1: Capsid protein G8P

Chain f:  86% 14%



- Molecule 1: Capsid protein G8P

Chain g:  86% 14%



- Molecule 1: Capsid protein G8P

Chain h:  86% 14%



- Molecule 1: Capsid protein G8P

Chain i:  86% 14%



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 3. Colouring as in section 4.1 above.

- Molecule 1: Capsid protein G8P

Chain A: 



- Molecule 1: Capsid protein G8P

Chain B: 



- Molecule 1: Capsid protein G8P

Chain C: 



- Molecule 1: Capsid protein G8P

Chain D: 



- Molecule 1: Capsid protein G8P

Chain E: 



- Molecule 1: Capsid protein G8P

Chain F: 



- Molecule 1: Capsid protein G8P

Chain G: 



- Molecule 1: Capsid protein G8P

Chain H:  86% 14%



- Molecule 1: Capsid protein G8P

Chain I:  86% 14%



- Molecule 1: Capsid protein G8P

Chain J:  86% 14%



- Molecule 1: Capsid protein G8P

Chain K:  86% 14%



- Molecule 1: Capsid protein G8P

Chain L:  86% 14%



- Molecule 1: Capsid protein G8P

Chain M:  86% 14%



- Molecule 1: Capsid protein G8P

Chain N:  86% 14%



- Molecule 1: Capsid protein G8P

Chain O:  86% 14%



- Molecule 1: Capsid protein G8P

Chain P:  86% 14%



- Molecule 1: Capsid protein G8P

Chain Q:  86% 14%



- Molecule 1: Capsid protein G8P

Chain R:  86% 14%



- Molecule 1: Capsid protein G8P

Chain S:  86% 14%



- Molecule 1: Capsid protein G8P

Chain T:  86% 14%



- Molecule 1: Capsid protein G8P

Chain U:  86% 14%



- Molecule 1: Capsid protein G8P

Chain V:  86% 14%



- Molecule 1: Capsid protein G8P

Chain W:  86% 14%



- Molecule 1: Capsid protein G8P

Chain X:  86% 14%



- Molecule 1: Capsid protein G8P

Chain Y:  86% 14%



- Molecule 1: Capsid protein G8P

Chain Z:  86% 14%



- Molecule 1: Capsid protein G8P

Chain a:  86% 14%



- Molecule 1: Capsid protein G8P

Chain b:  86% 14%



- Molecule 1: Capsid protein G8P

Chain c:  86% 14%



- Molecule 1: Capsid protein G8P

Chain d: 86% 14%



- Molecule 1: Capsid protein G8P

Chain e: 86% 14%




- Molecule 1: Capsid protein G8P

Chain f: 86% 14%



- Molecule 1: Capsid protein G8P

Chain g: 86% 14%



- Molecule 1: Capsid protein G8P

Chain h: 86% 14%



- Molecule 1: Capsid protein G8P

Chain i: 86% 14%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *Fold-and-dock*.

Of the 5000 calculated structures, 3 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Rosetta	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	292
Number of shifts mapped to atoms	292
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	1%

Note: This is a solid-state NMR structure, where hydrogen atoms are typically not assigned a chemical shift value, which may lead to lower completeness of assignment measure.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	33915	35070	35070	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	B	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	C	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	D	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	E	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	G	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	H	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	I	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	J	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	K	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	L	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	M	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	N	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	O	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	P	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	Q	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	R	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	S	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	T	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	U	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	V	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	W	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	X	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	Y	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	Z	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	a	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	b	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	c	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	d	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	e	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	f	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	g	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	h	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	i	43/50 (86%)	43±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
All	All	4515/5250 (86%)	4515 (100%)	0 (0%)	0 (0%)	100	100

There are no Ramachandran outliers.

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	B	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	C	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	D	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	E	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	F	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	G	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	H	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	I	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	J	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	K	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	L	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	M	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	N	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	O	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	P	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	Q	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	R	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	S	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	T	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	U	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	V	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	W	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	X	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	Y	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	Z	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	a	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	b	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	c	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	d	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	e	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	f	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	g	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	h	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
1	i	32/36 (89%)	32±0 (100±0%)	0±0 (0±0%)	100	100
All	All	3360/3780 (89%)	3360 (100%)	0 (0%)	100	100

There are no protein residues with a non-rotameric sidechain to report.

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 1% for the well-defined parts and 1% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	292
Number of shifts mapped to atoms	292
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	50	-0.66 ± 0.11	Should be checked
$^{13}\text{C}_\beta$	46	0.19 ± 0.14	None needed (< 0.5 ppm)
$^{13}\text{C}'$	49	-0.83 ± 0.22	Should be applied
^{15}N	48	-0.22 ± 0.29	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 1%, i.e. 251 atoms were assigned a chemical shift out of a possible 19950. 0 out of 210 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	127/7560 (2%)	0/3080 (0%)	85/3010 (3%)	42/1470 (3%)
Sidechain	95/10290 (1%)	0/6895 (0%)	91/3185 (3%)	4/210 (2%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	29/2100 (1%)	0/1015 (0%)	29/1050 (3%)	0/35 (0%)
Overall	251/19950 (1%)	0/10990 (0%)	205/7245 (3%)	46/1715 (3%)

Note: This is a solid-state NMR structure, where hydrogen atoms are typically not assigned a chemical shift value, which may lead to lower completeness of assignment measure.

7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	19	THR	N	11.50	91.89 – 138.78	-22.1

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

