



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 18, 2026 – 05:22 PM UTC

PDB ID : 4MJ2 / pdb_00004mj2
Title : Crystal structure of apo-iduronidase in the R3 form
Authors : Bie, H.; Yin, J.; He, X.; Kermode, A.R.; Goddard-Borger, E.D.; Withers, S.G.; James, M.N.G.
Deposited on : 2013-09-03
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	653	6% 90% • 9%
1	B	653	4% 90% • 7%
2	C	2	50% 50%
3	D	5	80% 20%
4	E	3	67% 33%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	F	7	29% 57% 14%

2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 10336 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-L-iduronidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	595	Total	C	N	O	S	0	1	0
			4745	3036	856	841	12			
1	B	607	Total	C	N	O	S	0	4	0
			4851	3099	878	860	14			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	33	GLN	HIS	SEE REMARK 999	UNP P35475
A	63	PRO	GLN	SEE REMARK 999	UNP P35475
A	105	GLN	ARG	SEE REMARK 999	UNP P35475
B	33	GLN	HIS	SEE REMARK 999	UNP P35475
B	63	PRO	GLN	SEE REMARK 999	UNP P35475
B	105	GLN	ARG	SEE REMARK 999	UNP P35475

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	E	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	F	7	Total	C	N	O	0	0	0
			83	46	2	35			

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	Cl	0	0
			1	1		
8	B	1	Total	Cl	0	0
			1	1		

- Molecule 9 is L(+)-TARTARIC ACID (CCD ID: TLA) (formula: C₄H₆O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	C	O	0	0
			10	4	6		
9	B	1	Total	C	O	0	0
			10	4	6		

- Molecule 10 is S,R MESO-TARTARIC ACID (CCD ID: SRT) (formula: C₄H₆O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	B	1	Total	C	O	0	0
			10	4	6		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	168	Total	O	0	0
			168	168		
11	B	265	Total	O	0	0
			265	265		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alpha-L-iduronidase



- Molecule 1: Alpha-L-iduronidase



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  80% 20%



- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  67% 33%



- Molecule 5: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  29% 57% 14%



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	259.36Å 259.36Å 71.81Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.01 – 2.10 49.01 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.5 (49.01-2.10) 99.5 (49.01-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.70 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.191 , 0.206 0.196 , 0.212	Depositor DCC
R_{free} test set	5233 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	29.3	Xtriage
Anisotropy	0.656	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 45.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.037 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10336	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, MAN, TLA, CL, BMA, GOL, SRT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/4890	0.65	0/6680
1	B	0.43	0/4999	0.67	2/6829 (0.0%)
All	All	0.42	0/9889	0.66	2/13509 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	429	GLY	CA-C-N	5.05	124.66	119.56
1	B	429	GLY	C-N-CA	5.05	124.66	119.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4745	0	4644	5	0
1	B	4851	0	4753	11	0
2	C	28	0	25	0	0
3	D	61	0	52	0	0
4	E	39	0	34	0	0
5	F	83	0	70	1	0
6	A	14	0	13	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	14	0	13	0	0
7	A	6	0	8	0	0
7	B	30	0	40	2	0
8	A	1	0	0	0	0
8	B	1	0	0	0	0
9	B	20	0	8	0	0
10	B	10	0	4	0	0
11	A	168	0	0	0	0
11	B	265	0	0	0	0
All	All	10336	0	9664	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:575:SER:OG	1:B:577[A]:CYS:SG	2.48	0.58
1:A:434:TRP:CH2	1:A:436:ALA:HB2	2.45	0.52
1:B:501:PHE:HA	1:B:504:MET:HE3	1.94	0.49
1:B:195:MET:HE1	1:B:236:GLY:HA3	1.98	0.46
1:B:434:TRP:CH2	1:B:436:ALA:HB2	2.51	0.46
1:A:383:ARG:CG	1:A:387:LEU:HD23	2.48	0.43
1:B:276:GLU:OE1	1:B:329:HIS:NE2	2.44	0.43
1:B:56:LEU:CD1	5:F:7:MAN:H3	2.49	0.42
1:B:334:LEU:HD21	1:B:343:TYR:HB3	2.02	0.41
1:B:570:ASP:CG	1:B:603:ASN:HD21	2.28	0.41
1:A:383:ARG:HG2	1:A:387:LEU:HD23	2.01	0.41
1:B:356:HIS:HB2	7:B:913:GOL:H32	2.02	0.41
1:B:471:TYR:HA	1:B:539:HIS:O	2.20	0.41
1:A:556:ALA:HB1	1:A:564:LEU:HD11	2.02	0.41
1:A:276:GLU:OE1	1:A:329:HIS:NE2	2.47	0.40
1:B:356:HIS:HB2	7:B:913:GOL:C3	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	588/653 (90%)	569 (97%)	19 (3%)	0	100	100
1	B	605/653 (93%)	586 (97%)	19 (3%)	0	100	100
All	All	1193/1306 (91%)	1155 (97%)	38 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	504/545 (92%)	501 (99%)	3 (1%)	78	86
1	B	516/545 (95%)	515 (100%)	1 (0%)	87	93
All	All	1020/1090 (94%)	1016 (100%)	4 (0%)	84	89

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33	GLN
1	A	564	LEU
1	A	598	LYS
1	B	99	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	GLN
1	A	185	HIS
1	A	196	GLN
1	A	282	GLN
1	A	330	GLN
1	A	358	HIS
1	B	113	HIS
1	B	185	HIS
1	B	196	GLN
1	B	262	HIS
1	B	285	GLN
1	B	330	GLN
1	B	348	ASN
1	B	561	GLN
1	B	603	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

17 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	1,2	14,14,15	0.53	0	17,19,21	0.89	1 (5%)
2	NAG	C	2	2	14,14,15	0.52	0	17,19,21	0.74	0
3	NAG	D	1	1,3	14,14,15	0.53	0	17,19,21	0.81	0
3	NAG	D	2	3	14,14,15	0.54	0	17,19,21	1.11	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BMA	D	3	3	11,11,12	0.29	0	15,15,17	0.70	0
3	MAN	D	4	3	11,11,12	0.59	0	15,15,17	0.56	0
3	MAN	D	5	3	11,11,12	0.56	0	15,15,17	0.69	0
4	NAG	E	1	1,4	14,14,15	0.50	0	17,19,21	1.07	1 (5%)
4	NAG	E	2	4	14,14,15	0.55	0	17,19,21	0.82	0
4	BMA	E	3	4	11,11,12	0.27	0	15,15,17	0.56	0
5	NAG	F	1	1,5	14,14,15	0.55	0	17,19,21	1.02	1 (5%)
5	NAG	F	2	5	14,14,15	0.62	0	17,19,21	0.88	1 (5%)
5	BMA	F	3	5	11,11,12	0.35	0	15,15,17	0.68	0
5	MAN	F	4	5	11,11,12	0.55	0	15,15,17	0.79	1 (6%)
5	MAN	F	5	5	11,11,12	0.71	0	15,15,17	1.01	1 (6%)
5	MAN	F	6	5	11,11,12	0.61	0	15,15,17	0.61	0
5	MAN	F	7	5	11,11,12	0.60	0	15,15,17	1.09	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
3	NAG	D	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	0/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1
3	MAN	D	4	3	-	2/2/19/22	0/1/1/1
3	MAN	D	5	3	-	2/2/19/22	0/1/1/1
4	NAG	E	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	E	2	4	-	0/6/23/26	0/1/1/1
4	BMA	E	3	4	-	0/2/19/22	0/1/1/1
5	NAG	F	1	1,5	-	0/6/23/26	0/1/1/1
5	NAG	F	2	5	-	0/6/23/26	0/1/1/1
5	BMA	F	3	5	-	0/2/19/22	0/1/1/1
5	MAN	F	4	5	-	2/2/19/22	0/1/1/1
5	MAN	F	5	5	-	2/2/19/22	0/1/1/1
5	MAN	F	6	5	-	0/2/19/22	0/1/1/1
5	MAN	F	7	5	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	2	NAG	C1-O5-C5	3.32	116.64	112.19
4	E	1	NAG	C1-O5-C5	3.03	116.25	112.19
5	F	5	MAN	C1-C2-C3	2.81	113.74	109.64
5	F	7	MAN	C1-C2-C3	2.63	113.47	109.64
5	F	1	NAG	O5-C1-C2	-2.41	107.57	111.29
5	F	7	MAN	C1-O5-C5	2.34	115.32	112.19
2	C	1	NAG	C1-O5-C5	2.32	115.30	112.19
5	F	4	MAN	C1-O5-C5	2.26	115.22	112.19
5	F	2	NAG	C1-O5-C5	2.11	115.02	112.19

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	2	NAG	O5-C5-C6-O6
3	D	4	MAN	O5-C5-C6-O6
3	D	4	MAN	C4-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6
5	F	5	MAN	O5-C5-C6-O6
5	F	5	MAN	C4-C5-C6-O6
3	D	5	MAN	O5-C5-C6-O6
3	D	5	MAN	C4-C5-C6-O6
5	F	4	MAN	C4-C5-C6-O6
5	F	4	MAN	O5-C5-C6-O6

There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	F	7	MAN	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.




Oligosaccharide Chain D



Oligosaccharide Chain E





5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 2 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	TLA	B	917	-	9,9,9	1.20	0	12,12,12	1.00	1 (8%)
9	TLA	B	919	-	9,9,9	1.13	0	12,12,12	1.07	2 (16%)
6	NAG	B	901	1	14,14,15	0.49	0	17,19,21	0.77	0
7	GOL	B	913	-	5,5,5	0.37	0	5,5,5	0.32	0
7	GOL	B	915	-	5,5,5	0.32	0	5,5,5	0.22	0
7	GOL	B	914	-	5,5,5	0.28	0	5,5,5	0.39	0
7	GOL	B	916	-	5,5,5	0.26	0	5,5,5	0.51	0
7	GOL	A	909	-	5,5,5	0.26	0	5,5,5	0.22	0
7	GOL	B	912	-	5,5,5	0.38	0	5,5,5	0.19	0
6	NAG	A	901	1	14,14,15	0.58	0	17,19,21	1.06	1 (5%)
10	SRT	B	918	-	9,9,9	1.17	0	12,12,12	1.04	1 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	TLA	B	917	-	-	0/12/12/12	-
9	TLA	B	919	-	-	3/12/12/12	-
6	NAG	B	901	1	-	0/6/23/26	0/1/1/1
7	GOL	B	913	-	-	0/4/4/4	-
7	GOL	B	915	-	-	0/4/4/4	-
7	GOL	B	914	-	-	2/4/4/4	-
7	GOL	B	916	-	-	0/4/4/4	-
7	GOL	A	909	-	-	2/4/4/4	-
7	GOL	B	912	-	-	4/4/4/4	-
6	NAG	A	901	1	-	0/6/23/26	0/1/1/1
10	SRT	B	918	-	-	12/12/12/12	-

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	901	NAG	C4-C3-C2	2.62	114.86	111.02
9	B	919	TLA	O41-C4-C3	2.18	119.37	113.31
10	B	918	SRT	O41-C4-C3	2.16	119.32	113.31
9	B	917	TLA	O11-C1-C2	2.03	118.95	113.31
9	B	919	TLA	O11-C1-C2	2.02	118.93	113.31

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	B	912	GOL	O1-C1-C2-C3
10	B	918	SRT	O1-C1-C2-O2
10	B	918	SRT	O11-C1-C2-O2
10	B	918	SRT	O2-C2-C3-O3
7	B	914	GOL	O1-C1-C2-O2
10	B	918	SRT	O2-C2-C3-C4
7	A	909	GOL	O1-C1-C2-C3
7	B	914	GOL	O1-C1-C2-C3
7	A	909	GOL	O1-C1-C2-O2
7	B	912	GOL	O1-C1-C2-O2
10	B	918	SRT	O3-C3-C4-O4
10	B	918	SRT	C1-C2-C3-O3
10	B	918	SRT	O3-C3-C4-O41
7	B	912	GOL	O2-C2-C3-O3
7	B	912	GOL	C1-C2-C3-O3
9	B	919	TLA	C2-C3-C4-O4
10	B	918	SRT	C1-C2-C3-C4
10	B	918	SRT	C2-C3-C4-O4
10	B	918	SRT	C2-C3-C4-O41
10	B	918	SRT	O11-C1-C2-C3
9	B	919	TLA	O3-C3-C4-O4
10	B	918	SRT	O1-C1-C2-C3
9	B	919	TLA	C2-C3-C4-O41

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	B	913	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	595/653 (91%)	0.42	36 (6%)	27 29	22, 36, 64, 86	1 (0%)
1	B	607/653 (92%)	0.11	26 (4%)	40 41	12, 30, 50, 68	4 (0%)
All	All	1202/1306 (92%)	0.26	62 (5%)	33 35	12, 33, 61, 86	5 (0%)

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	451	ASN	4.0
1	A	481	CYS	3.9
1	B	56	LEU	3.6
1	B	426	ARG	3.5
1	A	336	ASN	3.5
1	A	577	CYS	3.3
1	B	338	THR	3.2
1	B	521	GLY	3.2
1	B	189	ASP	3.0
1	A	639	LEU	3.0
1	A	338	THR	3.0
1	B	590	LYS	2.9
1	B	588	ASP	2.9
1	A	431	ALA	2.9
1	A	520	ALA	2.9
1	A	452	ARG	2.8
1	B	188	PHE	2.8
1	A	627	ALA	2.8
1	A	616	GLY	2.7
1	A	430	PRO	2.7
1	A	588	ASP	2.7
1	B	589	GLY	2.7
1	A	99	THR	2.7
1	B	642	PRO	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	375	ARG	2.6
1	A	64	TYR	2.6
1	B	107	LEU	2.5
1	A	531[A]	ARG	2.5
1	A	28	ALA	2.5
1	A	542	ALA	2.5
1	A	355	TYR	2.4
1	A	470	VAL	2.4
1	B	190	ASN	2.4
1	B	428	GLN	2.4
1	B	62	ASP	2.4
1	B	337	THR	2.4
1	B	250	GLY	2.3
1	B	103	THR	2.3
1	B	252	ALA	2.3
1	A	107	LEU	2.3
1	A	187	ASP	2.3
1	A	529	ALA	2.3
1	A	429	GLY	2.3
1	A	626	TRP	2.3
1	B	187	ASP	2.3
1	A	186	HIS	2.2
1	A	65	VAL	2.2
1	B	65	VAL	2.2
1	B	64	TYR	2.2
1	A	567	VAL	2.2
1	A	636	VAL	2.2
1	B	522	GLY	2.2
1	B	186	HIS	2.2
1	B	574	GLY	2.1
1	B	336	ASN	2.1
1	A	461	ARG	2.1
1	A	408	ALA	2.1
1	A	337	THR	2.1
1	A	454	VAL	2.0
1	A	339	SER	2.0
1	A	526	LEU	2.0
1	A	592	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MAN	D	4	11/12	0.51	0.19	80,84,86,87	0
4	BMA	E	3	11/12	0.53	0.18	85,86,87,88	0
5	MAN	F	7	11/12	0.55	0.20	75,77,79,80	0
5	MAN	F	5	11/12	0.61	0.21	72,74,77,79	0
3	MAN	D	5	11/12	0.67	0.18	73,73,74,75	0
4	NAG	E	2	14/15	0.73	0.16	68,74,76,81	0
2	NAG	C	2	14/15	0.73	0.14	66,72,73,74	0
5	MAN	F	4	11/12	0.83	0.16	57,58,61,66	0
5	MAN	F	6	11/12	0.85	0.13	58,61,66,72	0
3	NAG	D	2	14/15	0.88	0.12	47,48,55,55	0
4	NAG	E	1	14/15	0.88	0.11	43,48,55,62	0
2	NAG	C	1	14/15	0.89	0.10	44,47,53,59	0
3	BMA	D	3	11/12	0.90	0.13	60,65,70,74	0
5	BMA	F	3	11/12	0.90	0.11	42,46,51,53	0
3	NAG	D	1	14/15	0.91	0.10	37,41,44,44	0
5	NAG	F	2	14/15	0.92	0.09	33,35,38,40	0
5	NAG	F	1	14/15	0.97	0.06	27,30,32,32	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain D:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
9	TLA	B	919	10/10	0.79	0.17	66,68,71,71	0
6	NAG	A	901	14/15	0.82	0.12	51,57,60,60	0
7	GOL	B	912	6/6	0.85	0.16	42,43,43,43	0
10	SRT	B	918	10/10	0.86	0.11	50,53,53,54	0
7	GOL	B	913	6/6	0.87	0.14	39,40,41,41	0
7	GOL	A	909	6/6	0.88	0.14	46,48,49,52	0
7	GOL	B	914	6/6	0.88	0.13	37,38,38,38	0
6	NAG	B	901	14/15	0.90	0.10	35,40,43,43	0
7	GOL	B	915	6/6	0.91	0.13	48,50,53,54	0
9	TLA	B	917	10/10	0.92	0.09	37,39,40,42	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	GOL	B	916	6/6	0.95	0.08	31,32,33,34	0
8	CL	B	920	1/1	0.99	0.04	29,29,29,29	0
8	CL	A	910	1/1	0.99	0.04	31,31,31,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.