



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 09:32 AM UTC

PDB ID : 6MJ4 / pdb_00006mj4
Title : Crystal structure of MCD1D/INKTCR TERNARY COMPLEX bound to glycolipid (XXW)
Authors : Zajonc, D.M.; Bitra, A.; Janssens, J.
Deposited on : 2018-09-20
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	209	7% 93% . .
2	D	241	4% 93% 6% .
3	A	285	8% 86% 9% .
4	B	99	4% 91% 8% .
5	E	2	100%
6	F	3	67% 33%

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 7315 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called T cell receptor alpha variable 11,T cell receptor alpha variable 11,T cell receptor alpha joining 18,Human nkt tcr alpha chain, chimeric protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	201	Total	C	N	O	S	0	4	0
			1564	974	267	315	8			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	MET	-	initiating methionine	UNP A0A0B4J1J9
C	113	ILE	-	linker	UNP A0A0B4J1J9

- Molecule 2 is a protein called Beta-chain,T cell receptor chain,T cell receptor beta constant 2, CHIMERIC PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	239	Total	C	N	O	S	0	1	0
			1888	1184	336	362	6			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	95	ASP	-	linker	UNP A2NTY6
D	96	GLU	-	linker	UNP A2NTY6
D	97	GLY	-	linker	UNP A2NTY6
D	98	TYR	-	linker	UNP A2NTY6
D	130	ALA	ALA	linker	UNP A0N8J3
D	168	CYS	SER	variant	UNP A0A5B9
D	186	SER	CYS	variant	UNP A0A5B9

- Molecule 3 is a protein called Antigen-presenting glycoprotein CD1d1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	273	Total	C	N	O	S	0	4	0
			2205	1406	377	409	13			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	280	HIS	-	expression tag	UNP A0A0R4J090
A	281	HIS	-	expression tag	UNP A0A0R4J090
A	282	HIS	-	expression tag	UNP A0A0R4J090
A	283	HIS	-	expression tag	UNP A0A0R4J090
A	284	HIS	-	expression tag	UNP A0A0R4J090
A	285	HIS	-	expression tag	UNP A0A0R4J090

- Molecule 4 is a protein called Beta-2-microglobulin.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	B	98	Total	C	N	O	S	0	0	0
			793	506	134	146	7			

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	E	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 6 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	F	3	Total	C	N	O	0	0	0
			38	22	2	14			

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	C	1	Total	C	O	0	0
			6	3	3		
7	C	1	Total	C	O	0	0
			6	3	3		
7	C	1	Total	C	O	0	0
			6	3	3		
7	C	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	D	1	Total	Na	0	0
			1	1		

- Molecule 9 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 10 is N-[(2S,3S,4R)-3,4-dihydroxy-1-{[4-O-(prop-2-en-1-yl)-alpha-D-galactopyranosyl]oxy}octadecan-2-yl]hexacosanamide (CCD ID: JTG) (formula: C₅₃H₁₀₃NO₉).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
10	A	1	Total	C	N	O	0	0
			63	53	1	9		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	C	156	Total 156	O 156	0	0
11	D	246	Total 246	O 246	0	0
11	A	213	Total 213	O 213	0	0
11	B	70	Total 70	O 70	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: T cell receptor alpha variable 11,T cell receptor alpha variable 11,T cell receptor alpha joining 18,Human nkt tcr alpha chain, chimeric protein



- Molecule 2: Beta-chain,T cell receptor chain,T cell receptor beta constant 2, CHIMERIC PROTEIN



- Molecule 3: Antigen-presenting glycoprotein CD1d1



- Molecule 4: Beta-2-microglobulin



- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  100%

MAG1
MAG2

- Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  67% 33%

MAG1
MAG2
FUC3

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	78.95Å 190.72Å 151.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.10 – 2.00 47.10 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.6 (47.10-2.00) 99.6 (47.10-2.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0222	Depositor
R, R_{free}	0.185 , 0.214 (Not available) , 0.205	Depositor DCC
R_{free} test set	3839 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	28.0	Xtriage
Anisotropy	0.011	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 31.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7315	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, JTG, NA, GOL, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	C	0.57	0/1595	0.78	0/2171
2	D	0.57	0/1939	0.77	0/2642
3	A	0.55	0/2275	0.77	0/3095
4	B	0.56	0/819	0.76	0/1116
All	All	0.56	0/6628	0.77	0/9024

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1564	0	1498	4	0
2	D	1888	0	1794	8	0
3	A	2205	0	2108	19	0
4	B	793	0	746	7	0
5	E	28	0	25	0	0
6	F	38	0	34	0	0
7	A	12	0	16	2	0
7	C	24	0	32	0	0
8	D	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	A	14	0	13	0	0
10	A	63	0	0	0	0
11	A	213	0	0	0	0
11	B	70	0	0	2	0
11	C	156	0	0	0	0
11	D	246	0	0	1	0
All	All	7315	0	6266	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:301:GOL:H32	4:B:31:HIS:HB2	1.71	0.70
3:A:59:SER:H	3:A:62:GLN:HE21	1.40	0.70
3:A:215:PRO:O	3:A:267:HIS:HE1	1.84	0.61
1:C:22:GLN:OE1	1:C:74:THR:HG22	2.02	0.60
3:A:7:ASN:HD21	3:A:108:PRO:HD3	1.70	0.55
3:A:191:ALA:HA	3:A:209:HIS:O	2.08	0.54
3:A:59:SER:H	3:A:62:GLN:NE2	2.05	0.52
3:A:145:LEU:HB3	3:A:146:PRO:HD3	1.93	0.51
3:A:168:CYS:HB3	3:A:169:PRO:HD3	1.92	0.50
3:A:267:HIS:HD2	3:A:269:SER:OG	1.94	0.50
2:D:174:LEU:C	2:D:174:LEU:HD12	2.38	0.49
1:C:125:GLN:C	1:C:126:LEU:HD12	2.38	0.49
3:A:69:MET:HE2	3:A:69:MET:C	2.37	0.48
3:A:219:TRP:CZ2	3:A:221:MET:HG3	2.49	0.48
4:B:18:GLY:N	11:B:102:HOH:O	2.46	0.48
3:A:168:CYS:HB3	3:A:169:PRO:CD	2.44	0.47
3:A:219:TRP:HB3	3:A:266:LYS:HB2	1.95	0.47
3:A:258:GLU:HB3	3:A:279:TRP:CD1	2.50	0.47
3:A:200:ALA:HB1	3:A:203:HIS:H	1.81	0.46
4:B:67:HIS:HD2	11:B:105:HOH:O	1.98	0.46
1:C:126:LEU:HD12	1:C:126:LEU:N	2.32	0.45
2:D:78:LEU:HD12	2:D:78:LEU:N	2.31	0.45
3:A:188:LYS:HA	3:A:269:SER:OG	2.18	0.44
7:A:301:GOL:C3	4:B:31:HIS:HB2	2.44	0.44
2:D:217[B]:ASN:H	2:D:217[B]:ASN:ND2	2.15	0.43
4:B:21:ASN:HB3	4:B:70:PHE:CE1	2.54	0.43
3:A:59:SER:N	3:A:62:GLN:HE21	2.12	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:219:TRP:CH2	3:A:221:MET:HB2	2.54	0.42
3:A:81:ILE:HD12	3:A:98:ILE:CD1	2.49	0.42
2:D:115:ARG:NH1	2:D:116:ASN:HD21	2.18	0.42
4:B:4:THR:HA	4:B:5:PRO:HD3	1.97	0.42
2:D:2:ALA:N	11:D:407:HOH:O	2.54	0.41
2:D:176:GLU:OE2	2:D:186:SER:OG	2.38	0.41
2:D:208:GLN:HG3	2:D:231:ILE:HG23	2.03	0.41
3:A:100:LEU:HG	3:A:118:VAL:HG22	2.03	0.41
4:B:31:HIS:HA	4:B:32:PRO:C	2.46	0.40
1:C:32:HIS:CE1	2:D:98:TYR:CD1	3.09	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	179/188 (95%)	178 (99%)	1 (1%)	78	85
2	D	205/208 (99%)	202 (98%)	3 (2%)	57	64
3	A	240/249 (96%)	238 (99%)	2 (1%)	73	80
4	B	88/93 (95%)	88 (100%)	0	100	100
All	All	712/738 (96%)	706 (99%)	6 (1%)	70	80

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	181	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	129	LYS
2	D	168	CYS
2	D	184	ARG
3	A	69	MET
3	A	227	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	9	GLN
1	C	193	ASN
2	D	41	HIS
2	D	73	ASN
2	D	116	ASN
2	D	177	GLN
3	A	7	ASN
3	A	62	GLN
3	A	227	GLN
3	A	229	GLN
3	A	248	GLN
3	A	267	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	E	1	3,5	14,14,15	0.40	0	17,19,21	0.58	0
5	NAG	E	2	5	14,14,15	0.32	0	17,19,21	0.81	0
6	NAG	F	1	3,6	14,14,15	0.31	0	17,19,21	0.75	0
6	NAG	F	2	6	14,14,15	0.30	0	17,19,21	0.70	1 (5%)
6	FUC	F	3	6	10,10,11	0.51	0	14,14,16	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	E	1	3,5	-	0/6/23/26	0/1/1/1
5	NAG	E	2	5	-	2/6/23/26	0/1/1/1
6	NAG	F	1	3,6	-	0/6/23/26	0/1/1/1
6	NAG	F	2	6	-	2/6/23/26	0/1/1/1
6	FUC	F	3	6	-	-	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	F	2	NAG	C1-O5-C5	2.07	114.96	112.19

There are no chirality outliers.


All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	F	2	NAG	O5-C5-C6-O6
6	F	2	NAG	C4-C5-C6-O6
5	E	2	NAG	O5-C5-C6-O6
5	E	2	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
9	NAG	A	303	3	14,14,15	0.57	0	17,19,21	1.21	2 (11%)
10	JTG	A	309	-	63,63,63	0.71	1 (1%)	69,72,72	0.84	1 (1%)
7	GOL	A	302	-	5,5,5	0.28	0	5,5,5	0.28	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	GOL	C	602	-	5,5,5	0.32	0	5,5,5	0.17	0
7	GOL	A	301	-	5,5,5	0.37	0	5,5,5	0.49	0
7	GOL	C	603	-	5,5,5	0.32	0	5,5,5	0.13	0
7	GOL	C	601	-	5,5,5	0.35	0	5,5,5	0.63	0
7	GOL	C	604	-	5,5,5	0.34	0	5,5,5	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	NAG	A	303	3	-	0/6/23/26	0/1/1/1
10	JTG	A	309	-	-	11/62/82/82	0/1/1/1
7	GOL	A	302	-	-	2/4/4/4	-
7	GOL	C	602	-	-	4/4/4/4	-
7	GOL	A	301	-	-	2/4/4/4	-
7	GOL	C	603	-	-	2/4/4/4	-
7	GOL	C	601	-	-	1/4/4/4	-
7	GOL	C	604	-	-	3/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	A	309	JTG	CAX-CAU	3.69	1.52	1.29

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	A	303	NAG	C1-O5-C5	2.70	115.80	112.19
10	A	309	JTG	CAN-NAO-CBE	-2.50	119.24	123.40
9	A	303	NAG	C4-C3-C2	2.37	114.49	111.02

There are no chirality outliers.

All (25) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	C	602	GOL	C1-C2-C3-O3
7	A	301	GOL	O1-C1-C2-C3
7	A	301	GOL	O1-C1-C2-O2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
7	C	602	GOL	O1-C1-C2-C3
7	C	603	GOL	O1-C1-C2-C3
7	C	604	GOL	O1-C1-C2-C3
7	A	302	GOL	O1-C1-C2-C3
7	C	602	GOL	O2-C2-C3-O3
7	C	604	GOL	O1-C1-C2-O2
10	A	309	JTG	CCK-CCL-CCM-CCN
10	A	309	JTG	CCG-CCH-CCI-CCJ
10	A	309	JTG	CCA-CCB-CCC-CCD
10	A	309	JTG	CCE-CCF-CCG-CCH
10	A	309	JTG	CCB-CCC-CCD-CCE
7	C	603	GOL	O1-C1-C2-O2
10	A	309	JTG	O6-C2-C3-O2
10	A	309	JTG	CBP-CBQ-CBR-CBS
10	A	309	JTG	CBK-CBL-CBM-CBN
7	C	601	GOL	O2-C2-C3-O3
10	A	309	JTG	CAU-CAT-O4-C4
10	A	309	JTG	CCL-CCM-CCN-CCO
7	C	602	GOL	O1-C1-C2-O2
7	A	302	GOL	O1-C1-C2-O2
10	A	309	JTG	CCN-CCO-CCP-CCQ
7	C	604	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	301	GOL	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	C	201/209 (96%)	0.31	14 (6%)	22 21	9, 29, 60, 76	4 (1%)
2	D	239/241 (99%)	-0.02	10 (4%)	40 39	17, 27, 44, 62	1 (0%)
3	A	273/285 (95%)	0.25	22 (8%)	18 17	11, 28, 69, 93	4 (1%)
4	B	98/99 (98%)	0.44	4 (4%)	41 40	23, 36, 57, 76	0
All	All	811/834 (97%)	0.21	50 (6%)	26 25	9, 29, 60, 93	9 (1%)

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	B	4	THR	4.9
4	B	3	LYS	3.8
2	D	3	ALA	3.7
1	C	181	ASN	3.6
3	A	279	TRP	3.5
1	C	130	LYS	3.5
2	D	179	ALA	3.4
2	D	2	ALA	3.3
2	D	172	GLN	3.2
3	A	256	GLY	3.1
1	C	152	LYS	3.1
1	C	187	CYS	3.1
2	D	223	ASP	3.0
3	A	197	PRO	3.0
3	A	201	HIS	3.0
1	C	151	SER	2.7
4	B	99	MET	2.7
3	A	262	ALA	2.7
1	C	195	ILE	2.7
3	A	226	ASP	2.6
2	D	181	ASN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	A	223	MET	2.6
1	C	132	SER	2.6
1	C	133	ASP	2.6
2	D	180	LEU	2.6
1	C	150	GLN	2.5
1	C	115	ASP	2.5
3	A	253	VAL	2.5
3	A	233	HIS	2.5
3	A	198	SER	2.5
3	A	255	ALA	2.5
2	D	217[A]	ASN	2.5
3	A	110	ASN	2.4
3	A	108	PRO	2.4
2	D	182	ASP	2.3
3	A	230	GLN	2.3
1	C	2	THR	2.3
3	A	278	TYR	2.3
3	A	200	ALA	2.2
4	B	2	GLN	2.2
3	A	196	VAL	2.2
1	C	162	CYS	2.2
1	C	169	MET	2.2
1	C	196	ILE	2.1
3	A	109	GLY	2.1
3	A	22	SER	2.1
2	D	207	CYS	2.1
3	A	254	GLU	2.0
3	A	43	ASP	2.0
3	A	111	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	NAG	E	2	14/15	0.62	0.17	52,59,62,64	0
6	FUC	F	3	10/11	0.82	0.14	45,47,48,49	0
6	NAG	F	2	14/15	0.89	0.10	40,44,46,47	0
5	NAG	E	1	14/15	0.93	0.08	30,33,37,44	0
6	NAG	F	1	14/15	0.94	0.08	31,32,38,42	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	GOL	C	604	6/6	0.74	0.24	66,67,68,68	0
9	NAG	A	303	14/15	0.75	0.15	52,57,64,64	0
7	GOL	C	603	6/6	0.77	0.20	68,70,70,71	0
7	GOL	A	302	6/6	0.79	0.20	59,60,60,61	0
7	GOL	A	301	6/6	0.80	0.23	50,56,57,61	0
8	NA	D	301	1/1	0.82	0.30	43,43,43,43	0
7	GOL	C	601	6/6	0.85	0.19	34,38,38,38	0
7	GOL	C	602	6/6	0.85	0.15	45,49,50,52	0
10	JTG	A	309	63/63	0.94	0.10	19,29,38,41	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.