



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 12:25 PM UTC

PDB ID : 6MJQ / pdb_00006mjq
Title : Crystal structure of the mCD1d/xxp (JJ295) /iNKTCR ternary complex
Authors : Zajonc, D.M.; Bitra, A.; Janssens, J.
Deposited on : 2018-09-21
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	209	94%
1	G	209	95%
2	D	241	98%
2	H	241	98%
3	A	285	92%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	285	 91%5% .
4	B	99	 97%..
4	F	99	 97%..
5	I	2	 50%50%
5	K	2	 100%
5	L	2	 100%
6	J	4	 100%
7	M	5	 80%20%

2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 13257 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called T cell receptor alpha variable 11,T cell receptor alpha variable 11,T cell receptor alpha joining 18,Human nkt tcr alpha chain, chimeric protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	201	Total	C	N	O	S	0	1	0
			1543	957	265	313	8			
1	G	201	Total	C	N	O	S	0	1	0
			1542	957	263	314	8			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	MET	-	initiating methionine	UNP A0A0B4J1J9
C	113	ILE	-	linker	UNP A0A0B4J1J9
G	0	MET	-	initiating methionine	UNP A0A0B4J1J9
G	113	ILE	-	linker	UNP A0A0B4J1J9

- Molecule 2 is a protein called Beta-chain,T cell receptor chain,T cell receptor beta constant 2, CHIMERIC PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	240	Total	C	N	O	S	0	0	0
			1885	1182	334	363	6			
2	H	239	Total	C	N	O	S	0	1	0
			1884	1181	335	362	6			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	95	ASP	-	linker	UNP A2NTY6
D	96	GLU	-	linker	UNP A2NTY6
D	97	GLY	-	linker	UNP A2NTY6
D	98	TYR	-	linker	UNP A2NTY6
D	130	ALA	ALA	linker	UNP A0N8J3
D	168	CYS	SER	variant	UNP A0A5B9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	186	SER	CYS	variant	UNP A0A5B9
H	95	ASP	-	linker	UNP A2NTY6
H	96	GLU	-	linker	UNP A2NTY6
H	97	GLY	-	linker	UNP A2NTY6
H	98	TYR	-	linker	UNP A2NTY6
H	130	ALA	ALA	linker	UNP A0N8J3
H	168	CYS	SER	variant	UNP A0A5B9
H	186	SER	CYS	variant	UNP A0A5B9

- Molecule 3 is a protein called Antigen-presenting glycoprotein CD1d1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	274	Total	C	N	O	S	0	0	0
			2187	1394	376	404	13			
3	E	274	Total	C	N	O	S	0	0	0
			2193	1397	379	404	13			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	280	HIS	-	expression tag	UNP A0A0R4J090
A	281	HIS	-	expression tag	UNP A0A0R4J090
A	282	HIS	-	expression tag	UNP A0A0R4J090
A	283	HIS	-	expression tag	UNP A0A0R4J090
A	284	HIS	-	expression tag	UNP A0A0R4J090
A	285	HIS	-	expression tag	UNP A0A0R4J090
E	280	HIS	-	expression tag	UNP A0A0R4J090
E	281	HIS	-	expression tag	UNP A0A0R4J090
E	282	HIS	-	expression tag	UNP A0A0R4J090
E	283	HIS	-	expression tag	UNP A0A0R4J090
E	284	HIS	-	expression tag	UNP A0A0R4J090
E	285	HIS	-	expression tag	UNP A0A0R4J090

- Molecule 4 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	B	97	Total	C	N	O	S	0	0	0
			776	496	131	142	7			
4	F	97	Total	C	N	O	S	0	0	0
			776	496	131	142	7			


- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a

cetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	I	2	Total	C	N	O	0	0	0
			28	16	2	10			
5	K	2	Total	C	N	O	0	0	0
			28	16	2	10			
5	L	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 6 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	J	4	Total	C	N	O	0	0	0
			49	28	2	19			

- Molecule 7 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
7	M	5	Total	C	N	O	0	0	0
			60	34	2	24			

- Molecule 8 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	C	1	Total	C	O	0	0
			6	3	3		
8	E	1	Total	C	O	0	0
			6	3	3		
8	H	1	Total	C	O	0	0
			6	3	3		

- Molecule 9 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	D	1	Total	Na	0	0
			1	1		
9	H	2	Total	Na	0	0
			2	2		

- Molecule 10 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
10	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 11 is N-[(2S,3S,4R)-3,4-dihydroxy-1-[(4-O-{[4-(trifluoromethyl)phenyl]methyl}-alpha-D-galactopyranosyl)oxy]octadecan-2-yl]hexacosanamide (CCD ID: JUD) (formula: $C_{58}H_{104}F_3NO_9$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
11	A	1	Total	C	F	N	O	0
			71	58	3	1	9	
11	E	1	Total	C	F	N	O	0
			71	58	3	1	9	

- Molecule 12 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	C	14	Total 14	O 14	0	0
12	D	12	Total 12	O 12	0	0
12	A	15	Total 15	O 15	0	0
12	B	6	Total 6	O 6	0	0
12	E	15	Total 15	O 15	0	0
12	G	10	Total 10	O 10	0	0
12	H	17	Total 17	O 17	0	0
12	F	12	Total 12	O 12	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: T cell receptor alpha variable 11,T cell receptor alpha variable 11,T cell receptor alpha joining 18,Human nkt tcr alpha chain, chimeric protein

Chain C:  94%




- Molecule 1: T cell receptor alpha variable 11,T cell receptor alpha variable 11,T cell receptor alpha joining 18,Human nkt tcr alpha chain, chimeric protein

Chain G:  95%




- Molecule 2: Beta-chain,T cell receptor chain,T cell receptor beta constant 2, CHIMERIC PROTEIN

Chain D:  98%



- Molecule 2: Beta-chain,T cell receptor chain,T cell receptor beta constant 2, CHIMERIC PROTEIN

Chain H:  98%



- Molecule 3: Antigen-presenting glycoprotein CD1d1

Chain A:  92%





- Molecule 3: Antigen-presenting glycoprotein CD1d1

Chain E:  91% 5% .





- Molecule 4: Beta-2-microglobulin

Chain B:  97% ..



- Molecule 4: Beta-2-microglobulin

Chain F:  97% ..




- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  50% 50%



- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:  100%




- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L:  100%



- Molecule 6: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  100%

MAG1
MAG2
BMA3
FUC4

- Molecule 7: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M:



MAG1
MAG2
BMA3
MAN4
FUC5

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	78.97Å 191.86Å 150.96Å 90.00° 90.02° 90.00°	Depositor
Resolution (Å)	49.75 – 3.00 49.75 – 3.00	Depositor EDS
% Data completeness (in resolution range)	95.2 (49.75-3.00) 95.2 (49.75-3.00)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.73 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.8.0222	Depositor
R, R_{free}	0.215 , 0.258 0.218 , 0.259	Depositor DCC
R_{free} test set	2142 reflections (4.77%)	wwPDB-VP
Wilson B-factor (Å ²)	51.6	Xtriage
Anisotropy	0.054	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 15.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.428 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13257	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, MAN, GOL, NAG, JUD, FUC, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	C	0.53	0/1574	0.76	0/2140
1	G	0.52	0/1573	0.76	0/2140
2	D	0.54	0/1936	0.74	0/2639
2	H	0.53	0/1935	0.74	0/2638
3	A	0.52	0/2251	0.83	0/3062
3	E	0.52	0/2258	0.83	0/3071
4	B	0.55	0/802	0.72	0/1095
4	F	0.54	0/802	0.72	0/1095
All	All	0.53	0/13131	0.77	0/17880

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1543	0	1468	2	0
1	G	1542	0	1461	2	0
2	D	1885	0	1787	1	0
2	H	1884	0	1783	1	0
3	A	2187	0	2082	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	2193	0	2092	6	0
4	B	776	0	723	0	0
4	F	776	0	723	0	0
5	I	28	0	25	0	0
5	K	28	0	25	0	0
5	L	28	0	25	0	0
6	J	49	0	43	0	0
7	M	60	0	52	0	0
8	C	6	0	8	0	0
8	E	6	0	8	0	0
8	H	6	0	8	0	0
9	D	1	0	0	0	0
9	H	2	0	0	0	0
10	A	14	0	13	0	0
11	A	71	0	0	0	0
11	E	71	0	0	0	0
12	A	15	0	0	0	0
12	B	6	0	0	0	0
12	C	14	0	0	0	0
12	D	12	0	0	0	0
12	E	15	0	0	0	0
12	F	12	0	0	0	0
12	G	10	0	0	0	0
12	H	17	0	0	0	0
All	All	13257	0	12326	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:88:MET:HE1	3:E:145:LEU:HD23	1.76	0.67
3:A:88:MET:HE1	3:A:145:LEU:HD23	1.76	0.67
3:E:200:ALA:HB3	3:E:203:HIS:HB2	1.86	0.58
3:E:109:GLY:O	3:E:110:ASN:C	2.53	0.50
3:A:200:ALA:HB3	3:A:203:HIS:HB2	1.95	0.48
3:A:145:LEU:HB3	3:A:146:PRO:HD3	1.97	0.45
3:E:145:LEU:HB3	3:E:146:PRO:HD3	1.97	0.45
3:E:35:LEU:HD12	3:E:183:LEU:HD23	2.00	0.43
1:G:126:LEU:HD12	1:G:126:LEU:N	2.34	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:126:LEU:HD12	1:C:126:LEU:N	2.34	0.43
3:A:35:LEU:HD12	3:A:183:LEU:HD23	2.01	0.43
1:C:138:LEU:HD21	2:D:139:THR:HG21	2.01	0.42
3:A:216:LYS:N	3:A:217:PRO:CD	2.83	0.42
3:E:216:LYS:N	3:E:217:PRO:CD	2.83	0.42
1:G:138:LEU:HD21	2:H:139:THR:HG21	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	198/209 (95%)	184 (93%)	13 (7%)	1 (0%)	24	60
1	G	198/209 (95%)	183 (92%)	14 (7%)	1 (0%)	24	60
2	D	238/241 (99%)	233 (98%)	5 (2%)	0	100	100
2	H	238/241 (99%)	234 (98%)	4 (2%)	0	100	100
3	A	272/285 (95%)	257 (94%)	15 (6%)	0	100	100
3	E	272/285 (95%)	256 (94%)	14 (5%)	2 (1%)	18	53
4	B	95/99 (96%)	91 (96%)	4 (4%)	0	100	100
4	F	95/99 (96%)	91 (96%)	4 (4%)	0	100	100
All	All	1606/1668 (96%)	1529 (95%)	73 (4%)	4 (0%)	43	76

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	E	110	ASN
1	G	85	THR
1	C	85	THR
3	E	109	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	176/188 (94%)	175 (99%)	1 (1%)	78	88
1	G	176/188 (94%)	176 (100%)	0	100	100
2	D	204/208 (98%)	201 (98%)	3 (2%)	57	80
2	H	204/208 (98%)	201 (98%)	3 (2%)	57	80
3	A	235/249 (94%)	232 (99%)	3 (1%)	61	81
3	E	237/249 (95%)	232 (98%)	5 (2%)	47	75
4	B	85/93 (91%)	84 (99%)	1 (1%)	63	82
4	F	85/93 (91%)	84 (99%)	1 (1%)	63	82
All	All	1402/1476 (95%)	1385 (99%)	17 (1%)	63	82

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	1	LYS
2	D	25	GLN
2	D	168	CYS
2	D	235	GLU
3	A	6	LYS
3	A	227	GLN
3	A	276	ILE
4	B	45	LYS
3	E	110	ASN
3	E	173	ARG
3	E	201	HIS
3	E	227	GLN
3	E	276	ILE
2	H	25	GLN
2	H	168	CYS
2	H	235	GLU
4	F	45	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17)

such sidechains are listed below:

Mol	Chain	Res	Type
1	C	24	ASN
1	C	145	GLN
2	D	177	GLN
2	D	217	ASN
3	A	7	ASN
3	A	227	GLN
3	A	229	GLN
3	A	248	GLN
4	B	8	GLN
3	E	7	ASN
3	E	110	ASN
3	E	227	GLN
3	E	229	GLN
3	E	248	GLN
1	G	24	ASN
2	H	177	GLN
2	H	217	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

15 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	I	1	5,3	14,14,15	0.33	0	17,19,21	0.76	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	I	2	5	14,14,15	0.35	0	17,19,21	0.66	0
6	NAG	J	1	6,3	14,14,15	0.31	0	17,19,21	0.62	0
6	NAG	J	2	6	14,14,15	0.37	0	17,19,21	0.72	0
6	BMA	J	3	6	11,11,12	0.30	0	15,15,17	0.55	0
6	FUC	J	4	6	10,10,11	0.31	0	14,14,16	0.54	0
5	NAG	K	1	5,3	14,14,15	0.33	0	17,19,21	0.60	0
5	NAG	K	2	5	14,14,15	0.26	0	17,19,21	0.72	0
5	NAG	L	1	5,3	14,14,15	0.31	0	17,19,21	0.68	0
5	NAG	L	2	5	14,14,15	0.30	0	17,19,21	0.57	0
7	NAG	M	1	7,3	14,14,15	0.32	0	17,19,21	0.63	0
7	NAG	M	2	7	14,14,15	0.32	0	17,19,21	0.75	0
7	BMA	M	3	7	11,11,12	0.36	0	15,15,17	0.67	0
7	MAN	M	4	7	11,11,12	0.40	0	15,15,17	1.10	1 (6%)
7	FUC	M	5	7	10,10,11	0.32	0	14,14,16	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	I	1	5,3	-	0/6/23/26	0/1/1/1
5	NAG	I	2	5	-	2/6/23/26	0/1/1/1
6	NAG	J	1	6,3	-	0/6/23/26	0/1/1/1
6	NAG	J	2	6	-	0/6/23/26	0/1/1/1
6	BMA	J	3	6	-	0/2/19/22	0/1/1/1
6	FUC	J	4	6	-	-	0/1/1/1
5	NAG	K	1	5,3	-	0/6/23/26	0/1/1/1
5	NAG	K	2	5	-	0/6/23/26	0/1/1/1
5	NAG	L	1	5,3	-	0/6/23/26	0/1/1/1
5	NAG	L	2	5	-	0/6/23/26	0/1/1/1
7	NAG	M	1	7,3	-	0/6/23/26	0/1/1/1
7	NAG	M	2	7	-	2/6/23/26	0/1/1/1
7	BMA	M	3	7	-	0/2/19/22	0/1/1/1
7	MAN	M	4	7	-	0/2/19/22	0/1/1/1
7	FUC	M	5	7	-	-	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
7	M	4	MAN	C1-C2-C3	3.13	114.20	109.64
5	I	1	NAG	C1-O5-C5	2.27	115.22	112.19

There are no chirality outliers.


All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	M	2	NAG	O5-C5-C6-O6
5	I	2	NAG	O5-C5-C6-O6
7	M	2	NAG	C4-C5-C6-O6
5	I	2	NAG	C4-C5-C6-O6


There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 3 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
10	NAG	A	301	3	14,14,15	0.35	0	17,19,21	0.67	0
8	GOL	H	301	-	5,5,5	0.30	0	5,5,5	0.23	0
11	JUD	E	311	-	72,72,72	0.96	3 (4%)	84,87,87	0.79	0
8	GOL	C	301	-	5,5,5	0.29	0	5,5,5	0.19	0
8	GOL	E	310	-	5,5,5	0.27	0	5,5,5	0.26	0
11	JUD	A	308	-	72,72,72	0.97	3 (4%)	84,87,87	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	NAG	A	301	3	-	1/6/23/26	0/1/1/1
8	GOL	H	301	-	-	0/4/4/4	-
11	JUD	E	311	-	-	16/69/89/89	0/2/2/2
8	GOL	C	301	-	-	0/4/4/4	-
8	GOL	E	310	-	-	0/4/4/4	-
11	JUD	A	308	-	-	16/69/89/89	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	E	311	JUD	CAT-CAU	-4.26	1.40	1.50
11	A	308	JUD	CAT-CAU	-4.19	1.40	1.50
11	A	308	JUD	CBA-CAX	-4.11	1.41	1.49
11	E	311	JUD	CBA-CAX	-4.05	1.41	1.49
11	E	311	JUD	O1-C1	2.68	1.44	1.40
11	A	308	JUD	O1-C1	2.51	1.44	1.40

There are no bond angle outliers.

There are no chirality outliers.

All (33) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
11	A	308	JUD	O6-C2-C3-O2
11	E	311	JUD	CBN-CBO-CBP-CBQ
11	E	311	JUD	O6-C2-C3-O2
11	E	311	JUD	CBK-CBL-CBM-CBN
11	A	308	JUD	CBN-CBO-CBP-CBQ
11	A	308	JUD	CCK-CCL-CCM-CCN
11	A	308	JUD	CBK-CBL-CBM-CBN
11	A	308	JUD	CCG-CCH-CCI-CCJ
11	E	311	JUD	CCG-CCH-CCI-CCJ
11	A	308	JUD	CBP-CBQ-CBR-CBS
11	E	311	JUD	CCJ-CCK-CCL-CCM
11	E	311	JUD	CBW-CBX-CBY-CBZ
11	E	311	JUD	CCK-CCL-CCM-CCN
11	E	311	JUD	CBP-CBQ-CBR-CBS
11	A	308	JUD	CCC-CCD-CCE-CCF

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Atoms
11	A	308	JUD	CCA-CCB-CCC-CCD
11	A	308	JUD	CCJ-CCK-CCL-CCM
11	A	308	JUD	CBW-CBX-CBY-CBZ
11	A	308	JUD	CAU-CAT-O4-C4
11	E	311	JUD	CCA-CCB-CCC-CCD
11	E	311	JUD	CCC-CCD-CCE-CCF
10	A	301	NAG	C1-C2-N2-C7
11	E	311	JUD	CCO-CCP-CCQ-CCR
11	A	308	JUD	CCH-CCI-CCJ-CCK
11	E	311	JUD	CCH-CCI-CCJ-CCK
11	A	308	JUD	CBE-CBU-CBV-CBW
11	E	311	JUD	CBE-CBU-CBV-CBW
11	A	308	JUD	CCL-CCM-CCN-CCO
11	A	308	JUD	CAP-CAN-NAO-CBE
11	E	311	JUD	CAP-CAN-NAO-CBE
11	E	311	JUD	CAU-CAT-O4-C4
11	A	308	JUD	CCM-CCN-CCO-CCP
11	E	311	JUD	CCL-CCM-CCN-CCO

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	201/209 (96%)	-1.35	0 100 100	26, 53, 81, 87	1 (0%)
1	G	201/209 (96%)	-1.33	0 100 100	25, 51, 82, 88	1 (0%)
2	D	240/241 (99%)	-1.39	0 100 100	35, 49, 68, 78	0
2	H	239/241 (99%)	-1.36	0 100 100	24, 49, 67, 78	1 (0%)
3	A	274/285 (96%)	-1.34	0 100 100	35, 53, 90, 105	0
3	E	274/285 (96%)	-1.35	0 100 100	36, 53, 93, 117	0
4	B	97/99 (97%)	-1.30	0 100 100	43, 57, 76, 94	0
4	F	97/99 (97%)	-1.29	0 100 100	44, 59, 76, 93	0
All	All	1623/1668 (97%)	-1.35	0 100 100	24, 52, 81, 117	3 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	MAN	M	4	11/12	0.95	0.06	100,101,102,103	0
5	NAG	K	2	14/15	0.97	0.07	83,85,86,86	0
6	BMA	J	3	11/12	0.97	0.06	77,78,79,80	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	BMA	M	3	11/12	0.97	0.06	89,92,94,97	0
5	NAG	I	2	14/15	0.97	0.07	67,69,70,70	0
7	NAG	M	2	14/15	0.98	0.06	71,74,78,84	0
6	NAG	J	2	14/15	0.98	0.04	67,68,71,74	0
5	NAG	L	2	14/15	0.98	0.05	69,71,72,72	0
5	NAG	L	1	14/15	0.99	0.05	57,59,61,65	0
6	FUC	J	4	10/11	0.99	0.04	66,67,67,67	0
7	NAG	M	1	14/15	0.99	0.07	59,61,68,68	0
5	NAG	K	1	14/15	0.99	0.04	70,72,75,79	0
6	NAG	J	1	14/15	0.99	0.06	57,58,64,64	0
5	NAG	I	1	14/15	0.99	0.04	56,57,60,63	0
7	FUC	M	5	10/11	0.99	0.07	69,70,71,71	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain K:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain L:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain J:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain M:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	GOL	H	301	6/6	0.96	0.11	77,78,78,78	0
9	NA	H	302	1/1	0.98	0.12	57,57,57,57	0
10	NAG	A	301	14/15	0.98	0.04	64,65,65,65	0
9	NA	D	301	1/1	0.99	0.03	44,44,44,44	0
8	GOL	E	310	6/6	0.99	0.07	90,90,90,90	0
8	GOL	C	301	6/6	0.99	0.09	72,74,75,76	0
11	JUD	A	308	71/71	0.99	0.05	49,54,72,76	0
11	JUD	E	311	71/71	0.99	0.05	49,53,67,71	0
9	NA	H	303	1/1	1.00	0.04	44,44,44,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around JUD A 308:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around JUD E 311:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.