



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 14, 2026 – 11:04 PM UTC

PDB ID : 3MKA / pdb_00003mka
Title : Crystal Structure of Mycobacterium Tuberculosis Proteasome with propetide
and an T1A mutation at beta-subunit
Authors : Li, D.; Li, H.
Deposited on : 2010-04-14
Resolution : 2.51 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.










Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1	248	2% 51% 30% 7% 12%
1	A	248	% 47% 38% 5% 10%
1	B	248	48% 35% 6% 11%
1	D	248	2% 51% 30% 8% 10%
1	F	248	2% 51% 31% 6% 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	I	248	
1	K	248	
1	M	248	
1	O	248	
1	Q	248	
1	S	248	
1	U	248	
1	W	248	
1	Y	248	
2	2	291	
2	C	291	
2	E	291	
2	G	291	
2	H	291	
2	J	291	
2	L	291	
2	N	291	
2	P	291	
2	R	291	
2	T	291	
2	V	291	
2	X	291	
2	Z	291	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 49801 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	222	Total	C	N	O	S	0	0	0
			1713	1078	310	322	3			
1	B	220	Total	C	N	O	S	0	0	0
			1695	1064	308	320	3			
1	D	222	Total	C	N	O	S	0	0	0
			1717	1082	310	322	3			
1	F	221	Total	C	N	O	S	0	0	0
			1706	1073	309	321	3			
1	I	217	Total	C	N	O	S	0	0	0
			1680	1055	305	317	3			
1	K	221	Total	C	N	O	S	0	0	0
			1705	1072	309	321	3			
1	M	224	Total	C	N	O	S	0	0	0
			1730	1090	312	325	3			
1	O	218	Total	C	N	O	S	0	0	0
			1675	1049	306	317	3			
1	Q	222	Total	C	N	O	S	0	0	0
			1716	1081	310	322	3			
1	S	219	Total	C	N	O	S	0	0	0
			1686	1058	307	318	3			
1	U	221	Total	C	N	O	S	0	0	0
			1706	1073	309	321	3			
1	W	221	Total	C	N	O	S	0	0	0
			1710	1078	309	320	3			
1	Y	224	Total	C	N	O	S	0	0	0
			1730	1090	312	325	3			
1	1	219	Total	C	N	O	S	0	0	0
			1693	1065	307	318	3			

- Molecule 2 is a protein called Proteasome subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	250	Total	C	N	O	S	0	0	0
			1853	1165	320	363	5			
2	E	248	Total	C	N	O	S	0	0	0
			1834	1153	317	359	5			
2	G	251	Total	C	N	O	S	0	0	0
			1858	1168	321	364	5			
2	H	249	Total	C	N	O	S	0	0	0
			1844	1159	319	361	5			
2	J	252	Total	C	N	O	S	0	0	0
			1863	1170	322	366	5			
2	L	251	Total	C	N	O	S	0	0	0
			1857	1167	321	364	5			
2	N	252	Total	C	N	O	S	0	0	0
			1863	1170	322	366	5			
2	P	250	Total	C	N	O	S	0	0	0
			1853	1165	320	363	5			
2	R	246	Total	C	N	O	S	0	0	0
			1822	1145	315	357	5			
2	T	250	Total	C	N	O	S	0	0	0
			1854	1165	320	364	5			
2	V	249	Total	C	N	O	S	0	0	0
			1843	1158	319	361	5			
2	X	249	Total	C	N	O	S	0	0	0
			1848	1162	319	362	5			
2	Z	245	Total	C	N	O	S	0	0	0
			1817	1142	314	356	5			
2	2	246	Total	C	N	O	S	0	0	0
			1829	1150	316	358	5			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	301	ALA	THR	engineered mutation	UNP O33245
E	301	ALA	THR	engineered mutation	UNP O33245
G	301	ALA	THR	engineered mutation	UNP O33245
H	301	ALA	THR	engineered mutation	UNP O33245
J	301	ALA	THR	engineered mutation	UNP O33245
L	301	ALA	THR	engineered mutation	UNP O33245
N	301	ALA	THR	engineered mutation	UNP O33245
P	301	ALA	THR	engineered mutation	UNP O33245
R	301	ALA	THR	engineered mutation	UNP O33245
T	301	ALA	THR	engineered mutation	UNP O33245
V	301	ALA	THR	engineered mutation	UNP O33245
X	301	ALA	THR	engineered mutation	UNP O33245

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
Z	301	ALA	THR	engineered mutation	UNP O33245
2	301	ALA	THR	engineered mutation	UNP O33245

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	6	Total O 6 6	0	0
3	B	2	Total O 2 2	0	0
3	C	3	Total O 3 3	0	0
3	D	3	Total O 3 3	0	0
3	E	1	Total O 1 1	0	0
3	F	4	Total O 4 4	0	0
3	G	13	Total O 13 13	0	0
3	H	3	Total O 3 3	0	0
3	I	2	Total O 2 2	0	0
3	J	3	Total O 3 3	0	0
3	K	4	Total O 4 4	0	0
3	L	2	Total O 2 2	0	0
3	M	5	Total O 5 5	0	0
3	N	4	Total O 4 4	0	0
3	O	1	Total O 1 1	0	0
3	P	5	Total O 5 5	0	0
3	R	1	Total O 1 1	0	0
3	S	3	Total O 3 3	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	T	3	Total 3	O 3	0	0
3	U	2	Total 2	O 2	0	0
3	V	4	Total 4	O 4	0	0
3	W	6	Total 6	O 6	0	0
3	X	2	Total 2	O 2	0	0
3	Y	6	Total 6	O 6	0	0
3	Z	5	Total 5	O 5	0	0
3	2	5	Total 5	O 5	0	0
3	1	3	Total 3	O 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha








• Molecule 1: Proteasome subunit alpha




• Molecule 1: Proteasome subunit alpha




• Molecule 1: Proteasome subunit alpha






• Molecule 1: Proteasome subunit alpha

Chain M: 53% 31% 6% 10%




• Molecule 1: Proteasome subunit alpha


Chain O: 4% 46% 35% 6% 12%




• Molecule 1: Proteasome subunit alpha

Chain Q: 2% 43% 38% 8% 10%







• Molecule 1: Proteasome subunit alpha




• Molecule 1: Proteasome subunit alpha




• Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 1: Proteasome subunit alpha




- Molecule 2: Proteasome subunit beta




- Molecule 2: Proteasome subunit beta






• Molecule 2: Proteasome subunit beta




• Molecule 2: Proteasome subunit beta







• Molecule 2: Proteasome subunit beta




• Molecule 2: Proteasome subunit beta




• Molecule 2: Proteasome subunit beta





• Molecule 2: Proteasome subunit beta



• Molecule 2: Proteasome subunit beta



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	173.93Å 115.42Å 199.58Å 90.00° 112.89° 90.00°	Depositor
Resolution (Å)	29.77 – 2.51 29.77 – 2.51	Depositor EDS
% Data completeness (in resolution range)	93.9 (29.77-2.51) 94.1 (29.77-2.51)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.74 (at 2.51Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.217 , 0.247 0.215 , 0.245	Depositor DCC
R_{free} test set	11802 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	67.1	Xtriage
Anisotropy	0.083	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 62.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.023 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	49801	wwPDB-VP
Average B, all atoms (Å ²)	83.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	1	0.88	0/1720	1.12	7/2325 (0.3%)
1	A	0.92	0/1741	1.15	9/2353 (0.4%)
1	B	0.80	0/1722	1.14	15/2327 (0.6%)
1	D	0.87	2/1746 (0.1%)	1.21	17/2360 (0.7%)
1	F	0.90	1/1733 (0.1%)	1.19	11/2342 (0.5%)
1	I	0.91	2/1707 (0.1%)	1.21	11/2306 (0.5%)
1	K	0.85	0/1733	1.23	19/2342 (0.8%)
1	M	0.94	0/1759	1.17	9/2378 (0.4%)
1	O	0.84	0/1700	1.19	20/2297 (0.9%)
1	Q	0.86	0/1745	1.18	14/2359 (0.6%)
1	S	0.87	1/1712 (0.1%)	1.20	14/2313 (0.6%)
1	U	0.90	2/1733 (0.1%)	1.19	17/2342 (0.7%)
1	W	0.94	0/1739	1.17	13/2351 (0.6%)
1	Y	0.87	1/1759 (0.1%)	1.15	8/2378 (0.3%)
2	2	1.02	2/1858 (0.1%)	1.17	6/2520 (0.2%)
2	C	1.08	2/1882 (0.1%)	1.16	10/2553 (0.4%)
2	E	0.98	3/1863 (0.2%)	1.20	10/2527 (0.4%)
2	G	1.10	3/1887 (0.2%)	1.18	6/2560 (0.2%)
2	H	1.09	3/1872 (0.2%)	1.19	15/2538 (0.6%)
2	J	1.03	3/1892 (0.2%)	1.17	8/2566 (0.3%)
2	L	1.00	2/1886 (0.1%)	1.16	9/2558 (0.4%)
2	N	1.07	2/1892 (0.1%)	1.20	19/2566 (0.7%)
2	P	0.94	0/1882	1.17	13/2553 (0.5%)
2	R	0.94	3/1851 (0.2%)	1.14	6/2510 (0.2%)
2	T	1.05	1/1883 (0.1%)	1.16	6/2554 (0.2%)
2	V	1.10	5/1872 (0.3%)	1.19	9/2539 (0.4%)
2	X	1.01	2/1877 (0.1%)	1.19	7/2546 (0.3%)
2	Z	1.08	2/1846 (0.1%)	1.19	12/2503 (0.5%)
All	All	0.97	42/50492 (0.1%)	1.18	320/68366 (0.5%)

The worst 5 of 42 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	446	ALA	CA-CB	-8.40	1.40	1.53
1	Y	165	ASN	C-N	8.24	1.44	1.33
2	Z	450	MET	SD-CE	7.88	1.99	1.79
2	J	345	ILE	C-N	-7.84	1.22	1.33
1	U	165	ASN	C-N	7.25	1.43	1.33

The worst 5 of 320 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	-38	VAL	N-CA-C	-8.78	96.24	108.27
1	U	112	THR	N-CA-C	8.57	120.70	111.36
1	S	213	LEU	N-CA-C	-8.47	94.93	108.73
1	K	10	GLU	N-CA-C	8.40	120.43	111.28
1	Y	165	ASN	O-C-N	-8.40	113.42	122.07

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1	1693	0	1688	138	0
1	A	1713	0	1713	131	0
1	B	1695	0	1685	129	0
1	D	1717	0	1712	140	0
1	F	1706	0	1705	93	0
1	I	1680	0	1673	102	0
1	K	1705	0	1702	154	0
1	M	1730	0	1726	115	0
1	O	1675	0	1676	107	0
1	Q	1716	0	1710	151	0
1	S	1686	0	1685	107	0
1	U	1706	0	1705	161	0
1	W	1710	0	1705	128	0
1	Y	1730	0	1726	92	0
2	2	1829	0	1824	69	0
2	C	1853	0	1850	106	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	1834	0	1829	101	0
2	G	1858	0	1855	78	0
2	H	1844	0	1841	111	0
2	J	1863	0	1858	88	0
2	L	1857	0	1853	102	0
2	N	1863	0	1858	78	0
2	P	1853	0	1850	103	0
2	R	1822	0	1815	84	0
2	T	1854	0	1850	76	0
2	V	1843	0	1837	75	0
2	X	1848	0	1845	96	0
2	Z	1817	0	1810	86	0
3	1	3	0	0	3	0
3	2	5	0	0	0	0
3	A	6	0	0	3	0
3	B	2	0	0	0	0
3	C	3	0	0	1	0
3	D	3	0	0	1	0
3	E	1	0	0	0	0
3	F	4	0	0	1	0
3	G	13	0	0	0	0
3	H	3	0	0	1	0
3	I	2	0	0	1	0
3	J	3	0	0	1	0
3	K	4	0	0	0	0
3	L	2	0	0	1	0
3	M	5	0	0	5	0
3	N	4	0	0	0	0
3	O	1	0	0	1	0
3	P	5	0	0	0	0
3	R	1	0	0	0	0
3	S	3	0	0	0	0
3	T	3	0	0	0	0
3	U	2	0	0	0	0
3	V	4	0	0	1	0
3	W	6	0	0	0	0
3	X	2	0	0	0	0
3	Y	6	0	0	1	0
3	Z	5	0	0	0	0
All	All	49801	0	49586	2715	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

The worst 5 of 2715 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:Q:35:TYR:CE1	1:Q:37:GLY:HA3	1.51	1.43
1:O:35:TYR:CE1	1:O:37:GLY:HA3	1.56	1.40
2:H:407:TYR:CE1	2:H:417:ALA:HB3	1.59	1.36
1:S:35:TYR:CE1	1:S:37:GLY:HA3	1.56	1.36
1:B:35:TYR:CE1	1:B:37:GLY:HA3	1.60	1.35

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	1	215/248 (87%)	204 (95%)	10 (5%)	1 (0%)	24	43
1	A	218/248 (88%)	208 (95%)	10 (5%)	0	100	100
1	B	216/248 (87%)	205 (95%)	11 (5%)	0	100	100
1	D	218/248 (88%)	202 (93%)	16 (7%)	0	100	100
1	F	217/248 (88%)	208 (96%)	9 (4%)	0	100	100
1	I	213/248 (86%)	204 (96%)	9 (4%)	0	100	100
1	K	217/248 (88%)	206 (95%)	11 (5%)	0	100	100
1	M	220/248 (89%)	215 (98%)	5 (2%)	0	100	100
1	O	214/248 (86%)	202 (94%)	12 (6%)	0	100	100
1	Q	218/248 (88%)	205 (94%)	13 (6%)	0	100	100
1	S	215/248 (87%)	202 (94%)	13 (6%)	0	100	100
1	U	217/248 (88%)	203 (94%)	14 (6%)	0	100	100
1	W	217/248 (88%)	205 (94%)	11 (5%)	1 (0%)	24	43
1	Y	220/248 (89%)	212 (96%)	8 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	2	242/291 (83%)	241 (100%)	1 (0%)	0	100	100
2	C	246/291 (84%)	241 (98%)	5 (2%)	0	100	100
2	E	244/291 (84%)	239 (98%)	5 (2%)	0	100	100
2	G	247/291 (85%)	246 (100%)	1 (0%)	0	100	100
2	H	244/291 (84%)	239 (98%)	5 (2%)	0	100	100
2	J	248/291 (85%)	245 (99%)	3 (1%)	0	100	100
2	L	247/291 (85%)	240 (97%)	6 (2%)	1 (0%)	30	49
2	N	248/291 (85%)	244 (98%)	4 (2%)	0	100	100
2	P	246/291 (84%)	244 (99%)	2 (1%)	0	100	100
2	R	242/291 (83%)	240 (99%)	2 (1%)	0	100	100
2	T	246/291 (84%)	243 (99%)	3 (1%)	0	100	100
2	V	245/291 (84%)	244 (100%)	1 (0%)	0	100	100
2	X	245/291 (84%)	244 (100%)	1 (0%)	0	100	100
2	Z	241/291 (83%)	240 (100%)	1 (0%)	0	100	100
All	All	6466/7546 (86%)	6271 (97%)	192 (3%)	3 (0%)	100	100

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	W	4	PRO
2	L	-19	ALA
1	1	7	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	1	168/192 (88%)	147 (88%)	21 (12%)	4	9
1	A	171/192 (89%)	157 (92%)	14 (8%)	10	23
1	B	168/192 (88%)	151 (90%)	17 (10%)	7	15

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	171/192 (89%)	156 (91%)	15 (9%)	9	20
1	F	170/192 (88%)	153 (90%)	17 (10%)	7	15
1	I	167/192 (87%)	152 (91%)	15 (9%)	9	19
1	K	170/192 (88%)	152 (89%)	18 (11%)	6	14
1	M	173/192 (90%)	156 (90%)	17 (10%)	7	16
1	O	167/192 (87%)	151 (90%)	16 (10%)	8	17
1	Q	171/192 (89%)	150 (88%)	21 (12%)	4	10
1	S	168/192 (88%)	154 (92%)	14 (8%)	10	22
1	U	170/192 (88%)	152 (89%)	18 (11%)	6	14
1	W	170/192 (88%)	152 (89%)	18 (11%)	6	14
1	Y	173/192 (90%)	153 (88%)	20 (12%)	5	11
2	2	186/216 (86%)	172 (92%)	14 (8%)	12	26
2	C	188/216 (87%)	180 (96%)	8 (4%)	26	51
2	E	185/216 (86%)	167 (90%)	18 (10%)	8	17
2	G	188/216 (87%)	176 (94%)	12 (6%)	16	33
2	H	186/216 (86%)	179 (96%)	7 (4%)	29	56
2	J	189/216 (88%)	178 (94%)	11 (6%)	18	38
2	L	188/216 (87%)	175 (93%)	13 (7%)	14	30
2	N	189/216 (88%)	179 (95%)	10 (5%)	20	42
2	P	188/216 (87%)	176 (94%)	12 (6%)	16	33
2	R	184/216 (85%)	173 (94%)	11 (6%)	17	36
2	T	189/216 (88%)	174 (92%)	15 (8%)	11	24
2	V	186/216 (86%)	176 (95%)	10 (5%)	20	41
2	X	188/216 (87%)	171 (91%)	17 (9%)	9	19
2	Z	184/216 (85%)	168 (91%)	16 (9%)	9	21
All	All	4995/5712 (87%)	4580 (92%)	415 (8%)	10	22

5 of 415 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Q	171	TYR
1	U	59	ARG
1	1	30	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	Q	226	THR
1	S	92	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 129 such sidechains are listed below:

Mol	Chain	Res	Type
2	Z	456	GLN
2	2	415	GLN
1	K	129	HIS
1	K	101	ASN
2	2	456	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	H	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	H	-6:ALA	C	-5:GLN	N	4.40

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	1	219/248 (88%)	0.16	5 (2%) 61 57	48, 86, 121, 152	0
1	A	222/248 (89%)	0.15	3 (1%) 73 70	43, 81, 117, 145	0
1	B	220/248 (88%)	0.12	1 (0%) 87 85	48, 88, 127, 145	0
1	D	222/248 (89%)	0.36	5 (2%) 61 57	52, 92, 126, 144	0
1	F	221/248 (89%)	0.09	4 (1%) 67 64	45, 86, 126, 147	0
1	I	217/248 (87%)	0.05	2 (0%) 81 78	45, 80, 123, 149	0
1	K	221/248 (89%)	0.49	3 (1%) 73 70	48, 98, 135, 148	0
1	M	224/248 (90%)	0.09	1 (0%) 88 86	45, 78, 127, 147	0
1	O	218/248 (87%)	0.37	9 (4%) 41 37	50, 97, 137, 150	0
1	Q	222/248 (89%)	0.35	5 (2%) 61 57	48, 96, 131, 140	0
1	S	219/248 (88%)	0.09	3 (1%) 73 70	46, 85, 129, 140	0
1	U	221/248 (89%)	0.09	1 (0%) 87 85	49, 88, 130, 144	0
1	W	221/248 (89%)	-0.04	2 (0%) 81 78	48, 81, 125, 149	0
1	Y	224/248 (90%)	0.11	3 (1%) 75 71	53, 93, 132, 142	0
2	2	246/291 (84%)	-0.16	3 (1%) 76 73	48, 68, 97, 107	0
2	C	250/291 (85%)	-0.09	1 (0%) 88 86	51, 73, 111, 143	0
2	E	248/291 (85%)	0.08	7 (2%) 55 50	30, 77, 115, 132	0
2	G	251/291 (86%)	-0.13	1 (0%) 88 86	44, 67, 115, 141	0
2	H	249/291 (85%)	-0.08	4 (1%) 70 67	45, 71, 113, 136	0
2	J	252/291 (86%)	-0.21	3 (1%) 76 73	47, 70, 109, 135	0
2	L	251/291 (86%)	-0.07	4 (1%) 70 67	51, 75, 116, 140	0
2	N	252/291 (86%)	-0.20	3 (1%) 76 73	44, 70, 115, 144	0
2	P	250/291 (85%)	-0.07	3 (1%) 76 73	47, 75, 114, 144	0
2	R	246/291 (84%)	0.08	7 (2%) 55 50	30, 82, 118, 137	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
2	T	250/291 (85%)	-0.28	1 (0%) 88 86	40, 68, 110, 137	0
2	V	249/291 (85%)	-0.18	2 (0%) 82 80	47, 68, 112, 136	0
2	X	249/291 (85%)	-0.08	5 (2%) 65 61	44, 65, 115, 144	0
2	Z	245/291 (84%)	-0.02	7 (2%) 53 49	49, 73, 115, 131	0
All	All	6579/7546 (87%)	0.03	98 (1%) 72 68	30, 78, 124, 152	0

The worst 5 of 98 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	6	PHE	5.2
2	E	-19	ALA	4.4
2	X	351	VAL	4.2
1	F	235	VAL	4.0
2	V	-19	ALA	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.