



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 02:34 AM UTC

PDB ID : 3MKK / pdb_00003mkk
Title : The crystal structure of the D307A mutant of glycoside HYDROLASE (FAMILY 31) from Ruminococcus obeum ATCC 29174 in complex with isomaltose
Authors : Tan, K.; Tesar, C.; Freeman, L.; Babnigg, G.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2010-04-15
Resolution : 1.91 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1188 (1.92-1.92)
Clashscore	190562	1209 (1.92-1.92)
Ramachandran outliers	187476	1195 (1.92-1.92)
Sidechain outliers	187428	1195 (1.92-1.92)
RSRZ outliers	180081	1188 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	666	89% 9%
1	B	666	91% 9%
2	C	2	50% 50%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 11896 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called alpha-glucosidase GH31 family.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	664	Total	C	N	O	S	3	6	0
			5468	3510	895	1028	35			
1	B	664	Total	C	N	O	S	0	8	0
			5475	3518	897	1025	35			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP A5ZY13
A	-1	ASN	-	expression tag	UNP A5ZY13
A	0	ALA	-	expression tag	UNP A5ZY13
A	307	ALA	ASP	engineered mutation	UNP A5ZY13
B	-2	SER	-	expression tag	UNP A5ZY13
B	-1	ASN	-	expression tag	UNP A5ZY13
B	0	ALA	-	expression tag	UNP A5ZY13
B	307	ALA	ASP	engineered mutation	UNP A5ZY13

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-6)-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	C	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is alpha-D-glucopyranose (CCD ID: GLC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 4 is beta-D-glucopyranose (CCD ID: BGC) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			12	6	6		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	439	Total 439	O 439	0	0
5	B	468	Total 468	O 468	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: alpha-glucosidase GH31 family



- Molecule 1: alpha-glucosidase GH31 family



- Molecule 2: alpha-D-glucopyranose-(1-6)-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	68.45Å 125.51Å 87.87Å 90.00° 107.85° 90.00°	Depositor
Resolution (Å)	41.44 – 1.91 41.44 – 1.91	Depositor EDS
% Data completeness (in resolution range)	99.0 (41.44-1.91) 99.0 (41.44-1.91)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.27 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.170 , 0.218 0.193 , 0.236	Depositor DCC
R_{free} test set	5381 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	23.1	Xtriage
Anisotropy	0.621	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 26.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11896	wwPDB-VP
Average B, all atoms (Å ²)	9.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	2/5626 (0.0%)	0.86	1/7585 (0.0%)
1	B	0.83	0/5642	0.89	3/7603 (0.0%)
All	All	0.80	2/11268 (0.0%)	0.87	4/15188 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	508	VAL	CA-CB	7.11	1.57	1.54
1	A	59	GLY	N-CA	5.65	1.49	1.44

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	435	LEU	CA-C-N	-6.49	111.77	118.85
1	B	435	LEU	C-N-CA	-6.49	111.77	118.85
1	B	654	LYS	N-CA-C	5.52	118.45	110.28
1	A	428	ILE	N-CA-C	-5.02	105.81	110.53

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5468	0	5247	50	0
1	B	5475	0	5277	32	0
2	C	23	0	19	1	0
3	A	11	0	10	0	0
4	A	12	0	11	0	0
5	A	439	0	0	3	0
5	B	468	0	0	4	0
All	All	11896	0	10564	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:330:PHE:HB2	1:A:343:MET:HE3	1.39	1.05
1:A:330:PHE:CB	1:A:343:MET:HE3	2.02	0.89
1:B:422:LYS:H	1:B:427:HIS:HD2	1.21	0.84
1:A:422:LYS:H	1:A:427:HIS:HD2	1.28	0.80
1:B:136:GLU:HG3	5:B:857:HOH:O	1.83	0.77
1:A:189:ILE:HD11	1:A:494:ILE:HD12	1.71	0.72
1:B:403[A]:SER:HB3	1:B:410:MET:HE1	1.72	0.72
1:A:21:GLU:HG2	5:A:993:HOH:O	1.89	0.71
1:A:330:PHE:HB2	1:A:343:MET:CE	2.20	0.68
1:A:333:ASP:O	1:B:324:LYS:NZ	2.30	0.63
1:B:371:HIS:HE1	1:B:375:HIS:ND1	1.97	0.63
1:A:187:ASN:CB	1:A:494:ILE:HD11	2.30	0.62
1:A:362:HIS:NE2	1:A:371:HIS:HD2	1.97	0.62
1:A:38:PHE:CZ	1:A:132:ILE:HD11	2.37	0.60
1:A:166:GLN:NE2	1:A:184:TYR:OH	2.35	0.59
1:A:310:GLU:N	1:A:311:PRO:HA	2.18	0.58
1:B:362:HIS:NE2	1:B:371:HIS:HD2	2.02	0.57
1:B:422:LYS:N	1:B:427:HIS:HD2	1.99	0.57
1:A:494:ILE:HG23	5:A:887:HOH:O	2.03	0.56
1:A:371:HIS:HE1	1:A:375:HIS:ND1	2.03	0.56
1:A:187:ASN:HB3	1:A:494:ILE:HD11	1.87	0.56
1:A:659[B]:ARG:HB2	1:A:659[B]:ARG:HH21	1.71	0.55
1:B:371:HIS:CE1	1:B:375:HIS:ND1	2.75	0.55
1:A:118:MET:HG2	5:B:1061:HOH:O	2.07	0.55
1:A:309:ASN:ND2	1:A:403:SER:OG	2.40	0.54
1:A:422:LYS:N	1:A:427:HIS:HD2	2.02	0.54
1:A:189:ILE:HD11	1:A:494:ILE:CD1	2.36	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:607:ILE:CD1	1:A:633:ILE:HG22	2.39	0.53
1:A:326:PHE:CE2	1:A:343:MET:HE2	2.44	0.52
1:B:225:LYS:HE2	1:B:302:GLU:OE2	2.10	0.52
1:B:38:PHE:CZ	1:B:132:ILE:HD11	2.46	0.50
1:B:431:ASN:HD22	1:B:431:ASN:C	2.20	0.50
1:A:502:ASN:O	1:A:506:ARG:HG3	2.11	0.50
1:A:656:GLU:HB2	5:A:904:HOH:O	2.11	0.49
1:B:292:LYS:HE2	5:B:1083:HOH:O	2.12	0.49
1:B:576:ILE:HD12	1:B:586:GLU:HG2	1.96	0.47
1:A:576:ILE:CD1	1:A:586:GLU:HG3	2.45	0.47
1:B:643:TYR:CD2	1:B:643:TYR:C	2.91	0.47
1:A:157:PRO:HG3	1:A:445:TYR:CD1	2.50	0.46
1:A:326:PHE:CD2	1:A:343:MET:CE	2.98	0.46
1:A:326:PHE:CD2	1:A:343:MET:HE2	2.51	0.46
1:B:189:ILE:HD11	1:B:494:ILE:CG2	2.45	0.46
1:A:600:LEU:HD11	1:B:118:MET:HE1	1.98	0.45
1:B:576:ILE:CD1	1:B:586:GLU:HG2	2.47	0.45
1:B:93:VAL:O	1:B:98:THR:HA	2.16	0.45
1:B:422:LYS:H	1:B:427:HIS:CD2	2.14	0.45
1:A:371:HIS:CE1	1:A:375:HIS:ND1	2.84	0.45
1:B:142:ASP:O	1:B:146:GLN:HG3	2.17	0.45
1:A:422:LYS:H	1:A:427:HIS:CD2	2.19	0.44
1:A:330:PHE:CG	1:A:343:MET:HE3	2.52	0.44
1:B:431:ASN:C	1:B:431:ASN:ND2	2.75	0.44
1:A:59:GLY:HA2	1:A:433:LYS:HD3	1.99	0.44
1:B:460:ASP:HB3	1:B:600:LEU:HD11	2.00	0.44
1:A:501:ILE:O	1:A:504:ARG:HB3	2.18	0.43
1:B:92:ILE:HD12	1:B:92:ILE:N	2.32	0.43
1:B:50:VAL:O	1:B:62:LYS:HG2	2.18	0.43
1:A:609:SER:HA	1:A:635:TYR:CE2	2.54	0.43
1:B:182[A]:LYS:HG3	1:B:186:GLU:CD	2.43	0.43
2:C:1:BGC:H6C1	2:C:2:GLC:C5	2.49	0.43
1:A:348:LYS:HE2	1:A:348:LYS:HB3	1.83	0.43
1:A:38:PHE:HZ	1:A:132:ILE:HD11	1.82	0.43
1:B:310:GLU:N	1:B:311:PRO:HA	2.34	0.43
1:B:166:GLN:NE2	1:B:184:TYR:OH	2.52	0.42
1:A:270:VAL:HG12	1:A:271:TRP:H	1.84	0.42
1:A:560:GLN:O	1:A:561:ASN:HB2	2.19	0.42
1:A:355:GLU:O	1:A:359:ARG:HG3	2.19	0.42
1:B:156:ILE:HG22	1:B:398:ARG:HD3	2.01	0.42
1:B:417:TRP:CD1	1:B:419:GLY:H	2.37	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:110:THR:O	1:A:123:VAL:HA	2.20	0.41
1:A:306:ASN:OD1	1:A:401:MET:HE1	2.21	0.41
1:A:643:TYR:CD2	1:A:643:TYR:C	2.98	0.41
1:A:659[B]:ARG:HB2	1:A:659[B]:ARG:NH2	2.35	0.41
1:A:55:GLU:HB2	1:A:438:LEU:HD21	2.03	0.41
1:A:368:LYS:O	1:A:369:ILE:HD13	2.21	0.41
1:A:396:GLU:O	1:A:649:HIS:HA	2.21	0.41
1:A:344:GLN:HG3	5:B:1074:HOH:O	2.22	0.40
1:B:605:LEU:C	1:B:605:LEU:HD12	2.46	0.40
1:A:64:GLY:O	1:B:458:THR:HG21	2.22	0.40
1:B:1:MET:HB3	1:B:146:GLN:OE1	2.21	0.40
1:A:532:VAL:HG21	1:A:547:MET:HG3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	668/666 (100%)	639 (96%)	27 (4%)	2 (0%)	36	26
1	B	670/666 (101%)	649 (97%)	20 (3%)	1 (0%)	48	40
All	All	1338/1332 (100%)	1288 (96%)	47 (4%)	3 (0%)	43	34

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	58	ARG
1	A	270	VAL
1	B	270	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	577/574 (100%)	565 (98%)	12 (2%)	47	33
1	B	579/574 (101%)	564 (97%)	15 (3%)	40	24
All	All	1156/1148 (101%)	1129 (98%)	27 (2%)	43	30

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	192	ASP
1	A	270	VAL
1	A	284	GLU
1	A	348	LYS
1	A	355	GLU
1	A	380	TYR
1	A	494	ILE
1	A	501	ILE
1	A	508	VAL
1	A	654	LYS
1	A	655	LYS
1	A	656	GLU
1	B	18	GLU
1	B	25	GLU
1	B	136	GLU
1	B	192	ASP
1	B	211	GLU
1	B	270	VAL
1	B	281	LEU
1	B	320	LEU
1	B	335	GLU
1	B	368	LYS
1	B	369	ILE
1	B	457	THR
1	B	465	PHE
1	B	492	GLU
1	B	569	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	89	ASN
1	A	166	GLN
1	A	187	ASN
1	A	339	HIS
1	A	344	GLN
1	A	371	HIS
1	A	427	HIS
1	B	371	HIS
1	B	427	HIS
1	B	431	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BGC	C	1	2	12,12,12	0.64	0	17,17,17	3.09	8 (47%)
2	GLC	C	2	2	11,11,12	0.67	0	15,15,17	0.81	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	C	1	2	-	2/2/22/22	0/1/1/1
2	GLC	C	2	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	BGC	O5-C5-C4	6.22	120.91	109.70
2	C	1	BGC	O5-C1-C2	5.59	120.14	110.30
2	C	1	BGC	O4-C4-C3	5.30	122.87	110.38
2	C	1	BGC	O5-C5-C6	4.00	116.36	106.44
2	C	1	BGC	O1-C1-C2	3.88	120.23	108.98
2	C	1	BGC	O4-C4-C5	3.08	116.90	109.32
2	C	1	BGC	C6-C5-C4	2.47	119.08	113.02
2	C	1	BGC	C4-C3-C2	-2.15	107.06	110.83

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	1	BGC	O5-C5-C6-O6
2	C	1	BGC	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	BGC	1	0
2	C	2	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	BGC	A	802	-	12,12,12	0.71	0	17,17,17	1.61	3 (17%)
3	GLC	A	801	-	11,11,12	0.66	0	15,15,17	1.26	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BGC	A	802	-	-	2/2/22/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLC	A	801	-	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	801	GLC	C1-O5-C5	3.79	117.27	112.19
4	A	802	BGC	C4-C3-C2	-3.76	104.24	110.83
4	A	802	BGC	O2-C2-C3	2.68	116.69	110.38
4	A	802	BGC	O4-C4-C3	2.52	116.32	110.38

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	802	BGC	O5-C5-C6-O6
4	A	802	BGC	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	664/666 (99%)	-0.14	5 (0%) 82 88	2, 9, 17, 28	9 (1%)
1	B	664/666 (99%)	-0.34	6 (0%) 81 86	2, 7, 17, 28	10 (1%)
All	All	1328/1332 (99%)	-0.24	11 (0%) 82 88	2, 8, 17, 28	19 (1%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	281	LEU	4.1
1	B	341	TRP	3.5
1	A	341	TRP	2.9
1	A	463	LEU	2.8
1	B	0	ALA	2.7
1	B	21	GLU	2.6
1	A	25	GLU	2.5
1	B	653	ASP	2.5
1	A	211	GLU	2.4
1	B	629	ASN	2.3
1	B	628	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BGC	C	1	12/12	0.84	0.13	26,40,43,43	0
2	GLC	C	2	11/12	0.95	0.08	15,19,22,22	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	BGC	A	802	12/12	0.77	0.12	28,43,45,48	0
3	GLC	A	801	11/12	0.96	0.06	21,23,26,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.