



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 01:24 AM UTC

PDB ID : 3MKN / pdb_00003mkn
Title : Crystal structure of the E. coli pyrimidine nucleosidase YeiK bound to a competitive inhibitor
Authors : Garau, G.; Fornili, A.; Giabbai, B.; Degano, M.
Deposited on : 2010-04-15
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	316	12% 85% 9% ..
1	B	316	3% 88% 7% 5%
1	C	316	4% 88% .. 8%
1	D	316	6% 89% 7% .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9711 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative uncharacterized protein YeiK.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	303	Total	C	N	O	S	0	3	0
			2276	1444	379	433	20			
1	B	301	Total	C	N	O	S	0	5	0
			2287	1455	379	433	20			
1	C	292	Total	C	N	O	S	0	1	0
			2190	1391	364	415	20			
1	D	303	Total	C	N	O	S	0	4	0
			2298	1456	384	438	20			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP C3T3U2
A	-1	SER	-	expression tag	UNP C3T3U2
A	0	HIS	-	expression tag	UNP C3T3U2
B	-2	GLY	-	expression tag	UNP C3T3U2
B	-1	SER	-	expression tag	UNP C3T3U2
B	0	HIS	-	expression tag	UNP C3T3U2
C	-2	GLY	-	expression tag	UNP C3T3U2
C	-1	SER	-	expression tag	UNP C3T3U2
C	0	HIS	-	expression tag	UNP C3T3U2
D	-2	GLY	-	expression tag	UNP C3T3U2
D	-1	SER	-	expression tag	UNP C3T3U2
D	0	HIS	-	expression tag	UNP C3T3U2

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		
2	B	1	Total	Ca	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	1	Total	Ca	0	0
			1	1		
2	D	1	Total	Ca	0	0
			1	1		

- Molecule 3 is (2S,3S,4R,5R)-2-(3,4-diaminophenyl)-5-(hydroxymethyl)pyrrolidine-3,4-diol (CCD ID: DNB) (formula: C₁₁H₁₇N₃O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	11	3	3		
3	B	1	Total	C	N	O	0	0
			17	11	3	3		
3	C	1	Total	C	N	O	0	0
			17	11	3	3		
3	D	1	Total	C	N	O	0	0
			17	11	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	101	Total	O	0	0
			101	101		
4	B	221	Total	O	0	0
			221	221		
4	C	97	Total	O	0	0
			97	97		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	169	Total 169	O 169	0	0



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	84.85Å 86.45Å 97.82Å 90.00° 98.86° 90.00°	Depositor
Resolution (Å)	44.97 – 2.00 44.97 – 2.00	Depositor EDS
% Data completeness (in resolution range)	97.3 (44.97-2.00) 97.3 (44.97-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.37 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.174 , 0.196 0.175 , 0.195	Depositor DCC
R_{free} test set	4602 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	23.7	Xtriage
Anisotropy	0.576	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 54.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9711	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DNB, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.32	16/2330 (0.7%)	0.99	9/3176 (0.3%)
1	B	0.92	0/2357	0.90	1/3209 (0.0%)
1	C	0.71	0/2234	0.83	0/3043
1	D	0.85	0/2359	0.93	3/3213 (0.1%)
All	All	0.98	16/9280 (0.2%)	0.92	13/12641 (0.1%)

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	204	ARG	CZ-NH1	24.50	1.67	1.32
1	A	204	ARG	CD-NE	22.70	1.78	1.46
1	A	204	ARG	CZ-NH2	16.60	1.55	1.33
1	A	205	MET	CG-SD	14.00	2.15	1.80
1	A	208	ALA	C-N	12.42	1.53	1.33
1	A	213	GLY	C-O	11.99	1.38	1.23
1	A	208	ALA	C-O	11.56	1.38	1.24
1	A	205	MET	SD-CE	10.74	2.06	1.79
1	A	307	ARG	CZ-NH1	9.55	1.46	1.32
1	A	204	ARG	NE-CZ	7.80	1.41	1.33
1	A	204	ARG	CG-CD	7.73	1.75	1.52
1	A	207	ARG	CZ-NH1	7.50	1.43	1.32
1	A	310	ILE	C-N	7.21	1.43	1.33
1	A	213	GLY	C-N	6.84	1.43	1.34
1	A	207	ARG	C-O	5.64	1.30	1.24
1	A	203	ALA	C-O	5.29	1.30	1.24

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	204	ARG	NE-CZ-NH2	-14.02	106.58	119.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	204	ARG	CD-NE-CZ	-11.44	108.38	124.40
1	A	204	ARG	CG-CD-NE	-7.42	95.68	112.00
1	B	227	GLN	N-CA-C	-6.81	105.60	114.04
1	A	205	MET	CG-SD-CE	-6.76	86.02	100.90
1	A	204	ARG	NH1-CZ-NH2	6.74	128.06	119.30
1	D	236	GLY	CA-C-N	6.04	126.00	119.78
1	D	236	GLY	C-N-CA	6.04	126.00	119.78
1	A	213	GLY	O-C-N	5.75	127.70	122.18
1	A	208	ALA	O-C-N	5.72	128.91	122.22
1	A	204	ARG	CB-CG-CD	-5.22	99.30	111.30
1	D	86	GLY	N-CA-C	-5.15	107.96	115.63
1	A	228	PHE	N-CA-C	-5.07	106.60	112.89

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2276	0	2272	29	1
1	B	2287	0	2307	20	1
1	C	2190	0	2201	7	0
1	D	2298	0	2302	13	1
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	17	0	16	2	0
3	B	17	0	16	2	0
3	C	17	0	17	0	0
3	D	17	0	16	3	0
4	A	101	0	0	2	0
4	B	221	0	0	8	0
4	C	97	0	0	0	0
4	D	169	0	0	3	0
All	All	9711	0	9147	71	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:204:ARG:CD	1:A:204:ARG:CG	1.75	1.58
1:A:204:ARG:CD	1:A:204:ARG:NE	1.78	1.45
1:A:205:MET:SD	1:A:205:MET:CE	2.06	1.44
1:A:205:MET:SD	1:A:205:MET:CG	2.15	1.34
1:B:15[C]:ASP:OD2	1:B:238:VAL:HA	1.73	0.88
4:B:434:HOH:O	1:D:280:VAL:HG11	1.76	0.83
1:B:15[A]:ASP:HB2	4:B:394:HOH:O	1.78	0.81
1:A:204:ARG:CD	1:A:204:ARG:CZ	2.59	0.80
1:A:204:ARG:CD	1:A:204:ARG:CB	2.59	0.80
1:A:136:MET:HE1	1:B:136:MET:HE1	1.76	0.67
1:C:136:MET:HE1	1:D:136:MET:HE1	1.77	0.67
1:B:15[B]:ASP:HB2	4:B:394:HOH:O	1.93	0.67
1:B:228:PHE:HA	1:B:232:GLY:CA	2.27	0.65
1:A:204:ARG:CG	1:A:204:ARG:NE	2.62	0.63
1:A:195:THR:OG1	1:A:238:VAL:O	2.20	0.60
1:B:228:PHE:HA	1:B:232:GLY:HA2	1.82	0.60
1:D:195:THR:OG1	1:D:238:VAL:O	2.18	0.60
1:D:62:PRO:CG	4:D:374:HOH:O	2.50	0.59
1:A:215:LEU:HG	1:A:219:ILE:HD12	1.84	0.58
1:A:81:ILE:HG13	3:A:502:DNB:N4	2.19	0.58
1:D:62:PRO:HG2	4:D:374:HOH:O	2.05	0.56
1:C:291:ILE:HG22	1:C:292:THR:HG23	1.87	0.56
1:A:248:ILE:HG21	1:A:301[A]:LEU:HD11	1.87	0.55
1:A:196:VAL:HA	1:A:237:PRO:HA	1.87	0.55
1:A:90:PRO:HG3	1:A:219:ILE:HG13	1.89	0.54
4:B:434:HOH:O	1:D:280:VAL:CG1	2.47	0.53
1:A:196:VAL:HA	1:A:237:PRO:N	2.23	0.53
1:B:15[C]:ASP:HB3	4:B:394:HOH:O	2.09	0.53
1:B:291:ILE:HG22	1:B:292:THR:HG23	1.89	0.52
1:B:55:GLN:HA	1:B:97[A]:ARG:HE	1.74	0.52
1:B:222:PHE:O	1:B:226:THR:HG22	2.09	0.52
1:A:205:MET:CE	1:A:205:MET:CG	2.87	0.52
1:D:291:ILE:HG22	1:D:292:THR:HG23	1.91	0.51
1:A:205:MET:SD	1:A:205:MET:CB	2.98	0.50
1:D:191:LEU:HD11	1:D:253:ILE:HG23	1.93	0.50
1:A:195:THR:O	1:A:237:PRO:HA	2.12	0.49
1:A:97:ARG:NH1	4:A:391:HOH:O	2.41	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:15[C]:ASP:CG	4:B:447:HOH:O	2.55	0.49
1:C:248:ILE:HG21	1:C:301:LEU:HD11	1.95	0.49
1:A:198:THR:HB	1:A:199:PRO:HD2	1.94	0.48
1:A:196:VAL:HA	1:A:237:PRO:CA	2.42	0.48
1:B:15[A]:ASP:OD2	1:B:239:HIS:HB2	2.13	0.48
1:D:62:PRO:HG3	4:D:374:HOH:O	2.11	0.47
1:B:191[A]:LEU:HD11	1:B:253:ILE:HG23	1.95	0.47
1:A:291:ILE:HG22	1:A:292:THR:HG23	1.95	0.47
1:A:59:ILE:O	1:A:97:ARG:NH1	2.46	0.46
1:A:191:LEU:HD11	1:A:253:ILE:HG23	1.98	0.46
3:B:502:DNB:H2'	3:B:502:DNB:H2	1.58	0.46
3:D:502:DNB:H2	3:D:502:DNB:H2'	1.76	0.45
1:A:14:HIS:CD2	1:A:219:ILE:HG23	2.52	0.45
1:A:14:HIS:HD2	1:A:219:ILE:HG23	1.83	0.44
1:D:81:ILE:HG13	3:D:502:DNB:HN4A	1.83	0.44
1:B:55:GLN:HB2	1:B:97[A]:ARG:HG2	1.98	0.44
1:D:191:LEU:C	1:D:191:LEU:HD23	2.43	0.44
1:D:198:THR:OG1	1:D:200:ASP:OD1	2.33	0.44
3:A:502:DNB:H2	3:A:502:DNB:H2'	1.85	0.43
1:C:191:LEU:HD11	1:C:253:ILE:HG23	2.00	0.43
1:C:47:LEU:HD11	1:C:99:ALA:HB2	2.01	0.43
1:C:12:PRO:HA	1:C:13:GLY:HA3	1.78	0.43
3:B:502:DNB:C2	3:B:502:DNB:H5'	2.50	0.42
1:A:135:ARG:HD3	1:A:135:ARG:HA	1.87	0.42
1:A:195:THR:C	1:A:237:PRO:HA	2.45	0.42
1:C:236:GLY:HA2	1:C:237:PRO:HD3	1.87	0.42
1:B:52:ASN:ND2	4:B:314:HOH:O	2.52	0.41
1:B:91:VAL:HG22	4:B:471:HOH:O	2.20	0.41
1:D:240:ASP:OD1	3:D:502:DNB:O3'	2.38	0.41
1:B:47:LEU:HD11	1:B:99:ALA:HB2	2.04	0.40
1:B:191[A]:LEU:CD1	1:B:253:ILE:HG23	2.52	0.40
1:A:97:ARG:HD3	4:A:391:HOH:O	2.21	0.40
1:B:25:LYS:CG	1:B:308:GLY:HA3	2.51	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:144:ILE:O	1:A:207:ARG:NH2[2_646]	1.95	0.25
1:B:60:ASN:O	1:D:101:SER:OG[1_655]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	302/316 (96%)	294 (97%)	7 (2%)	1 (0%)	36	35
1	B	303/316 (96%)	293 (97%)	9 (3%)	1 (0%)	36	35
1	C	287/316 (91%)	280 (98%)	7 (2%)	0	100	100
1	D	303/316 (96%)	293 (97%)	9 (3%)	1 (0%)	36	35
All	All	1195/1264 (94%)	1160 (97%)	32 (3%)	3 (0%)	36	35

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	232	GLY
1	A	39	GLY
1	D	39	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	248/259 (96%)	244 (98%)	4 (2%)	55	62
1	B	252/259 (97%)	250 (99%)	2 (1%)	73	80
1	C	239/259 (92%)	238 (100%)	1 (0%)	84	89
1	D	253/259 (98%)	249 (98%)	4 (2%)	55	62
All	All	992/1036 (96%)	981 (99%)	11 (1%)	65	73

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	80	ASN
1	A	97	ARG
1	A	190	ASP
1	A	207	ARG
1	B	190	ASP
1	B	231	TYR
1	C	190	ASP
1	D	91	VAL
1	D	97	ARG
1	D	190	ASP
1	D	196	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	52	ASN
1	A	60	ASN
1	A	265	ASN
1	B	52	ASN
1	B	265	ASN
1	C	52	ASN
1	C	265	ASN
1	D	52	ASN
1	D	265	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	DNB	D	502	2	18,18,18	1.23	3 (16%)	22,26,26	2.00	8 (36%)
3	DNB	C	502	2	18,18,18	1.11	1 (5%)	22,26,26	1.59	3 (13%)
3	DNB	B	502	2	18,18,18	1.15	1 (5%)	22,26,26	1.84	4 (18%)
3	DNB	A	502	2	18,18,18	1.18	2 (11%)	22,26,26	2.02	5 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DNB	D	502	2	-	3/6/22/22	0/2/2/2
3	DNB	C	502	2	-	4/6/22/22	0/2/2/2
3	DNB	B	502	2	-	4/6/22/22	0/2/2/2
3	DNB	A	502	2	-	4/6/22/22	0/2/2/2

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	502	DNB	C1-C1'	-3.29	1.47	1.52
3	A	502	DNB	C1-C1'	-3.11	1.47	1.52
3	C	502	DNB	C1-C1'	-2.97	1.48	1.52
3	B	502	DNB	C1-C1'	-2.90	1.48	1.52
3	A	502	DNB	C4'-N4'	-2.30	1.44	1.48
3	D	502	DNB	C4'-N4'	-2.25	1.44	1.48
3	D	502	DNB	C5'-C4'	2.11	1.56	1.52

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	DNB	C1-C1'-N4'	-6.64	102.44	112.98
3	B	502	DNB	C1-C1'-N4'	-5.83	103.74	112.98

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	502	DNB	C1-C1'-N4'	-5.28	104.61	112.98
3	D	502	DNB	C1-C1'-N4'	-4.61	105.66	112.98
3	D	502	DNB	C2'-C3'-C4'	3.81	108.33	102.55
3	A	502	DNB	C2'-C3'-C4'	3.03	107.15	102.55
3	D	502	DNB	C6-C1-C1'	-2.94	116.10	120.78
3	D	502	DNB	C3-C2-C1	-2.90	117.49	121.42
3	D	502	DNB	C6-C5-C4	-2.78	118.63	121.42
3	B	502	DNB	C6-C1-C2	2.65	121.81	118.74
3	A	502	DNB	O3'-C3'-C4'	-2.61	105.95	112.92
3	D	502	DNB	O3'-C3'-C4'	-2.58	106.02	112.92
3	D	502	DNB	C6-C1-C2	2.56	121.70	118.74
3	A	502	DNB	C6-C1-C1'	-2.51	116.79	120.78
3	B	502	DNB	C5-C6-C1	-2.49	118.69	121.18
3	D	502	DNB	C5-C4-C3	2.38	121.26	118.71
3	C	502	DNB	C2'-C3'-C4'	2.37	106.15	102.55
3	C	502	DNB	O3'-C3'-C4'	-2.34	106.66	112.92
3	B	502	DNB	C2'-C3'-C4'	2.30	106.05	102.55
3	A	502	DNB	C6-C5-C4	-2.01	119.40	121.42

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502	DNB	C6-C1-C1'-C2'
3	A	502	DNB	C2-C1-C1'-C2'
3	B	502	DNB	C6-C1-C1'-C2'
3	B	502	DNB	C2-C1-C1'-C2'
3	C	502	DNB	C6-C1-C1'-C2'
3	C	502	DNB	C2-C1-C1'-C2'
3	D	502	DNB	C6-C1-C1'-C2'
3	D	502	DNB	C2-C1-C1'-C2'
3	C	502	DNB	C6-C1-C1'-N4'
3	C	502	DNB	C2-C1-C1'-N4'
3	A	502	DNB	C2-C1-C1'-N4'
3	B	502	DNB	C2-C1-C1'-N4'
3	A	502	DNB	C6-C1-C1'-N4'
3	B	502	DNB	C6-C1-C1'-N4'
3	D	502	DNB	C2-C1-C1'-N4'

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	502	DNB	3	0
3	B	502	DNB	2	0
3	A	502	DNB	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	303/316 (95%)	0.35	39 (12%) 7 6	11, 39, 87, 110	5 (1%)
1	B	301/316 (95%)	-0.57	10 (3%) 49 48	8, 15, 38, 71	5 (1%)
1	C	292/316 (92%)	0.07	14 (4%) 35 34	13, 35, 82, 96	2 (0%)
1	D	303/316 (95%)	-0.26	18 (5%) 28 27	10, 22, 51, 83	5 (1%)
All	All	1199/1264 (94%)	-0.10	81 (6%) 23 22	8, 25, 77, 110	17 (1%)

All (81) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	91	VAL	6.6
1	D	81	ILE	6.1
1	A	238	VAL	5.8
1	C	78	ALA	5.5
1	D	80	ASN	5.3
1	A	91	VAL	5.3
1	A	228	PHE	5.1
1	A	229	GLU	4.8
1	A	311	LYS	4.8
1	D	82	HIS	4.7
1	A	310	ILE	4.5
1	B	226	THR	4.4
1	A	237	PRO	4.3
1	D	86	GLY	4.3
1	D	79	ASP	4.3
1	A	198	THR	4.2
1	C	222	PHE	4.0
1	D	85	THR	3.9
1	A	207	ARG	3.9
1	D	83	GLY	3.8
1	D	78	ALA	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	85	THR	3.7
1	A	202	ILE	3.6
1	B	78	ALA	3.6
1	D	228	PHE	3.6
1	A	196	VAL	3.5
1	A	226	THR	3.5
1	B	232	GLY	3.5
1	C	86	GLY	3.5
1	D	84	ASP	3.4
1	A	82	HIS	3.4
1	A	209	GLY	3.4
1	A	93	GLU	3.4
1	C	84	ASP	3.4
1	A	203	ALA	3.4
1	A	197	CYS	3.2
1	C	60	ASN	3.2
1	A	84	ASP	3.2
1	A	200	ASP	3.1
1	B	14	HIS	3.1
1	D	224	LEU	3.1
1	A	312	THR	3.0
1	A	224	LEU	2.9
1	D	226	THR	2.8
1	A	199	PRO	2.8
1	A	208	ALA	2.8
1	A	307	ARG	2.8
1	B	86	GLY	2.7
1	A	81	ILE	2.7
1	D	236	GLY	2.7
1	A	205	MET	2.6
1	A	222	PHE	2.6
1	A	221	ASN	2.6
1	A	225	LYS	2.6
1	B	25	LYS	2.6
1	A	80	ASN	2.6
1	C	200	ASP	2.5
1	A	201	VAL	2.5
1	C	221	ASN	2.5
1	B	231	TYR	2.4
1	A	116	ASP	2.4
1	A	309	TYR	2.4
1	D	101	SER	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	230	ASN	2.3
1	A	210	GLY	2.3
1	C	310	ILE	2.3
1	C	223	THR	2.3
1	A	83	GLY	2.3
1	C	219	ILE	2.2
1	A	219	ILE	2.2
1	A	223	THR	2.2
1	A	86	GLY	2.1
1	C	14	HIS	2.1
1	D	60[A]	ASN	2.1
1	D	238	VAL	2.1
1	D	116	ASP	2.1
1	A	3	LYS	2.1
1	B	228	PHE	2.1
1	D	225	LYS	2.0
1	C	77	VAL	2.0
1	B	234	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	DNB	C	502	17/17	0.73	0.19	48,60,64,64	0
3	DNB	B	502	17/17	0.77	0.17	18,42,49,49	0
3	DNB	D	502	17/17	0.84	0.13	19,35,43,44	0
3	DNB	A	502	17/17	0.90	0.11	35,41,49,49	0
2	CA	A	501	1/1	0.97	0.04	24,24,24,24	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CA	C	501	1/1	0.97	0.04	27,27,27,27	0
2	CA	B	501	1/1	0.99	0.03	13,13,13,13	0
2	CA	D	501	1/1	0.99	0.03	16,16,16,16	0

6.5 Other polymers [i](#)

There are no such residues in this entry.