



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 11:14 AM UTC

PDB ID : 4MK7 / pdb_00004mk7
Title : Hepatitis C Virus polymerase NS5B genotype 1b (BK) in complex with inhibitor 2 (3-(3-tert-butyl-4-methoxyphenyl)pyridin-2(1H)-one)
Authors : Harris, S.F.; Wong, A.
Deposited on : 2013-09-04
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	570	
1	B	570	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9024 atoms, of which 14 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA-DIRECTED RNA POLYMERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	557	Total	C	N	O	S	0	3	0
			4362	2752	771	806	33			
1	B	559	Total	C	N	O	S	0	4	0
			4376	2759	774	808	35			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	HIS	-	expression tag	UNP P26663
B	1	HIS	-	expression tag	UNP P26663

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	O	S	0	0
			10	2	6	1	1		
3	B	1	Total	C	H	O	S	0	0
			10	2	6	1	1		

- Molecule 4 is 3-(3-tert-butyl-4-methoxyphenyl)pyridin-2(1H)-one (CCD ID: 28O) (formula: $C_{16}H_{19}NO_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	H	N	O	0	0
			20	16	1	1	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	H	N	O	0	0
			20	16	1	1	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	111	Total	O	0	0
			111	111		
5	B	114	Total	O	0	0
			114	114		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RNA-DIRECTED RNA POLYMERASE



• Molecule 1: RNA-DIRECTED RNA POLYMERASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	89.90Å 112.42Å 125.93Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.00 – 2.80 38.00 – 2.80	Depositor EDS
% Data completeness (in resolution range)	97.2 (38.00-2.80) 97.3 (38.00-2.80)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.01 (at 2.81Å)	Xtriage
Refinement program	BUSTER-TNT, BUSTER 2.11.4	Depositor
R, R_{free}	0.220 , 0.272 (Not available) , 0.240	Depositor DCC
R_{free} test set	1581 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	35.3	Xtriage
Anisotropy	0.232	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 60.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	9024	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 45.02 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.4040e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DMS, 28O, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	0/4457	1.35	8/6046 (0.1%)
1	B	0.82	1/4471 (0.0%)	1.36	11/6064 (0.2%)
All	All	0.82	1/8928 (0.0%)	1.35	19/12110 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	93	PRO	CA-C	5.26	1.54	1.51

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	129	ASP	CA-CB-CG	6.11	118.70	112.60
1	A	552	VAL	N-CA-C	-6.10	106.96	111.90
1	B	552	VAL	N-CA-C	-6.08	106.98	111.90
1	A	129	ASP	CA-CB-CG	5.86	118.46	112.60
1	A	107	ASP	CA-C-N	5.41	127.38	120.56
1	A	107	ASP	C-N-CA	5.41	127.38	120.56
1	B	107	ASP	CA-C-N	5.41	127.37	120.56
1	B	107	ASP	C-N-CA	5.41	127.37	120.56
1	A	334	ALA	CA-C-N	5.30	127.38	120.28
1	A	334	ALA	C-N-CA	5.30	127.38	120.28
1	A	490	ARG	CA-C-N	5.29	127.32	120.44
1	A	490	ARG	C-N-CA	5.29	127.32	120.44
1	B	490	ARG	CA-C-N	5.24	127.25	120.44
1	B	490	ARG	C-N-CA	5.24	127.25	120.44
1	B	351	GLY	N-CA-C	-5.10	106.32	112.49
1	B	164	ASP	CA-C-N	5.01	127.69	120.38
1	B	164	ASP	C-N-CA	5.01	127.69	120.38
1	B	315	VAL	CA-C-N	5.01	128.34	121.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	315	VAL	C-N-CA	5.01	128.34	121.33

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4362	0	4381	14	0
1	B	4376	0	4391	17	0
2	A	1	0	0	0	0
3	A	4	6	6	0	0
3	B	4	6	6	0	0
4	A	19	1	19	0	0
4	B	19	1	19	0	0
5	A	111	0	0	1	0
5	B	114	0	0	5	0
All	All	9010	14	8822	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:262:ILE:HG23	5:B:800:HOH:O	1.67	0.94
1:B:192:GLY:HA3	1:B:316:ASN:HD22	1.54	0.72
1:A:527:ASN:HD21	1:A:534:LEU:H	1.49	0.60
1:B:257:THR:CG2	5:B:800:HOH:O	2.50	0.58
1:A:86:GLU:HA	1:A:111:LEU:HD21	1.85	0.58
1:B:86:GLU:HA	1:B:111:LEU:HD21	1.85	0.57
1:A:450:ALA:HB1	5:A:792:HOH:O	2.06	0.55
1:B:200:ARG:HD2	5:B:732:HOH:O	2.07	0.54
1:A:215:MET:HB2	1:A:326:SER:HB2	1.90	0.54
1:B:192:GLY:HA3	1:B:316:ASN:ND2	2.20	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:340:THR:HG23	1:B:350:PRO:HG3	1.89	0.53
1:A:187:MET:HE1	1:A:292:THR:HG22	1.91	0.53
1:B:187:MET:HE1	1:B:292:THR:HG22	1.92	0.52
1:B:257:THR:HG22	5:B:800:HOH:O	2.09	0.50
1:B:200:ARG:HH12	1:B:316:ASN:HD21	1.58	0.49
1:A:309:GLN:O	1:A:324:CYS:HB2	2.13	0.49
1:B:309:GLN:O	1:B:324:CYS:HB2	2.13	0.49
1:B:535:LYS:H	1:B:535:LYS:HD2	1.79	0.47
1:A:82:LEU:HD13	1:A:249:ALA:HB2	1.97	0.46
1:A:230:GLU:HG3	1:A:262:ILE:HD12	1.99	0.45
1:B:144:VAL:HB	1:B:394:ARG:HG2	1.98	0.45
1:A:340:THR:HG23	1:A:350:PRO:HG3	1.99	0.44
1:B:257:THR:HG23	5:B:800:HOH:O	2.12	0.44
1:A:144:VAL:HB	1:A:394:ARG:HG2	1.99	0.44
1:A:308:LEU:HB3	1:A:311:CYS:SG	2.58	0.44
1:B:308:LEU:HB3	1:B:311:CYS:SG	2.60	0.42
1:A:555:TYR:CD1	1:A:560:ILE:HG13	2.56	0.41
1:B:555:TYR:CD1	1:B:560:ILE:HG13	2.56	0.41
1:A:419:LEU:HD23	1:A:485:VAL:HG21	2.03	0.40
1:A:138:ILE:HD11	1:A:159:LEU:HG	2.03	0.40
1:B:419:LEU:HD23	1:B:485:VAL:HG21	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	556/570 (98%)	543 (98%)	13 (2%)	0	100	100
1	B	559/570 (98%)	542 (97%)	17 (3%)	0	100	100
All	All	1115/1140 (98%)	1085 (97%)	30 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	478/485 (99%)	459 (96%)	19 (4%)	28	63
1	B	479/485 (99%)	458 (96%)	21 (4%)	25	59
All	All	957/970 (99%)	917 (96%)	40 (4%)	26	61

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	CYS
1	A	31	LEU
1	A	47	LEU
1	A	57	LEU
1	A	83	LEU
1	A	159	LEU
1	A	213	ASN
1	A	220	ASP
1	A	222	ARG
1	A	237	GLU
1	A	278	ARG
1	A	308	LEU
1	A	315	VAL
1	A	316	ASN
1	A	320	LEU
1	A	363	ILE
1	A	379	LYS
1	A	419	LEU
1	A	466	LEU
1	B	14	CYS
1	B	31	LEU
1	B	57	LEU
1	B	83	LEU
1	B	100	LYS
1	B	159	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	213	ASN
1	B	222	ARG
1	B	237	GLU
1	B	278	ARG
1	B	308	LEU
1	B	315	VAL
1	B	316	ASN
1	B	320	LEU
1	B	363	ILE
1	B	419	LEU
1	B	461	GLN
1	B	466	LEU
1	B	534	LEU
1	B	535	LYS
1	B	545	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (21) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	ASN
1	A	35	ASN
1	A	63	HIS
1	A	95	HIS
1	A	120	HIS
1	A	184	GLN
1	A	273	ASN
1	A	411	ASN
1	A	438	GLN
1	A	527	ASN
1	B	24	ASN
1	B	35	ASN
1	B	63	HIS
1	B	95	HIS
1	B	120	HIS
1	B	148	GLN
1	B	184	GLN
1	B	316	ASN
1	B	411	ASN
1	B	438	GLN
1	B	483	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	28O	A	603	-	20,20,20	0.37	0	29,29,29	0.59	0
3	DMS	A	602	-	3,3,3	0.31	0	3,3,3	0.25	0
3	DMS	B	601	-	3,3,3	0.41	0	3,3,3	0.11	0
4	28O	B	602	-	20,20,20	0.44	0	29,29,29	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	28O	A	603	-	-	0/12/12/12	0/2/2/2
4	28O	B	602	-	-	0/12/12/12	0/2/2/2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	557/570 (97%)	0.17	9 (1%) 70 61	13, 34, 57, 76	3 (0%)
1	B	559/570 (98%)	0.08	6 (1%) 78 70	14, 31, 52, 91	4 (0%)
All	All	1116/1140 (97%)	0.13	15 (1%) 75 66	13, 33, 54, 91	7 (0%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	148	GLN	3.2
1	B	379	LYS	3.0
1	A	148	GLN	2.9
1	B	152	GLY	2.8
1	A	379	LYS	2.4
1	B	531	LYS	2.3
1	A	16	ALA	2.2
1	A	536	LEU	2.2
1	A	15	ALA	2.2
1	A	352	ASP	2.1
1	B	514	GLN	2.1
1	A	544	ARG	2.1
1	A	102	GLY	2.0
1	B	524	TYR	2.0
1	A	270	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	DMS	B	601	4/4	0.89	0.15	42,44,46,46	0
3	DMS	A	602	4/4	0.93	0.11	49,49,51,51	0
2	MG	A	601	1/1	0.93	0.11	37,37,37,37	0
4	28O	B	602	19/19	0.93	0.09	20,25,29,30	0
4	28O	A	603	19/19	0.94	0.08	32,35,37,37	0

6.5 Other polymers

There are no such residues in this entry.