



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 05:41 PM UTC

PDB ID : 5MKG / pdb_00005mkg
Title : PA3825-EAL Ca-CdG Structure
Authors : Horrell, S.; Bellini, D.; Strange, R.; Wagner, A.; Walsh, M.
Deposited on : 2016-12-04
Resolution : 2.44 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2340 (2.46-2.42)
Clashscore	190562	2400 (2.46-2.42)
Ramachandran outliers	187476	2379 (2.46-2.42)
Sidechain outliers	187428	2379 (2.46-2.42)
RSRZ outliers	180081	2340 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	256	29% 90% 6% ..
2	B	254	21% 87% 9% ..

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4066 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Diguanylate phosphodiesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	248	Total	C	N	O	S	0	0	0
			1940	1233	336	364	7			


- Molecule 2 is a protein called Diguanylate phosphodiesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	249	Total	C	N	O	S	0	0	0
			1951	1243	336	365	7			

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Ca	0	0
			3	3		
3	B	4	Total	Ca	0	0
			4	4		

- Molecule 4 is 9,9'-[(2R,3R,3aS,5S,7aR,9R,10R,10aS,12S,14aR)-3,5,10,12-tetrahydroxy-5,12-dioxidoctahydro-2H,7H-difuro[3,2-d:3',2'-j][1,3,7,9,2,8]tetraoxadiphosphacyclododecine-2,9-diyl]bis(2-amino-1,9-dihydro-6H-purin-6-one) (CCD ID: C2E) (formula: C₂₀H₂₄N₁₀O₁₄P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 46	C 20	N 10	O 14	P 2	0	0
4	B	1	Total 46	C 20	N 10	O 14	P 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	35	Total	O	0	0
			35	35		
5	B	41	Total	O	0	0
			41	41		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Diguanylate phosphodiesterase



- Molecule 2: Diguanylate phosphodiesterase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	112.23Å 59.40Å 92.78Å 90.00° 115.03° 90.00°	Depositor
Resolution (Å)	51.29 – 2.44 51.29 – 2.44	Depositor EDS
% Data completeness (in resolution range)	98.0 (51.29-2.44) 98.0 (51.29-2.44)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.44 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.238 , 0.285 0.239 , 0.285	Depositor DCC
R_{free} test set	1060 reflections (5.19%)	wwPDB-VP
Wilson B-factor (Å ²)	55.1	Xtriage
Anisotropy	0.073	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 53.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4066	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 29.29 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.5893e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, C2E

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.61	0/1979	0.90	4/2682 (0.1%)
2	B	0.61	0/1991	0.90	2/2699 (0.1%)
All	All	0.61	0/3970	0.90	6/5381 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	175	PHE	CA-C-N	5.95	132.41	121.70
1	A	175	PHE	C-N-CA	5.95	132.41	121.70
2	B	53	ARG	CA-C-N	5.70	125.82	119.32
2	B	53	ARG	C-N-CA	5.70	125.82	119.32
1	A	53	ARG	CA-C-N	5.53	125.62	119.32
1	A	53	ARG	C-N-CA	5.53	125.62	119.32

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1940	0	1935	8	0
2	B	1951	0	1947	14	0
3	A	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	4	0	0	0	0
4	A	46	0	22	0	0
4	B	46	0	22	0	0
5	A	35	0	0	0	0
5	B	41	0	0	1	0
All	All	4066	0	3926	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:72:ARG:HH22	2:B:108:ALA:HB3	1.54	0.70
2:B:104:CYS:O	2:B:143:THR:HG22	1.96	0.65
1:A:36:ILE:HD11	1:A:235:TYR:CD1	2.35	0.61
2:B:36:ILE:HD11	2:B:235:TYR:CD1	2.39	0.57
2:B:159:ILE:HD11	2:B:177:VAL:HG11	1.88	0.55
1:A:116:LEU:HD22	1:A:127:LEU:HD13	1.94	0.51
1:A:90:GLU:HG3	1:A:91:PRO:HD2	1.93	0.50
2:B:172:LEU:HD13	2:B:210:LEU:HD13	1.94	0.50
1:A:175:PHE:HA	1:A:176:GLN:HB2	1.95	0.48
1:A:186:PHE:HB3	1:A:200:LEU:HD12	1.95	0.48
1:A:36:ILE:CD1	1:A:235:TYR:CE1	2.97	0.47
2:B:36:ILE:CD1	2:B:235:TYR:CE1	2.98	0.47
2:B:171:TYR:O	2:B:173:ARG:N	2.48	0.47
1:A:184:GLN:HG2	1:A:218:GLY:HA3	1.98	0.44
2:B:145:ARG:NH1	5:B:402:HOH:O	2.50	0.44
2:B:170:ALA:O	2:B:172:LEU:N	2.48	0.44
2:B:13:ARG:HG3	2:B:14:ALA:N	2.33	0.43
2:B:139:SER:OG	2:B:143:THR:OG1	2.37	0.43
2:B:116:LEU:HD22	2:B:127:LEU:HD13	2.01	0.42
2:B:36:ILE:CD1	2:B:235:TYR:CD1	3.03	0.42
1:A:36:ILE:CD1	1:A:235:TYR:CD1	3.01	0.41
2:B:15:LEU:HD21	2:B:69:PRO:HB2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	244/256 (95%)	233 (96%)	7 (3%)	4 (2%)	7	6
2	B	245/254 (96%)	233 (95%)	8 (3%)	4 (2%)	7	6
All	All	489/510 (96%)	466 (95%)	15 (3%)	8 (2%)	7	6

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	141	GLU
1	A	142	VAL
2	B	109	LEU
2	B	171	TYR
2	B	172	LEU
2	B	18	ASN
1	A	176	GLN
1	A	139	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/215 (97%)	206 (99%)	3 (1%)	59	70
2	B	210/214 (98%)	206 (98%)	4 (2%)	50	63
All	All	419/429 (98%)	412 (98%)	7 (2%)	53	66

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	184	GLN
1	A	248	GLU
1	A	253	SER
2	B	8	SER
2	B	13	ARG
2	B	107	LEU
2	B	148	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	184	GLN
1	A	260	GLN
2	B	184	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 7 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	C2E	B	304	3	52,52,52	1.22	5 (9%)	78,82,82	1.88	14 (17%)
4	C2E	A	304	3	52,52,52	1.18	6 (11%)	78,82,82	1.80	14 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	C2E	B	304	3	-	7/30/62/62	0/6/7/7
4	C2E	A	304	3	-	6/30/62/62	0/6/7/7

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	304	C2E	C51-C41	3.45	1.48	1.38
4	B	304	C2E	C5-C4	3.22	1.47	1.38
4	A	304	C2E	C5-C4	3.03	1.47	1.38
4	A	304	C2E	C51-C41	3.01	1.47	1.38
4	B	304	C2E	C51-N71	-2.76	1.33	1.39
4	A	304	C2E	C5-N7	-2.74	1.33	1.39
4	B	304	C2E	C61-N11	-2.53	1.34	1.38
4	A	304	C2E	C6-N1	-2.44	1.34	1.38
4	A	304	C2E	C4-N9	-2.28	1.32	1.38
4	B	304	C2E	C6-N1	-2.26	1.34	1.38
4	A	304	C2E	C61-N11	-2.08	1.35	1.38

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	304	C2E	C51-C41-N31	-6.21	118.51	128.39
4	B	304	C2E	C5-C4-N3	-5.66	119.38	128.39
4	A	304	C2E	C5-C4-N3	-5.48	119.67	128.39
4	A	304	C2E	C51-C41-N31	-5.34	119.90	128.39
4	B	304	C2E	C2-N3-C4	5.16	121.19	112.30
4	A	304	C2E	C21-N31-C41	4.79	120.55	112.30
4	B	304	C2E	N91-C41-N31	4.73	135.41	125.95
4	B	304	C2E	C21-N31-C41	4.42	119.91	112.30
4	A	304	C2E	C2-N3-C4	4.33	119.76	112.30
4	A	304	C2E	N9-C4-N3	4.17	134.30	125.95
4	B	304	C2E	C6-C5-N7	4.07	137.70	130.29

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	304	C2E	N9-C4-N3	4.01	133.97	125.95
4	A	304	C2E	N91-C41-N31	3.80	133.56	125.95
4	A	304	C2E	C61-C51-N71	3.71	137.04	130.29
4	A	304	C2E	C6-C5-N7	2.91	135.58	130.29
4	B	304	C2E	C4-C5-N7	-2.73	106.34	110.67
4	A	304	C2E	C41-C51-N71	-2.63	106.50	110.67
4	B	304	C2E	O3'-P11-O11	-2.58	101.55	109.81
4	A	304	C2E	C2'-C3'-C4'	2.52	107.65	103.24
4	A	304	C2E	C4-C5-N7	-2.44	106.80	110.67
4	A	304	C2E	O6-C6-C5	-2.42	120.15	126.53
4	B	304	C2E	C61-C51-N71	2.34	134.55	130.29
4	B	304	C2E	O61-C61-C51	-2.30	120.47	126.53
4	B	304	C2E	C41-C51-N71	-2.27	107.07	110.67
4	B	304	C2E	O2P-P1-O1P	2.27	122.99	112.44
4	A	304	C2E	O21-P11-O11	2.24	122.85	112.44
4	B	304	C2E	C5-C6-N1	2.02	118.40	113.25
4	A	304	C2E	O61-C61-C51	-2.01	121.24	126.53

There are no chirality outliers.

All (13) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	304	C2E	C3A-O3A-P1-O2P
4	B	304	C2E	C3A-C4A-C5A-O5A
4	B	304	C2E	O4A-C4A-C5A-O5A
4	A	304	C2E	C3A-O3A-P1-O1P
4	B	304	C2E	C3'-O3'-P11-O21
4	B	304	C2E	C4A-C5A-O5A-P11
4	A	304	C2E	C3'-C4'-C5'-O5'
4	A	304	C2E	C4'-C5'-O5'-P1
4	A	304	C2E	C2A-C3A-O3A-P1
4	B	304	C2E	C4'-C3'-O3'-P11
4	B	304	C2E	C2'-C3'-O3'-P11
4	B	304	C2E	C3'-O3'-P11-O11
4	A	304	C2E	C4A-C3A-O3A-P1

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	248/256 (96%)	1.42	73 (29%)  	27, 57, 83, 114	58 (23%)
2	B	249/254 (98%)	1.15	54 (21%)  	25, 51, 86, 107	56 (22%)
All	All	497/510 (97%)	1.28	127 (25%)  	25, 55, 85, 114	114 (22%)

All (127) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	137	ILE	11.6
1	A	142	VAL	8.2
1	A	143	THR	7.7
1	A	144	ASP	7.6
2	B	175	PHE	6.9
1	A	174	LYS	6.9
1	A	139	SER	6.3
1	A	140	SER	6.3
2	B	172	LEU	5.9
1	A	138	GLU	5.7
1	A	176	GLN	5.6
2	B	173	ARG	5.4
2	B	45	ARG	5.2
2	B	105	HIS	5.1
1	A	75	MET	5.1
2	B	255	THR	4.8
1	A	255	THR	4.7
1	A	13	ARG	4.6
2	B	104	CYS	4.6
2	B	12	ARG	4.5
1	A	134	ARG	4.5
1	A	250	LEU	4.4
2	B	256	VAL	4.3
2	B	254	LEU	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	137	ILE	4.1
2	B	145	ARG	4.0
2	B	174	LYS	4.0
2	B	52	ILE	4.0
1	A	47	PRO	4.0
2	B	46	HIS	3.9
2	B	63	ARG	3.7
2	B	170	ALA	3.7
1	A	10	GLU	3.7
1	A	107	LEU	3.7
1	A	259	GLY	3.7
2	B	136	LEU	3.7
2	B	109	LEU	3.5
2	B	107	LEU	3.5
1	A	254	LEU	3.4
1	A	209	LYS	3.3
1	A	141	GLU	3.3
1	A	153	LEU	3.3
1	A	196	SER	3.3
2	B	103	HIS	3.3
1	A	247	LEU	3.2
2	B	116	LEU	3.2
1	A	116	LEU	3.2
1	A	9	SER	3.2
1	A	135	GLU	3.2
1	A	88	LYS	3.1
1	A	16	GLU	3.1
2	B	135	GLU	3.1
2	B	196	SER	3.1
1	A	15	LEU	3.1
1	A	214	ILE	3.1
1	A	72	ARG	3.0
1	A	191	GLY	3.0
2	B	171	TYR	3.0
2	B	176	GLN	3.0
1	A	252	SER	3.0
2	B	133	GLU	2.9
1	A	60	PHE	2.9
1	A	97	PHE	2.9
2	B	194	THR	2.9
1	A	127	LEU	2.9
1	A	104	CYS	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	21	ILE	2.8
2	B	251	LEU	2.8
1	A	71	THR	2.7
2	B	138	GLU	2.7
1	A	68	VAL	2.7
1	A	147	PHE	2.7
1	A	177	VAL	2.7
1	A	76	ARG	2.6
1	A	22	PRO	2.6
1	A	227	TYR	2.6
1	A	211	ASP	2.6
2	B	111	ASP	2.6
1	A	50	GLY	2.6
1	A	12	ARG	2.5
1	A	44	TRP	2.5
2	B	163	GLY	2.5
2	B	16	GLU	2.5
2	B	258	GLU	2.5
2	B	66	LEU	2.5
1	A	103	HIS	2.5
1	A	114	ARG	2.5
1	A	251	LEU	2.5
2	B	71	THR	2.5
2	B	72	ARG	2.5
1	A	124	HIS	2.5
2	B	44	TRP	2.5
1	A	45	ARG	2.5
2	B	252	SER	2.5
2	B	112	ASP	2.4
1	A	18	ASN	2.4
1	A	53	ARG	2.4
1	A	175	PHE	2.4
2	B	50	GLY	2.3
1	A	192	ILE	2.3
2	B	147	PHE	2.3
2	B	149	GLU	2.3
1	A	205	GLU	2.3
1	A	159	ILE	2.3
2	B	211	ASP	2.3
1	A	106	GLU	2.3
1	A	96	GLY	2.2
2	B	257	GLN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	248	GLU	2.2
2	B	18	ASN	2.2
1	A	129	LEU	2.2
1	A	136	LEU	2.2
2	B	64	SER	2.2
1	A	148	ASP	2.2
2	B	124	HIS	2.2
2	B	88	LYS	2.2
1	A	122	PRO	2.1
1	A	5	ALA	2.1
1	A	14	ALA	2.1
1	A	74	LEU	2.1
2	B	15	LEU	2.1
1	A	99	ILE	2.1
2	B	248	GLU	2.1
2	B	139	SER	2.1
1	A	223	GLU	2.1
1	A	186	PHE	2.0
2	B	51	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CA	A	302	1/1	0.79	0.12	98,98,98,98	0
3	CA	B	303	1/1	0.84	0.11	102,102,102,102	0
3	CA	A	301	1/1	0.91	0.16	98,98,98,98	0
3	CA	B	301	1/1	0.92	0.10	95,95,95,95	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	C2E	A	304	46/46	0.94	0.08	41,46,49,52	0
4	C2E	B	304	46/46	0.95	0.07	38,41,43,49	0
3	CA	B	305	1/1	0.96	0.09	56,56,56,56	0
3	CA	B	302	1/1	0.99	0.02	38,38,38,38	0
3	CA	A	303	1/1	1.00	0.03	51,51,51,51	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around C2E A 304:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around C2E B 304:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.