



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 2, 2026 – 02:27 AM UTC

PDB ID : 6MKD / pdb\_00006mkd  
Title : 4699 TCR bound to I-Ab Padi4  
Authors : Stadinski, B.D.; Blevins, S.J.; Huseby, E.S.  
Deposited on : 2018-09-25  
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	179	 3% (red), 61% (green), 30% (yellow), 7% (grey)
2	D	217	 60% (green), 32% (yellow), 7% (grey)
3	A	208	 2% (red), 66% (green), 26% (yellow), 7% (grey)
4	B	239	 70% (green), 29% (yellow), 1% (grey)

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5916 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H-2 class II histocompatibility antigen, A-B alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	C	166	1211	783	198	227	3	0	0	0

- Molecule 2 is a protein called Padi4 (92-105) peptide and MHC Class II I-Ab beta chain,H-2 class II histocompatibility antigen, A beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	201	1584	1001	276	300	7	0	0	0

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-12	GLY	-	linker	UNP Q9Z183
D	-11	GLY	-	linker	UNP Q9Z183
D	-10	GLY	-	linker	UNP Q9Z183
D	-9	GLY	-	linker	UNP Q9Z183
D	-8	SER	-	linker	UNP Q9Z183
D	-7	LEU	-	linker	UNP Q9Z183
D	-6	VAL	-	linker	UNP Q9Z183
D	-5	PRO	-	linker	UNP Q9Z183
D	-4	ARG	-	linker	UNP Q9Z183
D	-3	GLY	-	linker	UNP Q9Z183
D	-2	SER	-	linker	UNP Q9Z183
D	-1	GLY	-	linker	UNP Q9Z183
D	0	GLY	-	linker	UNP Q9Z183
D	1	GLY	-	linker	UNP Q9Z183
D	2	GLY	-	linker	UNP Q9Z183

- Molecule 3 is a protein called 4699 TCR alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	A	193	1339	853	222	258	6	0	0	0


- Molecule 4 is a protein called 4699 TCR beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	B	239	1782	1125	316	335	6	0	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: H-2 class II histocompatibility antigen, A-B alpha chain



- Molecule 2: Padi4 (92-105) peptide and MHC Class II I-Ab beta chain, H-2 class II histocompatibility antigen, A beta chain




- Molecule 3: 4699 TCR alpha chain





● Molecule 4: 4699 TCR beta chain

Chain B: 70% 29%



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	255.45Å 73.64Å 64.57Å 90.00° 90.39° 90.00°	Depositor
Resolution (Å)	29.24 – 3.20 29.24 – 3.20	Depositor EDS
% Data completeness (in resolution range)	97.8 (29.24-3.20) 97.7 (29.24-3.20)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.59 (at 3.18Å)	Xtrriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, $R_{free}$	0.233 , 0.278 0.234 , 0.280	Depositor DCC
$R_{free}$ test set	1172 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	70.3	Xtrriage
Anisotropy	0.625	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 53.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.037 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	5916	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.92% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	C	0.43	0/1242	0.68	0/1701
2	D	0.45	0/1627	0.70	0/2226
3	A	0.40	0/1372	0.73	5/1884 (0.3%)
4	B	0.38	0/1835	0.60	0/2519
All	All	0.42	0/6076	0.68	5/8330 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	152	LYS	N-CA-C	-8.35	101.42	111.69
3	A	170	ASP	CA-C-N	6.90	134.12	121.70
3	A	170	ASP	C-N-CA	6.90	134.12	121.70
3	A	153	ASP	CA-C-N	5.47	130.81	121.29
3	A	153	ASP	C-N-CA	5.47	130.81	121.29

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1211	0	1062	61	0
2	D	1584	0	1446	67	0
3	A	1339	0	1111	35	0
4	B	1782	0	1584	55	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5916	0	5203	196	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 196 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:23:GLN:HG2	4:B:30:MET:HE3	1.41	1.02
1:C:13:TYR:CE2	1:C:18:ASP:HA	2.13	0.84
4:B:156:TRP:HB2	4:B:205:ARG:HB3	1.59	0.84
3:A:13:TRP:CZ3	3:A:144:SER:OG	2.31	0.83
1:C:13:TYR:OH	1:C:18:ASP:CB	2.29	0.80

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	154/179 (86%)	142 (92%)	12 (8%)	0	100	100
2	D	197/217 (91%)	174 (88%)	23 (12%)	0	100	100
3	A	185/208 (89%)	167 (90%)	17 (9%)	1 (0%)	24	59
4	B	237/239 (99%)	219 (92%)	18 (8%)	0	100	100
All	All	773/843 (92%)	702 (91%)	70 (9%)	1 (0%)	48	79

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	119	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	117/160 (73%)	113 (97%)	4 (3%)	32	64
2	D	166/192 (86%)	165 (99%)	1 (1%)	78	84
3	A	118/185 (64%)	118 (100%)	0	100	100
4	B	173/204 (85%)	170 (98%)	3 (2%)	53	75
All	All	574/741 (78%)	566 (99%)	8 (1%)	59	77

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	B	72	PHE
4	B	62	TYR
2	D	139	GLU
1	C	125	SER
4	B	4	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
4	B	22	ASN
4	B	69	GLN
4	B	203	HIS
4	B	71	ASN
1	C	111	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	C	166/179 (92%)	0.27	5 (3%) 52 33	43, 66, 99, 109	0
2	D	201/217 (92%)	-0.12	0 100 100	36, 53, 68, 80	0
3	A	193/208 (92%)	0.32	4 (2%) 63 43	28, 74, 104, 117	0
4	B	239/239 (100%)	-0.00	0 100 100	46, 67, 83, 95	0
All	All	799/843 (94%)	0.10	9 (1%) 78 61	28, 64, 96, 117	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	118	ASN	3.8
1	C	99	LEU	2.9
1	C	177	HIS	2.8
3	A	143	ASP	2.4
3	A	170	ASP	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.