



Full wwPDB NMR Structure Validation Report ⓘ

Mar 10, 2026 – 12:18 AM UTC

PDB ID : 2MLG / pdb_00002mlg
BMRB ID : 19821
Title : Stf76 from the Sulfolobus islandicus plasmid-virus pSSVx
Authors : Farina, B.; Russo, L.; Fattorusso, R.
Deposited on : 2014-02-27

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 49%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	84	

2 Ensemble composition and analysis

This entry contains 20 models. The atoms present in the NMR models are not consistent. Some calculations may have failed as a result. All residues are included in the validation scores. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:6-A:68 (63)	0.89	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 4, 5, 6, 13, 14, 16, 17, 18
2	8, 9, 11, 12, 15, 20
3	3, 7, 10, 19

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 1340 atoms, of which 683 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Sulfolobus transcription factor 76 aminoacid protein, Stf76.


Mol	Chain	Residues	Atoms						Trace
1	A	77	Total	C	H	N	O	S	0
			1340	415	683	127	113	2	

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Sulfolobus transcription factor 76 aminoacid protein, Stf76




4.2 Scores per residue for each member of the ensemble

Colouring as in section [4.1](#) above.


4.2.1 Score per residue for model 1 (medoid)

- Molecule 1: Sulfolobus transcription factor 76 aminoacid protein, Stf76



4.2.2 Score per residue for model 2


- Molecule 1: Sulfolobus transcription factor 76 aminoacid protein, Stf76



4.2.3 Score per residue for model 3

- Molecule 1: Sulfolobus transcription factor 76 aminoacid protein, Stf76


Chain A:  73% 17% 8%



4.2.4 Score per residue for model 4

- Molecule 1: Sulfolobus transcription factor 76 aminoacid protein, Stf76


Chain A:  74% 17% 8%



4.2.5 Score per residue for model 5

- Molecule 1: Sulfolobus transcription factor 76 aminoacid protein, Stf76


Chain A:  74% 17% 8%



4.2.6 Score per residue for model 6

- Molecule 1: Sulfolobus transcription factor 76 aminoacid protein, Stf76


Chain A:  75% 17% 8%



4.2.7 Score per residue for model 7

- Molecule 1: Sulfolobus transcription factor 76 aminoacid protein, Stf76


Chain A:  75% 17% 8%



4.2.8 Score per residue for model 8

- Molecule 1: Sulfolobus transcription factor 76 aminoacid protein, Stf76


Chain A:  74% 17% 8%



4.2.9 Score per residue for model 9

- Molecule 1: Sulfolobus transcription factor 76 aminoacid protein, Stf76


Chain A:  73% 17% 8%



4.2.10 Score per residue for model 10

- Molecule 1: Sulfolobus transcription factor 76 aminoacid protein, Stf76


Chain A:  73% 17% 8%



4.2.11 Score per residue for model 11

- Molecule 1: Sulfolobus transcription factor 76 aminoacid protein, Stf76


Chain A:  75% 17% 8%



4.2.12 Score per residue for model 12

- Molecule 1: Sulfolobus transcription factor 76 aminoacid protein, Stf76


Chain A:  75% 17% 8%



4.2.13 Score per residue for model 13

- Molecule 1: Sulfolobus transcription factor 76 aminoacid protein, Stf76


Chain A:  74% 17% 8%



4.2.14 Score per residue for model 14

- Molecule 1: Sulfolobus transcription factor 76 aminoacid protein, Stf76


Chain A:  75% 17% 8%



4.2.15 Score per residue for model 15

- Molecule 1: Sulfolobus transcription factor 76 aminoacid protein, Stf76


Chain A:  75% 17% 8%



4.2.16 Score per residue for model 16

- Molecule 1: Sulfolobus transcription factor 76 aminoacid protein, Stf76


Chain A:  75% 17% 8%



4.2.17 Score per residue for model 17

- Molecule 1: Sulfolobus transcription factor 76 aminoacid protein, Stf76


Chain A:  74% 17% 8%



4.2.18 Score per residue for model 18

- Molecule 1: Sulfolobus transcription factor 76 aminoacid protein, Stf76


Chain A:  75% 17% 8%



4.2.19 Score per residue for model 19

- Molecule 1: Sulfolobus transcription factor 76 aminoacid protein, Stf76


Chain A:  74% 17% 8%



4.2.20 Score per residue for model 20

- Molecule 1: Sulfolobus transcription factor 76 aminoacid protein, Stf76

Chain A:  75% 17% 8%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *MFR*.

Of the 3000 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CS-ROSETTA	structure solution	
CS-ROSETTA	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	566
Number of shifts mapped to atoms	566
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	49%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.53±0.00	0±0/548 (0.0± 0.0%)	0.70±0.01	0±0/736 (0.0± 0.1%)
All	All	0.53	0/10960 (0.0%)	0.70	7/14720 (0.0%)

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	21	CYS	N-CA-CB	-5.68	102.75	111.56	4	2
1	A	67	CYS	N-CA-CB	-5.67	102.78	111.56	3	5

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	539	565	565	0±1
All	All	10780	11300	11300	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:39:LEU:O	1:A:39:LEU:HD23	0.60	1.96	13	1
1:A:13:LEU:C	1:A:13:LEU:HD23	0.52	2.30	10	1
1:A:13:LEU:HD23	1:A:13:LEU:O	0.50	2.07	10	1
1:A:39:LEU:HD23	1:A:39:LEU:C	0.46	2.35	13	1
1:A:13:LEU:O	1:A:13:LEU:HD23	0.42	2.13	5	1
1:A:16:LEU:O	1:A:20:GLY:N	0.42	2.53	9	1
1:A:13:LEU:HD23	1:A:13:LEU:C	0.42	2.40	5	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	63/84 (75%)	63±1 (100±1%)	0±1 (0±1%)	0±0 (0±0%)	100	100
All	All	1260/1680 (75%)	1254 (100%)	6 (0%)	0 (0%)	100	100

There are no Ramachandran outliers.

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	59/78 (76%)	59±0 (100±0%)	0±0 (0±0%)	87	98
All	All	1180/1560 (76%)	1179 (100%)	1 (0%)	87	98

All 1 unique residues with a non-rotameric sidechain are listed below.

Mol	Chain	Res	Type	Models (Total)
1	A	68	LEU	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 49% for the well-defined parts and 48% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	566
Number of shifts mapped to atoms	566
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	77	-0.42 ± 0.16	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	0	—	None (insufficient data)
$^{13}\text{C}'$	0	—	None (insufficient data)
^{15}N	77	-1.69 ± 0.21	Should be applied

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 49%, i.e. 478 atoms were assigned a chemical shift out of a possible 981. 0 out of 15 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	255/318 (80%)	129/129 (100%)	63/126 (50%)	63/63 (100%)
Sidechain	223/588 (38%)	223/380 (59%)	0/173 (0%)	0/35 (0%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	0/75 (0%)	0/36 (0%)	0/34 (0%)	0/5 (0%)
Overall	478/981 (49%)	352/545 (65%)	63/333 (19%)	63/103 (61%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 48%, i.e. 566 atoms were assigned a chemical shift out of a possible 1186. 0 out of 17 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	311/388 (80%)	157/157 (100%)	77/154 (50%)	77/77 (100%)
Sidechain	255/707 (36%)	255/455 (56%)	0/211 (0%)	0/41 (0%)
Aromatic	0/91 (0%)	0/44 (0%)	0/38 (0%)	0/9 (0%)
Overall	566/1186 (48%)	412/656 (63%)	77/403 (19%)	77/127 (61%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

