



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 10:16 AM UTC

PDB ID : 4MLV / pdb_00004mlv
Title : Crystal Structure of Bacillus megaterium porphobilinogen deaminase
Authors : Azim, N.; Deery, E.; Warren, M.J.; Erskine, P.; Cooper, J.B.; Coker, A.;
Wood, S.P.; Akhtar, M.
Deposited on : 2013-09-06
Resolution : 1.46 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1756 (1.46-1.46)
Clashscore	190562	1795 (1.46-1.46)
Ramachandran outliers	187476	1776 (1.46-1.46)
Sidechain outliers	187428	1776 (1.46-1.46)
RSRZ outliers	180081	1756 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	312	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2662 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Porphobilinogen deaminase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	290	2272	1416	398	446	12	0	8	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	SER	-	expression tag	UNP Q8GCA8
A	0	HIS	-	expression tag	UNP Q8GCA8


- Molecule 2 is 3-[5-{{3-(2-carboxyethyl)-4-(carboxymethyl)-5-methyl-1H-pyrrol-2-yl)methyl}-4-(carboxymethyl)-1H-pyrrol-3-yl]propanoic acid (CCD ID: DPM) (formula: C₂₀H₂₄N₂O₈).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	30	20	2	8	0	1


- Molecule 3 is 3-[(5S)-5-{{3-(2-carboxyethyl)-4-(carboxymethyl)-5-methyl-1H-pyrrol-2-yl)m

ethyl}-4-(carboxymethyl)-2-oxo-2,5-dihydro-1H-pyrrol-3-yl]propanoic acid (CCD ID: 29P)
(formula: C₂₀H₂₄N₂O₉).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
3	A	1	31	20	2	9	0	1

- Molecule 4 is ACETIC ACID (CCD ID: ACY) (formula: C₂H₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	4	2	2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	325	Total 325	O 325	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Porphobilinogen deaminase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	53.32Å 65.78Å 97.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.75 – 1.46 46.75 – 1.46	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.75-1.46) 100.0 (46.75-1.46)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.30 (at 1.46Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.141 , 0.186 0.139 , 0.184	Depositor DCC
R_{free} test set	3073 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	18.7	Xtrriage
Anisotropy	0.038	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 38.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2662	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ACY, DPM, 29P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.51	13/2352 (0.6%)	1.35	10/3162 (0.3%)

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	243	VAL	CA-CB	10.37	1.67	1.54
1	A	190	ILE	CA-CB	-9.97	1.43	1.54
1	A	174	ARG	CZ-NH2	-7.00	1.24	1.33
1	A	190	ILE	N-CA	-6.46	1.39	1.46
1	A	2	ARG	C-O	6.22	1.31	1.23
1	A	33	GLU	N-CA	-5.68	1.39	1.45
1	A	98	LEU	C-O	-5.60	1.17	1.23
1	A	206	GLU	N-CA	-5.59	1.39	1.46
1	A	181	THR	N-CA	-5.28	1.39	1.46
1	A	32	PHE	N-CA	-5.27	1.39	1.46
1	A	191	SER	CA-C	-5.24	1.47	1.53
1	A	17	GLN	CD-NE2	-5.19	1.22	1.33
1	A	17	GLN	CA-CB	5.15	1.61	1.53

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	150	ILE	CB-CA-C	-10.07	98.85	112.04
1	A	64	ILE	N-CA-C	-7.53	103.11	110.72
1	A	190	ILE	N-CA-C	6.96	121.56	113.42
1	A	10	ARG	N-CA-C	6.20	120.84	113.28
1	A	0	HIS	CA-C-O	6.01	127.33	120.84
1	A	139	SER	N-CA-C	5.63	118.35	111.82
1	A	242	GLN	CB-CG-CD	5.49	121.94	112.60
1	A	228	ARG	CB-CG-CD	5.31	123.51	111.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	231	ARG	CD-NE-CZ	-5.26	117.03	124.40
1	A	138	ARG	NE-CZ-NH2	5.05	123.75	119.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2272	0	2323	15	0
2	A	30	0	18	0	0
3	A	31	0	18	0	0
4	A	4	0	3	0	0
5	A	325	0	0	5	0
All	All	2662	0	2362	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:149:ASN:HB2	5:A:718:HOH:O	1.79	0.82
1:A:287[B]:GLU:OE1	1:A:291:ARG:NH2	2.29	0.63
1:A:122:VAL:HG11	1:A:144:LYS:HE3	1.83	0.60
1:A:122:VAL:HG11	1:A:144:LYS:CE	2.44	0.47
1:A:300:LEU:HD23	1:A:300:LEU:C	2.40	0.47
1:A:150:ILE:N	1:A:150:ILE:HD13	2.29	0.46
1:A:1:MET:HG2	5:A:724:HOH:O	2.16	0.46
1:A:17:GLN:NE2	5:A:565:HOH:O	2.48	0.45
1:A:39:MET:HE3	1:A:39:MET:HB3	1.82	0.45
1:A:10:ARG:HG3	5:A:791:HOH:O	2.16	0.44
1:A:17:GLN:HG2	1:A:174:ARG:HD3	1.98	0.44
1:A:236:GLU:HG2	5:A:814:HOH:O	2.16	0.44
1:A:12:LYS:HE3	1:A:175:MET:CE	2.48	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:262:LEU:HA	1:A:272:TYR:O	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	296/312 (95%)	293 (99%)	3 (1%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	253/261 (97%)	247 (98%)	6 (2%)	43 12

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	MET
1	A	100[A]	GLU
1	A	100[B]	GLU
1	A	144	LYS
1	A	150	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	151	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	149	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	29P	A	402[B]	1	32,32,32	2.55	14 (43%)	35,45,45	2.26	13 (37%)
4	ACY	A	403	-	3,3,3	0.72	0	3,3,3	0.71	0
2	DPM	A	401[A]	1	31,31,31	2.37	15 (48%)	39,43,43	1.72	11 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	29P	A	402[B]	1	-	7/22/38/38	0/2/2/2
2	DPM	A	401[A]	1	-	2/22/22/22	0/2/2/2

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402[B]	29P	C4B-C3B	-7.68	1.34	1.48
3	A	402[B]	29P	CHB-C4A	5.40	1.56	1.50
2	A	401[A]	DPM	C4B-C3B	-4.25	1.32	1.38
2	A	401[A]	DPM	C8A-C9A	4.12	1.60	1.50
3	A	402[B]	29P	C8A-C9A	4.09	1.60	1.50
2	A	401[A]	DPM	O4A-C9A	-3.96	1.17	1.30
3	A	402[B]	29P	O1A-C6A	-3.70	1.18	1.30
2	A	401[A]	DPM	C1B-NB	-3.62	1.32	1.37
2	A	401[A]	DPM	O2B-C6B	-3.44	1.19	1.30
2	A	401[A]	DPM	C5B-C2B	3.27	1.55	1.51
3	A	402[B]	29P	C1A-C2A	-3.24	1.32	1.38
2	A	401[A]	DPM	C1A-C2A	-3.23	1.32	1.38
2	A	401[A]	DPM	O4B-C9B	-3.06	1.20	1.30
2	A	401[A]	DPM	O2A-C6A	-2.99	1.20	1.30
2	A	401[A]	DPM	C5A-C6A	2.84	1.55	1.51
3	A	402[B]	29P	C5B-C6B	2.82	1.55	1.51
2	A	401[A]	DPM	CHB-C1B	2.73	1.59	1.49
3	A	402[B]	29P	C2B-C3B	2.64	1.36	1.33
3	A	402[B]	29P	C5A-C6A	2.49	1.54	1.51
3	A	402[B]	29P	O3A-C9A	-2.48	1.22	1.30
3	A	402[B]	29P	C5B-C2B	2.48	1.56	1.50
3	A	402[B]	29P	C4A-C3A	-2.42	1.33	1.39
2	A	401[A]	DPM	C3B-C2B	-2.41	1.35	1.42
2	A	401[A]	DPM	C4A-NA	-2.26	1.32	1.37
2	A	401[A]	DPM	C4A-C3A	-2.23	1.33	1.39
2	A	401[A]	DPM	C1A-NA	-2.15	1.31	1.37
3	A	402[B]	29P	O2B-C6B	-2.11	1.23	1.30
3	A	402[B]	29P	C7B-C3B	2.09	1.56	1.51
3	A	402[B]	29P	C1B-C2B	-2.04	1.44	1.51

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402[B]	29P	CHB-C1B-NB	-5.72	106.72	113.73

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402[B]	29P	CHA-C1A-C2A	-4.07	124.34	131.20
3	A	402[B]	29P	C1B-CHB-C4A	4.05	120.15	113.32
3	A	402[B]	29P	CHB-C4A-NA	3.99	126.44	121.10
3	A	402[B]	29P	O5B-C4B-C3B	-3.80	124.04	128.03
2	A	401[A]	DPM	CHA-C1A-C2A	-3.67	125.00	131.20
3	A	402[B]	29P	CHB-C4A-C3A	-3.54	124.86	130.50
2	A	401[A]	DPM	C2A-C1A-NA	3.44	110.57	108.13
2	A	401[A]	DPM	C2B-C1B-NB	-3.36	105.96	109.84
3	A	402[B]	29P	C2A-C1A-NA	3.30	110.47	108.13
2	A	401[A]	DPM	C7B-C3B-C4B	-3.29	120.58	127.16
3	A	402[B]	29P	O3A-C9A-C8A	2.88	123.09	114.00
2	A	401[A]	DPM	C5B-C2B-C1B	-2.67	123.81	126.54
3	A	402[B]	29P	C1A-C2A-C3A	-2.58	106.49	107.61
2	A	401[A]	DPM	C6B-C5B-C2B	2.55	120.19	113.76
3	A	402[B]	29P	O2A-C6A-C5A	-2.51	114.97	122.11
3	A	402[B]	29P	C6A-C5A-C2A	2.45	119.92	113.76
2	A	401[A]	DPM	O1B-C6B-C5B	-2.43	115.19	122.11
2	A	401[A]	DPM	C1A-C2A-C3A	-2.42	106.56	107.61
2	A	401[A]	DPM	C3B-C4B-NB	-2.35	106.20	109.22
3	A	402[B]	29P	O4A-C9A-C8A	-2.34	115.68	123.09
2	A	401[A]	DPM	O3B-C9B-C8B	-2.20	116.11	123.09
3	A	402[B]	29P	C1B-NB-C4B	-2.20	110.04	113.42
2	A	401[A]	DPM	C6A-C5A-C2A	2.19	119.27	113.76

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	402[B]	29P	C2A-C5A-C6A-O1A
2	A	401[A]	DPM	C2B-C5B-C6B-O1B
3	A	402[B]	29P	C2B-C5B-C6B-O1B
3	A	402[B]	29P	C7B-C8B-C9B-O4B
3	A	402[B]	29P	C2B-C5B-C6B-O2B
3	A	402[B]	29P	C7B-C8B-C9B-O3B
3	A	402[B]	29P	NA-C4A-CHB-C1B
2	A	401[A]	DPM	C2B-C5B-C6B-O2B
3	A	402[B]	29P	C2A-C5A-C6A-O2A


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	290/312 (92%)	-0.12	4 (1%) 73 77	10, 20, 40, 81	8 (2%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	-1	SER	4.4
1	A	177	TRP	2.5
1	A	309	ASP	2.3
1	A	181	THR	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	29P	A	402[B]	31/31	0.70	0.19	11,17,23,29	31
2	DPM	A	401[A]	30/30	0.92	0.10	12,18,22,25	30
4	ACY	A	403	4/4	0.95	0.08	24,27,28,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around DPM A 401 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**6.5 Other polymers** [\(i\)](#)

There are no such residues in this entry.