



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:28 AM UTC

PDB ID : 5MLZ / pdb_00005mlz
Title : Dolichyl phosphate mannose synthase in complex with GDP and Mg2+
Authors : Gandini, R.; Reichenbach, T.; Tan, T.C.; Divne, C.
Deposited on : 2016-12-08
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	374	25% 76% 16% • 5%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3071 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dolichol monophosphate mannose synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	355	Total	C	N	O	S	0	1	0
			2872	1893	483	491	5			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MET	-	initiating methionine	UNP Q8U4M3
A	-20	HIS	-	expression tag	UNP Q8U4M3
A	-19	HIS	-	expression tag	UNP Q8U4M3
A	-18	HIS	-	expression tag	UNP Q8U4M3
A	-17	HIS	-	expression tag	UNP Q8U4M3
A	-16	HIS	-	expression tag	UNP Q8U4M3
A	-15	HIS	-	expression tag	UNP Q8U4M3
A	-14	SER	-	expression tag	UNP Q8U4M3
A	-13	SER	-	expression tag	UNP Q8U4M3
A	-12	GLY	-	expression tag	UNP Q8U4M3
A	-11	VAL	-	expression tag	UNP Q8U4M3
A	-10	ASP	-	expression tag	UNP Q8U4M3
A	-9	LEU	-	expression tag	UNP Q8U4M3
A	-8	GLY	-	expression tag	UNP Q8U4M3
A	-7	THR	-	expression tag	UNP Q8U4M3
A	-6	GLU	-	expression tag	UNP Q8U4M3
A	-5	ASN	-	expression tag	UNP Q8U4M3
A	-4	LEU	-	expression tag	UNP Q8U4M3
A	-3	TYR	-	expression tag	UNP Q8U4M3
A	-2	PHE	-	expression tag	UNP Q8U4M3
A	-1	GLN	-	expression tag	UNP Q8U4M3
A	0	SER	-	expression tag	UNP Q8U4M3

- Molecule 2 is GUANOSINE-5'-DIPHOSPHATE (CCD ID: GDP) (formula: C₁₀H₁₅N₅O₁₁P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			28	10	5	11	2		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		

- Molecule 5 is LAURYL DIMETHYLAMINE-N-OXIDE (CCD ID: LDA) (formula: C₁₄H₃₁NO).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			16	14	1	1		
5	A	1	Total	C	N	O	0	0
			16	14	1	1		
5	A	1	Total	C	N	O	0	0
			16	14	1	1		
5	A	1	Total	C	N	O	0	0
			16	14	1	1		
5	A	1	Total	C	N	O	0	0
			16	14	1	1		
5	A	1	Total	C	N	O	0	0
			16	14	1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	57	Total	O	0	0
			57	57		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dolichol monophosphate mannose synthase



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	90.87Å 146.21Å 95.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.02 – 2.00 41.02 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (41.02-2.00) 99.8 (41.02-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.67 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.241 , 0.257 0.246 , 0.260	Depositor DCC
R_{free} test set	1995 reflections (4.63%)	wwPDB-VP
Wilson B-factor (Å ²)	46.5	Xtriage
Anisotropy	0.559	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 47.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3071	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, CL, LDA, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	4/2942 (0.1%)	0.89	12/3976 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	62	ARG	C-N	13.14	1.52	1.33
1	A	205	GLY	C-N	13.05	1.51	1.33
1	A	63	ARG	C-N	6.08	1.42	1.33
1	A	90	ALA	C-N	-5.80	1.25	1.33

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	62	ARG	CA-C-N	-8.53	110.15	122.94
1	A	62	ARG	C-N-CA	-8.53	110.15	122.94
1	A	46	TRP	N-CA-C	7.74	119.72	111.28
1	A	45	THR	CA-C-N	7.69	130.58	120.28
1	A	45	THR	C-N-CA	7.69	130.58	120.28
1	A	209	LEU	CB-CA-C	-6.83	98.07	109.27
1	A	13	ARG	CB-CA-C	-6.21	102.71	111.65
1	A	42	PRO	CA-C-N	5.82	130.08	120.94
1	A	42	PRO	C-N-CA	5.82	130.08	120.94

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	214	ILE	N-CA-C	5.69	115.88	110.53
1	A	13	ARG	NE-CZ-NH2	5.54	124.19	119.20
1	A	43	ASP	N-CA-CB	-5.00	102.96	110.17

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	204	ARG	Sidechain
1	A	45	THR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2872	0	2993	53	0
2	A	28	0	12	0	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
5	A	112	0	217	8	0
6	A	57	0	0	0	0
All	All	3071	0	3222	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:41:SER:HB3	1:A:43:ASP:HB3	1.47	0.97
1:A:50:GLN:O	1:A:53:SER:HB3	1.78	0.84
1:A:63:ARG:HH22	1:A:71:SER:HB3	1.47	0.79
1:A:63:ARG:HD2	1:A:66:GLU:HB2	1.65	0.78
1:A:202:ARG:HH22	1:A:208:LYS:NZ	1.82	0.77
1:A:45:THR:HA	1:A:47:GLU:N	2.01	0.75

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:TYR:O	1:A:208:LYS:HE3	1.89	0.72
1:A:12:GLU:O	1:A:45:THR:HG21	1.90	0.71
5:A:408:LDA:H102	5:A:409:LDA:H21	1.76	0.68
1:A:270:VAL:HG11	5:A:409:LDA:H31	1.75	0.66
1:A:45:THR:HA	1:A:46:TRP:C	2.25	0.62
1:A:202:ARG:NH2	1:A:206:GLU:O	2.33	0.61
1:A:112:ILE:HB	1:A:192:VAL:HG22	1.83	0.60
1:A:63:ARG:HH22	1:A:71:SER:CB	2.13	0.58
1:A:202:ARG:NH2	1:A:208:LYS:NZ	2.50	0.58
1:A:11:ASN:HA	1:A:43:ASP:OD2	2.03	0.58
1:A:63:ARG:NH2	1:A:71:SER:HB3	2.17	0.58
1:A:215:PHE:O	1:A:219:ARG:HG3	2.04	0.57
1:A:46:TRP:O	1:A:49:ALA:HB3	2.07	0.54
1:A:7:ILE:HD13	1:A:19:LEU:HD21	1.90	0.54
1:A:212:LYS:HG2	1:A:212:LYS:O	2.08	0.53
1:A:253:TRP:CD1	5:A:405:LDA:H11	2.43	0.53
1:A:70:SER:O	1:A:74:ILE:HG12	2.08	0.53
1:A:156:VAL:HB	1:A:178:LYS:HE2	1.91	0.52
1:A:11:ASN:HD22	1:A:11:ASN:N	2.09	0.51
1:A:63:ARG:NH2	1:A:72:ALA:N	2.59	0.51
1:A:-2:PHE:C	1:A:0:SER:H	2.19	0.51
1:A:124:VAL:HG21	1:A:127:TRP:CE2	2.47	0.50
1:A:45:THR:HA	1:A:47:GLU:H	1.73	0.50
1:A:44:ARG:HG3	1:A:45:THR:N	2.27	0.49
1:A:91:ASP:OD1	1:A:93:GLN:HG3	2.12	0.48
1:A:305:ALA:HA	1:A:341:ARG:HD2	1.95	0.48
1:A:63:ARG:HH22	1:A:71:SER:C	2.22	0.48
1:A:124:VAL:HG23	1:A:199:PHE:HB3	1.96	0.48
1:A:167:VAL:HG23	1:A:168:GLU:OE2	2.13	0.48
1:A:202:ARG:CZ	1:A:206:GLU:O	2.61	0.48
1:A:63:ARG:HH22	1:A:72:ALA:N	2.12	0.48
1:A:214:ILE:HG23	1:A:215:PHE:N	2.29	0.47
5:A:408:LDA:H21	5:A:408:LDA:HM23	1.52	0.47
5:A:404:LDA:HM23	5:A:404:LDA:H21	1.61	0.47
1:A:267:ILE:HB	1:A:268:PRO:HD3	1.96	0.47
1:A:91:ASP:CG	1:A:93:GLN:HG3	2.40	0.47
1:A:-1:GLN:HB2	1:A:30:LEU:HD22	1.96	0.46
5:A:407:LDA:H21	5:A:407:LDA:HM23	1.55	0.46
1:A:266:VAL:O	1:A:270:VAL:HG12	2.16	0.46
1:A:11:ASN:HD21	1:A:42:PRO:HD2	1.81	0.46
1:A:61:ILE:HG12	1:A:79:GLU:HG3	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:119:VAL:HG21	1:A:194:GLU:CB	2.47	0.45
5:A:408:LDA:H52	5:A:409:LDA:H111	1.99	0.45
1:A:119:VAL:HG21	1:A:194:GLU:HB3	1.98	0.44
1:A:46:TRP:CD1	1:A:47:GLU:HA	2.53	0.44
5:A:408:LDA:H71	5:A:409:LDA:H32	1.99	0.44
1:A:2:LYS:HB3	1:A:82:GLY:HA2	2.00	0.43
1:A:124:VAL:HG21	1:A:127:TRP:CD2	2.54	0.43
1:A:203:ALA:C	1:A:204:ARG:HG3	2.44	0.42
1:A:265:ALA:O	1:A:268:PRO:HD2	2.20	0.41
1:A:18:GLU:HG2	1:A:92:LEU:HD21	2.03	0.41
1:A:210:LYS:HA	1:A:211:GLY:HA2	1.61	0.41
1:A:41:SER:C	1:A:43:ASP:N	2.79	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	354/374 (95%)	344 (97%)	10 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	311/327 (95%)	306 (98%)	5 (2%)	55 62

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	0	SER
1	A	45	THR
1	A	50	GLN
1	A	216	GLU
1	A	287	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	ASN
1	A	15	ASN
1	A	248	ASN
1	A	345	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	LDA	A	410	-	13,15,15	2.42	2 (15%)	14,17,17	0.54	0
5	LDA	A	404	-	13,15,15	2.40	2 (15%)	14,17,17	0.61	0
5	LDA	A	408	-	13,15,15	2.42	2 (15%)	14,17,17	0.55	0
5	LDA	A	409	-	13,15,15	2.47	2 (15%)	14,17,17	0.52	0
5	LDA	A	406	-	13,15,15	2.39	2 (15%)	14,17,17	0.61	0
5	LDA	A	405	-	13,15,15	2.43	2 (15%)	14,17,17	0.44	0
5	LDA	A	407	-	13,15,15	2.40	2 (15%)	14,17,17	0.47	0
2	GDP	A	401	3	29,30,30	1.14	2 (6%)	45,47,47	1.67	5 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	LDA	A	410	-	-	8/13/13/13	-
5	LDA	A	404	-	-	6/13/13/13	-
5	LDA	A	408	-	-	3/13/13/13	-
5	LDA	A	409	-	-	6/13/13/13	-
5	LDA	A	406	-	-	9/13/13/13	-
5	LDA	A	405	-	-	8/13/13/13	-
5	LDA	A	407	-	-	4/13/13/13	-
2	GDP	A	401	3	-	0/16/32/32	0/3/3/3

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	409	LDA	O1-N1	-6.82	1.25	1.42
5	A	404	LDA	O1-N1	-6.81	1.25	1.42
5	A	410	LDA	O1-N1	-6.80	1.25	1.42
5	A	407	LDA	O1-N1	-6.78	1.25	1.42
5	A	408	LDA	O1-N1	-6.78	1.25	1.42
5	A	406	LDA	O1-N1	-6.77	1.25	1.42
5	A	405	LDA	O1-N1	-6.77	1.25	1.42
5	A	409	LDA	C1-N1	-5.64	1.45	1.51
5	A	405	LDA	C1-N1	-5.50	1.45	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	408	LDA	C1-N1	-5.42	1.45	1.51
5	A	410	LDA	C1-N1	-5.39	1.45	1.51
5	A	407	LDA	C1-N1	-5.33	1.46	1.51
5	A	404	LDA	C1-N1	-5.28	1.46	1.51
5	A	406	LDA	C1-N1	-5.23	1.46	1.51
2	A	401	GDP	C5-C4	3.07	1.47	1.38
2	A	401	GDP	C6-N1	-2.33	1.34	1.38

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	GDP	C5-C4-N3	-5.63	119.43	128.39
2	A	401	GDP	C2-N3-C4	4.79	120.55	112.30
2	A	401	GDP	N9-C4-N3	4.03	134.01	125.95
2	A	401	GDP	C6-C5-N7	3.53	136.71	130.29
2	A	401	GDP	C4-C5-N7	-2.67	106.44	110.67

There are no chirality outliers.

All (44) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	404	LDA	C2-C1-N1-CM2
5	A	406	LDA	C2-C1-N1-CM1
5	A	406	LDA	N1-C1-C2-C3
5	A	409	LDA	C2-C1-N1-CM1
5	A	409	LDA	C2-C1-N1-CM2
5	A	405	LDA	C1-C2-C3-C4
5	A	405	LDA	C7-C8-C9-C10
5	A	410	LDA	C11-C10-C9-C8
5	A	407	LDA	C11-C10-C9-C8
5	A	408	LDA	C7-C8-C9-C10
5	A	410	LDA	C2-C3-C4-C5
5	A	407	LDA	C7-C8-C9-C10
5	A	405	LDA	C4-C5-C6-C7
5	A	408	LDA	C1-C2-C3-C4
5	A	404	LDA	C7-C8-C9-C10
5	A	407	LDA	C1-C2-C3-C4
5	A	404	LDA	C1-C2-C3-C4
5	A	410	LDA	C1-C2-C3-C4
5	A	407	LDA	C9-C10-C11-C12
5	A	405	LDA	C9-C10-C11-C12
5	A	410	LDA	C7-C8-C9-C10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	A	409	LDA	C9-C10-C11-C12
5	A	405	LDA	C3-C4-C5-C6
5	A	404	LDA	C11-C10-C9-C8
5	A	410	LDA	C9-C10-C11-C12
5	A	410	LDA	C5-C6-C7-C8
5	A	406	LDA	C7-C8-C9-C10
5	A	406	LDA	C2-C1-N1-CM2
5	A	410	LDA	C2-C1-N1-CM2
5	A	406	LDA	C2-C1-N1-O1
5	A	409	LDA	C2-C1-N1-O1
5	A	409	LDA	C4-C5-C6-C7
5	A	409	LDA	C7-C8-C9-C10
5	A	410	LDA	C3-C4-C5-C6
5	A	404	LDA	C4-C5-C6-C7
5	A	406	LDA	C6-C7-C8-C9
5	A	405	LDA	C2-C3-C4-C5
5	A	406	LDA	C9-C10-C11-C12
5	A	405	LDA	C5-C6-C7-C8
5	A	406	LDA	C1-C2-C3-C4
5	A	406	LDA	C5-C6-C7-C8
5	A	408	LDA	C5-C6-C7-C8
5	A	405	LDA	C11-C10-C9-C8
5	A	404	LDA	C9-C10-C11-C12


There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	404	LDA	1	0
5	A	408	LDA	4	0
5	A	409	LDA	4	0
5	A	405	LDA	1	0
5	A	407	LDA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	355/374 (94%)	1.35	93 (26%) 1 1	26, 55, 94, 104	1 (0%)

All (93) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	215	PHE	6.9
1	A	46	TRP	6.6
1	A	13	ARG	6.1
1	A	-2	PHE	5.5
1	A	44	ARG	4.8
1	A	207	SER	4.6
1	A	64	THR	4.5
1	A	-1	GLN	4.2
1	A	52	LEU	4.2
1	A	63	ARG	4.1
1	A	127	TRP	4.0
1	A	129	PHE	4.0
1	A	210	LYS	3.9
1	A	177	PHE	3.8
1	A	203	ALA	3.8
1	A	211	GLY	3.8
1	A	43	ASP	3.7
1	A	130	TYR	3.7
1	A	61	ILE	3.7
1	A	42	PRO	3.6
1	A	14	GLU	3.6
1	A	60	VAL	3.6
1	A	39	ASP	3.6
1	A	65	LYS	3.4
1	A	123	LYS	3.4
1	A	45	THR	3.4
1	A	209	LEU	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	232	ASP	3.3
1	A	267	ILE	3.3
1	A	67	LYS	3.3
1	A	208	LYS	3.3
1	A	62	ARG	3.2
1	A	17	GLU	3.2
1	A	58	VAL	3.2
1	A	212	LYS	3.2
1	A	56	TYR	3.2
1	A	0	SER	3.2
1	A	15	ASN	3.0
1	A	213	THR	3.0
1	A	20	PHE	3.0
1	A	128	PRO	3.0
1	A	214	ILE	3.0
1	A	201	ILE	2.9
1	A	11	ASN	2.9
1	A	16	LEU	2.9
1	A	29	GLY	2.9
1	A	133	LEU	2.8
1	A	260	ILE	2.8
1	A	159	PHE	2.8
1	A	10	TYR	2.8
1	A	200	GLY	2.8
1	A	35	VAL	2.8
1	A	175	ILE	2.7
1	A	264	ILE	2.7
1	A	12	GLU	2.7
1	A	92	LEU	2.7
1	A	25	ASN	2.6
1	A	338[A]	PHE	2.6
1	A	204	ARG	2.6
1	A	326	TYR	2.6
1	A	266	VAL	2.6
1	A	327	LEU	2.5
1	A	37	VAL	2.4
1	A	262	LYS	2.4
1	A	306	LEU	2.4
1	A	30	LEU	2.4
1	A	331	LEU	2.4
1	A	265	ALA	2.3
1	A	323	GLY	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	48	LYS	2.3
1	A	40	ASP	2.3
1	A	50	GLN	2.3
1	A	68	GLY	2.3
1	A	38	ASP	2.2
1	A	21	SER	2.2
1	A	168	GLU	2.2
1	A	206	GLU	2.2
1	A	181	MET	2.2
1	A	328	ILE	2.2
1	A	1	MET	2.2
1	A	9	THR	2.2
1	A	34	ILE	2.1
1	A	69	LEU	2.1
1	A	59	LYS	2.1
1	A	118	TYR	2.1
1	A	162	LEU	2.1
1	A	57	PRO	2.1
1	A	101	LYS	2.1
1	A	253	TRP	2.1
1	A	49	ALA	2.1
1	A	258	LEU	2.0
1	A	131	ARG	2.0
1	A	176	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	LDA	A	408	16/16	0.69	0.34	60,72,89,90	0
5	LDA	A	405	16/16	0.70	0.26	67,79,109,113	0
5	LDA	A	409	16/16	0.73	0.29	74,86,95,95	0
5	LDA	A	410	16/16	0.76	0.22	60,77,95,95	0
5	LDA	A	404	16/16	0.77	0.28	59,69,91,93	0
5	LDA	A	407	16/16	0.83	0.22	66,75,89,90	0
5	LDA	A	406	16/16	0.88	0.18	51,56,79,81	0
2	GDP	A	401	28/28	0.90	0.12	48,63,70,71	0
3	MG	A	402	1/1	0.93	0.07	59,59,59,59	0
4	CL	A	403	1/1	0.94	0.10	52,52,52,52	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.