



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 08:29 AM UTC

PDB ID : 6ML1 / pdb_00006ml1
Title : Structure of the USP15 deubiquitinase domain in complex with an affinity-matured inhibitory Ubv
Authors : Singer, A.U.; Teyra, J.; Boehmelt, G.; Lenter, M.; Sicheri, F.; Sidhu, S.S.
Deposited on : 2018-09-26
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	349	 6% 88% 9%
1	B	349	 6% 85% 7% 8%
2	C	92	 12% 92% 5%
2	E	92	 11% 82% 9% 10%
3	G	26	 8% 19% 81%

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 6889 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitin carboxyl-terminal hydrolase 15,Ubiquitin carboxyl-terminal hydrolase 15,Ubiquitin carboxyl-terminal hydrolase 15.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	316	Total	C	N	O	S	0	2	0
			2547	1646	421	464	16			
1	B	321	Total	C	N	O	S	0	4	0
			2583	1661	425	481	16			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	270	SER	-	expression tag	UNP Q9Y4E8
A	271	GLY	-	expression tag	UNP Q9Y4E8
A	272	ALA	-	expression tag	UNP Q9Y4E8
A	273	ALA	-	expression tag	UNP Q9Y4E8
A	274	ALA	-	expression tag	UNP Q9Y4E8
A	872	GLY	-	linker	UNP Q9Y4E8
A	935	SER	-	expression tag	UNP Q9Y4E8
A	936	SER	-	expression tag	UNP Q9Y4E8
A	937	GLY	-	expression tag	UNP Q9Y4E8
B	270	SER	-	expression tag	UNP Q9Y4E8
B	271	GLY	-	expression tag	UNP Q9Y4E8
B	272	ALA	-	expression tag	UNP Q9Y4E8
B	273	ALA	-	expression tag	UNP Q9Y4E8
B	274	ALA	-	expression tag	UNP Q9Y4E8
B	872	GLY	-	linker	UNP Q9Y4E8
B	935	SER	-	expression tag	UNP Q9Y4E8
B	936	SER	-	expression tag	UNP Q9Y4E8
B	937	GLY	-	expression tag	UNP Q9Y4E8

- Molecule 2 is a protein called Ubiquitin variant 15.1a.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	83	Total 647	C 411	N 111	O 124	S 1	0	0	0
2	C	87	Total 651	C 421	N 107	O 122	S 1	0	1	0

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	-1	ALA	-	insertion	UNP A0A0L7RG06
E	0	ALA	GLY	conflict	UNP A0A0L7RG06
E	2	LEU	GLN	conflict	UNP A0A0L7RG06
E	9	SER	THR	conflict	UNP A0A0L7RG06
E	12	PHE	THR	conflict	UNP A0A0L7RG06
E	14	SER	THR	conflict	UNP A0A0L7RG06
E	19	PRO	ALA	conflict	UNP A0A0L7RG06
E	?	-	ASP	deletion	UNP A0A0L7RG06
E	47	GLY	GLN	conflict	UNP A0A0L7RG06
E	48	LYS	GLN	conflict	UNP A0A0L7RG06
E	51	SER	ILE	conflict	UNP A0A0L7RG06
E	53	TYR	ALA	conflict	UNP A0A0L7RG06
E	54	ARG	GLY	conflict	UNP A0A0L7RG06
E	71	HIS	GLU	conflict	UNP A0A0L7RG06
E	75	GLN	HIS	conflict	UNP A0A0L7RG06
E	77	LEU	VAL	conflict	UNP A0A0L7RG06
E	78	VAL	LEU	conflict	UNP A0A0L7RG06
E	79	ILE	ARG	conflict	UNP A0A0L7RG06
E	80	SER	LEU	conflict	UNP A0A0L7RG06
E	83	ILE	-	expression tag	UNP A0A0L7RG06
E	84	LEU	-	expression tag	UNP A0A0L7RG06
E	85	TYR	-	expression tag	UNP A0A0L7RG06
E	86	GLY	-	expression tag	UNP A0A0L7RG06
E	87	SER	-	expression tag	UNP A0A0L7RG06
E	88	SER	-	expression tag	UNP A0A0L7RG06
E	89	GLY	-	expression tag	UNP A0A0L7RG06
C	-1	ALA	-	insertion	UNP A0A0L7RG06
C	0	ALA	GLY	conflict	UNP A0A0L7RG06
C	2	LEU	GLN	conflict	UNP A0A0L7RG06
C	9	SER	THR	conflict	UNP A0A0L7RG06
C	12	PHE	THR	conflict	UNP A0A0L7RG06
C	14	SER	THR	conflict	UNP A0A0L7RG06
C	19	PRO	ALA	conflict	UNP A0A0L7RG06
C	?	-	ASP	deletion	UNP A0A0L7RG06
C	47	GLY	GLN	conflict	UNP A0A0L7RG06

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	48	LYS	GLN	conflict	UNP A0A0L7RG06
C	51	SER	ILE	conflict	UNP A0A0L7RG06
C	53	TYR	ALA	conflict	UNP A0A0L7RG06
C	54	ARG	GLY	conflict	UNP A0A0L7RG06
C	71	HIS	GLU	conflict	UNP A0A0L7RG06
C	75	GLN	HIS	conflict	UNP A0A0L7RG06
C	77	LEU	VAL	conflict	UNP A0A0L7RG06
C	78	VAL	LEU	conflict	UNP A0A0L7RG06
C	79	ILE	ARG	conflict	UNP A0A0L7RG06
C	80	SER	LEU	conflict	UNP A0A0L7RG06
C	83	ILE	-	expression tag	UNP A0A0L7RG06
C	84	LEU	-	expression tag	UNP A0A0L7RG06
C	85	TYR	-	expression tag	UNP A0A0L7RG06
C	86	GLY	-	expression tag	UNP A0A0L7RG06
C	87	SER	-	expression tag	UNP A0A0L7RG06
C	88	SER	-	expression tag	UNP A0A0L7RG06
C	89	GLY	-	expression tag	UNP A0A0L7RG06

- Molecule 3 is a protein called Proteolyzed N-terminal tag of Ubv.15.1a construct.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	G	5	Total	C	N	O	0	0	0
			40	24	11	5			

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Ca	0	0
			2	2		
4	B	1	Total	Ca	0	0
			1	1		
4	C	1	Total	Ca	0	0
			1	1		


- Molecule 5 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Zn	0	0
			1	1		
5	B	1	Total	Zn	0	0
			1	1		

- Molecule 6 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	Na	0	0
			2	2		
6	B	2	Total	Na	0	0
			2	2		

- Molecule 7 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	1	Total	Cl	0	1
			2	2		

- Molecule 9 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	C	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 10 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	155	Total	O	0	2
			157	157		
10	B	175	Total	O	0	2
			177	177		
10	E	33	Total	O	0	0
			33	33		
10	C	23	Total	O	0	0
			23	23		
10	G	3	Total	O	0	0
			3	3		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 15,Ubiquitin carboxyl-terminal hydrolase 15,Ubiquitin carboxyl-terminal hydrolase 15

Chain A: 




- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 15,Ubiquitin carboxyl-terminal hydrolase 15,Ubiquitin carboxyl-terminal hydrolase 15

Chain B: 




- Molecule 2: Ubiquitin variant 15.1a

Chain E: 




- Molecule 2: Ubiquitin variant 15.1a

Chain C: 



- Molecule 3: Proteolyzed N-terminal tag of Ubv.15.1a construct

Chain G:  8% 19% 81%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	42.77Å 115.29Å 95.32Å 90.00° 91.92° 90.00°	Depositor
Resolution (Å)	49.32 – 1.90 49.32 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.0 (49.32-1.90) 98.0 (49.32-1.90)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.55 (at 1.90Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.189 , 0.220 0.191 , 0.222	Depositor DCC
R_{free} test set	3427 reflections (4.06%)	wwPDB-VP
Wilson B-factor (Å ²)	28.8	Xtriage
Anisotropy	0.144	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 48.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.033 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6889	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, NA, CA, MES, ZN, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/2613	0.49	0/3539
1	B	0.32	0/2648	0.50	0/3586
2	C	0.34	1/661 (0.2%)	0.43	0/894
2	E	0.26	0/654	0.43	0/880
3	G	0.25	0/42	0.49	0/56
All	All	0.32	1/6618 (0.0%)	0.48	0/8955

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	50	LEU	CA-C	5.15	1.61	1.52

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2547	0	2466	6	0
1	B	2583	0	2471	18	0
2	C	651	0	638	1	0
2	E	647	0	660	6	0
3	G	40	0	27	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	2	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	2	0	0	0	0
6	B	2	0	0	0	0
7	B	4	0	6	0	0
8	B	2	0	0	1	0
9	C	12	0	12	0	0
10	A	157	0	0	1	0
10	B	177	0	0	5	0
10	C	23	0	0	0	0
10	E	33	0	0	1	0
10	G	3	0	0	0	0
All	All	6889	0	6280	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:420:GLU:OE1	10:B:1101:HOH:O	1.97	0.82
1:B:460:PRO:O	2:E:51:SER:HB2	1.91	0.69
2:E:6:LYS:O	2:E:75:GLN:NE2	2.27	0.68
1:B:841:SER:H	1:B:844:MET:HE3	1.59	0.68
1:A:449:PRO:HD3	1:A:818:ALA:HA	1.78	0.65
1:A:782:LYS:HE3	1:A:849:ASP:OD1	1.98	0.63
2:E:2:LEU:HD11	2:E:14:SER:HB2	1.81	0.63
1:B:409[B]:ASP:OD1	10:B:1102:HOH:O	2.16	0.61
1:B:381:CYS:SG	1:B:382:GLN:N	2.74	0.61
1:B:299:PHE:CE2	1:B:382:GLN:HA	2.36	0.60
1:B:449:PRO:HD3	1:B:818:ALA:HA	1.85	0.58
1:B:873:CYS:SG	10:B:1263:HOH:O	2.58	0.56
1:B:798:LYS:HD2	1:B:798:LYS:H	1.73	0.54
8:B:1004[A]:CL:CL	10:B:1137:HOH:O	2.56	0.52
2:E:55:LYS:NZ	10:E:101:HOH:O	2.14	0.51
1:A:786:VAL:HG22	1:A:790:ASP:HB2	1.93	0.49
1:B:470:MET:HA	1:B:781:PRO:C	2.39	0.48
1:B:321:LYS:NZ	10:B:1111:HOH:O	2.46	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:382:GLN:HB2	1:B:926:TYR:CE1	2.50	0.46
1:A:460:PRO:O	2:C:51:SER:HB2	2.16	0.46
1:A:800:LYS:HA	1:A:819:THR:HA	1.98	0.45
1:B:786:VAL:HG22	1:B:790:ASP:HB2	1.97	0.45
1:A:844:MET:HA	10:A:1128:HOH:O	2.18	0.42
1:B:467:PRO:HG2	1:B:794:LEU:HB3	2.01	0.42
1:B:451:CYS:SG	1:B:811:ASN:ND2	2.82	0.42
1:B:320:ASP:OD1	1:B:347:LYS:NZ	2.33	0.42
1:B:383:GLU:H	1:B:383:GLU:HG2	1.68	0.42
2:E:8:LEU:HD23	2:E:8:LEU:HA	1.76	0.41
1:B:903:LYS:HB3	1:B:903:LYS:HE2	1.78	0.41
2:E:6:LYS:HD2	2:E:12:PHE:CZ	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	308/349 (88%)	305 (99%)	3 (1%)	0	100	100
1	B	315/349 (90%)	307 (98%)	8 (2%)	0	100	100
2	C	86/92 (94%)	84 (98%)	2 (2%)	0	100	100
2	E	79/92 (86%)	79 (100%)	0	0	100	100
3	G	3/26 (12%)	3 (100%)	0	0	100	100
All	All	791/908 (87%)	778 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	276/306 (90%)	276 (100%)	0	100	100
1	B	279/306 (91%)	279 (100%)	0	100	100
2	C	66/80 (82%)	66 (100%)	0	100	100
2	E	71/80 (89%)	71 (100%)	0	100	100
3	G	3/22 (14%)	3 (100%)	0	100	100
All	All	695/794 (88%)	695 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	857	ASN
1	A	883	HIS
1	B	365	GLN
2	E	56	GLN
2	C	75	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 14 ligands modelled in this entry, 12 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	EDO	B	1003	-	3,3,3	0.48	0	2,2,2	0.52	0
9	MES	C	102	-	12,12,12	2.32	1 (8%)	15,16,16	2.37	7 (46%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	EDO	B	1003	-	-	0/1/1/1	-
9	MES	C	102	-	-	2/6/14/14	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	C	102	MES	C8-S	-7.71	1.66	1.77

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	C	102	MES	C5-N4-C3	5.39	120.45	108.84
9	C	102	MES	C7-N4-C3	3.16	119.65	111.24
9	C	102	MES	O3S-S-C8	2.84	111.56	106.00
9	C	102	MES	C7-N4-C5	2.78	118.64	111.24
9	C	102	MES	C8-C7-N4	-2.58	102.60	112.36
9	C	102	MES	C6-C5-N4	-2.45	106.39	110.12
9	C	102	MES	C2-C3-N4	-2.44	106.41	110.12

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
9	C	102	MES	N4-C7-C8-S
9	C	102	MES	C8-C7-N4-C3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	316/349 (90%)	0.24	21 (6%) 24 26	13, 31, 58, 74	2 (0%)
1	B	321/349 (91%)	0.30	20 (6%) 26 28	11, 31, 57, 77	4 (1%)
2	C	87/92 (94%)	1.22	11 (12%) 8 8	24, 49, 66, 77	1 (1%)
2	E	83/92 (90%)	0.94	10 (12%) 9 9	26, 46, 63, 80	0
3	G	5/26 (19%)	1.80	2 (40%) 1 0	41, 43, 47, 63	0
All	All	812/908 (89%)	0.45	64 (7%) 18 20	11, 34, 61, 80	7 (0%)

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	52[A]	PHE	7.4
2	C	83	ILE	4.2
2	E	84	LEU	4.0
2	E	50	LEU	3.9
2	C	-1	ALA	3.9
3	G	12	ALA	3.9
2	C	85	TYR	3.8
1	B	382	GLN	3.7
2	E	83	ILE	3.7
2	E	-1	ALA	3.6
1	B	294	LEU	3.6
1	A	381	CYS	3.5
1	B	295	GLY	3.3
1	B	937	GLY	3.3
1	A	840	TYR	3.2
1	B	381	CYS	3.2
2	C	8	LEU	3.2
1	A	890	GLY	3.2
2	E	79	ILE	3.1
1	B	299	PHE	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	E	51	SER	3.1
2	C	51	SER	3.0
1	B	470	MET	3.0
2	E	52	PHE	2.8
1	A	801	LEU	2.7
1	A	802	GLY	2.7
1	A	812	CYS	2.7
1	A	838	PHE	2.7
1	B	806	PRO	2.7
2	C	84	LEU	2.7
1	A	848	LEU	2.6
1	A	294	LEU	2.6
2	C	50	LEU	2.6
2	C	49	ARG	2.6
3	G	11	MET	2.6
1	A	295	GLY	2.6
1	A	412	GLY	2.5
1	B	803	ALA	2.5
1	B	844	MET	2.5
1	A	843	TYR	2.5
2	C	79	ILE	2.4
1	B	798	LYS	2.4
1	A	407	LEU	2.3
1	A	808	TYR	2.3
1	B	356	TYR	2.3
1	B	463	TYR	2.3
1	B	840	TYR	2.3
1	A	299	PHE	2.3
1	B	373	PHE	2.3
1	B	371	PRO	2.3
1	A	806	PRO	2.2
1	B	935	SER	2.2
1	B	283	ASN	2.2
2	E	10	GLY	2.2
1	B	885	GLY	2.2
1	A	798	LYS	2.2
2	E	53	TYR	2.1
1	B	412	GLY	2.1
1	A	839	SER	2.1
1	A	845	ARG	2.1
1	A	442	PHE	2.0
1	A	453	LYS	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	12	PHE	2.0
2	E	35	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
9	MES	C	102	12/12	0.62	0.18	70,87,99,109	0
7	EDO	B	1003	4/4	0.76	0.15	45,47,47,52	0
6	NA	B	1006	1/1	0.92	0.12	45,45,45,45	0
8	CL	B	1004[A]	1/1	0.93	0.08	47,47,47,47	1
8	CL	B	1004[B]	1/1	0.93	0.08	37,37,37,37	1
4	CA	C	101	1/1	0.93	0.07	62,62,62,62	0
6	NA	A	1004	1/1	0.94	0.10	50,50,50,50	0
4	CA	B	1001	1/1	0.94	0.07	62,62,62,62	0
5	ZN	B	1002	1/1	0.96	0.05	39,39,39,39	0
6	NA	B	1005	1/1	0.97	0.07	28,28,28,28	0
5	ZN	A	1003	1/1	0.98	0.06	47,47,47,47	0
4	CA	A	1002	1/1	0.99	0.07	31,31,31,31	0
4	CA	A	1001	1/1	0.99	0.02	19,19,19,19	0
6	NA	A	1005	1/1	0.99	0.03	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.