



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:52 PM UTC

PDB ID : 1MM7 / pdb_00001mm7
Title : Crystal Structure of the GluR2 Ligand Binding Core (S1S2J) in Complex with Quisqualate in a Zinc Crystal Form at 1.65 Angstroms Resolution
Authors : Jin, R.; Horning, M.; Mayer, M.L.; Gouaux, E.
Deposited on : 2002-09-03
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2662 (1.66-1.66)
Ramachandran outliers	187476	2621 (1.66-1.66)
Sidechain outliers	187428	2621 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	263	 89% 9% .
1	B	263	 87% 10% ..
1	C	263	 83% 13% ...

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6669 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLUTAMATE RECEPTOR 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	258	Total	C	N	O	S	0	0	0
			1950	1245	321	370	14			
1	B	259	Total	C	N	O	S	0	0	0
			1978	1260	327	377	14			
1	C	258	Total	C	N	O	S	0	0	0
			1923	1225	319	366	13			


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	cloning artifact	UNP P19491
A	2	ALA	-	cloning artifact	UNP P19491
A	118	GLY	-	linker	UNP P19491
A	119	THR	-	linker	UNP P19491
B	1	GLY	-	cloning artifact	UNP P19491
B	2	ALA	-	cloning artifact	UNP P19491
B	118	GLY	-	linker	UNP P19491
B	119	THR	-	linker	UNP P19491
C	1	GLY	-	cloning artifact	UNP P19491
C	2	ALA	-	cloning artifact	UNP P19491
C	118	GLY	-	linker	UNP P19491
C	119	THR	-	linker	UNP P19491

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	2	Total	Zn	0	0
			2	2		
2	C	2	Total	Zn	0	0
			2	2		

- Molecule 3 is (S)-2-AMINO-3-(3,5-DIOXO-[1,2,4]OXADIAZOLIDIN-2-YL)-PROPIONIC ACID (CCD ID: QUS) (formula: C₅H₇N₃O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			13	5	3	5		
3	B	1	Total	C	N	O	0	0
			13	5	3	5		
3	C	1	Total	C	N	O	0	0
			13	5	3	5		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	277	Total	O	0	0
			277	277		
4	B	289	Total	O	0	0
			289	289		
4	C	208	Total	O	0	0
			208	208		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: GLUTAMATE RECEPTOR 2

Chain A:  89% 9% .




• Molecule 1: GLUTAMATE RECEPTOR 2

Chain B:  87% 10% ..



• Molecule 1: GLUTAMATE RECEPTOR 2

Chain C:  83% 13% . .



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	113.91Å 163.45Å 47.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.85 – 1.65	Depositor
% Data completeness (in resolution range)	90.2 (29.85-1.65)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.204 , 0.237	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6669	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: QUS, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/1986	0.96	8/2679 (0.3%)
1	B	0.57	0/2014	0.95	8/2716 (0.3%)
1	C	0.56	0/1958	0.98	10/2641 (0.4%)
All	All	0.57	0/5958	0.96	26/8036 (0.3%)

There are no bond length outliers.

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	259	GLY	N-CA-C	-7.46	101.39	111.54
1	B	27	GLU	N-CA-C	6.73	119.09	108.79
1	A	116	LYS	N-CA-C	-6.35	100.44	109.96
1	B	116	LYS	N-CA-C	-6.17	100.46	110.20
1	A	184	SER	N-CA-C	6.02	120.25	113.02
1	C	260	GLU	N-CA-C	5.68	119.76	112.26
1	A	256	TYR	N-CA-C	5.62	120.64	113.55
1	C	116	LYS	N-CA-C	-5.57	101.40	110.20
1	B	150	SER	N-CA-C	5.54	118.27	110.23
1	C	165	ALA	N-CA-C	5.51	118.01	110.24
1	B	29	ASN	N-CA-C	5.47	118.17	111.82
1	B	140	SER	N-CA-C	5.45	119.23	112.58
1	C	153	ALA	N-CA-C	5.43	122.37	110.80
1	B	103	SER	N-CA-C	-5.39	103.57	110.53
1	A	103	SER	N-CA-C	-5.33	103.65	110.53
1	A	57	GLY	N-CA-C	5.32	119.32	112.83
1	C	256	TYR	N-CA-C	5.30	116.91	111.03
1	A	151	LYS	N-CA-C	-5.28	106.07	112.88
1	A	138	LEU	N-CA-C	-5.22	102.32	110.10
1	B	57	GLY	N-CA-C	5.22	118.96	112.64
1	C	57	GLY	N-CA-C	5.16	119.13	112.83

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	119	THR	CA-C-N	5.15	126.28	119.84
1	C	119	THR	C-N-CA	5.15	126.28	119.84
1	C	105	PRO	N-CA-C	5.13	119.14	111.14
1	B	105	PRO	N-CA-C	5.11	119.11	111.14
1	A	37	VAL	N-CA-C	-5.09	105.74	110.53

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1950	0	1913	10	0
1	B	1978	0	1958	17	0
1	C	1923	0	1879	22	0
2	A	1	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
3	A	13	0	6	0	0
3	B	13	0	6	0	0
3	C	13	0	6	0	0
4	A	277	0	0	4	0
4	B	289	0	0	6	0
4	C	208	0	0	3	0
All	All	6669	0	5768	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:64:ARG:HH12	1:C:66:ALA:HA	1.39	0.87
1:B:130:GLN:HE22	1:B:133:ILE:H	1.21	0.84
1:A:25:MET:HG2	4:A:951:HOH:O	1.83	0.77

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:210:LYS:HE2	4:A:1177:HOH:O	1.89	0.71
1:C:199:TYR:CZ	1:C:203:ARG:HD2	2.28	0.68
1:B:249:LYS:HE2	4:B:1117:HOH:O	1.95	0.67
1:B:252:ASN:HA	1:B:256:TYR:HD2	1.63	0.63
1:B:81:GLY:HA2	4:B:1166:HOH:O	2.01	0.59
1:B:163:ARG:CG	4:B:1088:HOH:O	2.50	0.58
1:A:244:GLN:HB3	4:A:1174:HOH:O	2.02	0.58
1:C:46:HIS:ND1	4:C:1003:HOH:O	2.29	0.58
1:C:131:THR:CA	4:C:1066:HOH:O	2.52	0.57
1:C:64:ARG:HH12	1:C:66:ALA:CA	2.18	0.54
1:C:64:ARG:NH1	1:C:65:ASP:O	2.40	0.54
1:B:130:GLN:HE22	1:B:133:ILE:N	2.01	0.53
1:B:252:ASN:HA	1:B:256:TYR:CD2	2.43	0.53
1:A:244:GLN:HA	1:A:244:GLN:OE1	2.10	0.52
4:A:1174:HOH:O	1:B:166:GLU:HB2	2.08	0.52
1:A:209:MET:HE2	1:A:211:VAL:HG12	1.92	0.51
1:B:256:TYR:HE1	4:B:1089:HOH:O	1.92	0.51
1:C:148:ARG:NH1	4:C:1050:HOH:O	2.45	0.50
1:B:23:HIS:CD2	1:B:26:LEU:HD12	2.47	0.49
1:C:130:GLN:OE1	1:C:132:GLU:N	2.44	0.49
1:C:64:ARG:HD2	1:C:69:LYS:HA	1.94	0.49
1:A:42:GLU:HG3	1:A:246:LEU:HD21	1.95	0.49
1:C:134:ALA:O	1:C:188:TYR:HA	2.13	0.48
1:A:22:ASN:CB	1:A:25:MET:HE3	2.44	0.47
1:C:136:GLY:HA3	1:C:169:VAL:O	2.15	0.46
1:C:94:LEU:O	1:C:98:GLU:HG3	2.16	0.46
1:C:130:GLN:NE2	1:C:133:ILE:HD12	2.30	0.46
1:B:199:TYR:O	1:B:203:ARG:HG2	2.16	0.46
1:B:27:GLU:CB	4:B:1172:HOH:O	2.63	0.45
1:B:226:LYS:HE3	4:B:1055:HOH:O	2.15	0.45
1:B:130:GLN:NE2	1:B:133:ILE:H	2.01	0.44
1:C:15:PRO:HB3	1:C:199:TYR:CE1	2.53	0.44
1:C:88:ALA:HB1	1:C:89:PRO:HD2	2.00	0.43
1:A:88:ALA:HB1	1:A:89:PRO:HD2	1.99	0.43
1:C:203:ARG:HA	1:C:260:GLU:HG2	2.01	0.43
1:C:206:CYS:CB	1:C:261:CYS:SG	3.06	0.43
1:B:100:ILE:HD12	1:B:223:ALA:HB1	2.00	0.43
1:C:22:ASN:CG	1:C:22:ASN:O	2.61	0.42
1:B:128:SER:O	1:B:157:LYS:HE2	2.20	0.42
1:B:94:LEU:O	1:B:98:GLU:HG3	2.21	0.41
1:C:119:THR:HA	1:C:120:PRO:HD3	1.82	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:23:HIS:HA	1:C:26:LEU:HD12	2.03	0.41
1:C:46:HIS:CE1	1:C:240:LYS:HD3	2.55	0.41
1:A:251:LYS:O	1:A:255:TRP:HB2	2.21	0.40
1:C:180:ARG:O	1:C:184:SER:HB3	2.21	0.40
1:A:91:THR:HG21	1:A:218:LYS:HD2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	256/263 (97%)	252 (98%)	3 (1%)	1 (0%)	30	15
1	B	257/263 (98%)	251 (98%)	6 (2%)	0	100	100
1	C	256/263 (97%)	247 (96%)	8 (3%)	1 (0%)	30	15
All	All	769/789 (98%)	750 (98%)	17 (2%)	2 (0%)	36	21

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	67	ASP
1	C	153	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	199/219 (91%)	198 (100%)	1 (0%)	81	72
1	B	206/219 (94%)	203 (98%)	3 (2%)	57	37
1	C	196/219 (90%)	191 (97%)	5 (3%)	40	17
All	All	601/657 (92%)	592 (98%)	9 (2%)	57	37

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	109	LEU
1	B	109	LEU
1	B	130	GLN
1	B	139	ASP
1	C	15	PRO
1	C	22	ASN
1	C	64	ARG
1	C	109	LEU
1	C	130	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	23	HIS
1	B	130	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 8 ligands modelled in this entry, 5 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	QUS	C	803	-	9,13,13	1.07	0	10,18,18	1.46	2 (20%)
3	QUS	A	801	-	9,13,13	1.16	1 (11%)	10,18,18	1.31	1 (10%)
3	QUS	B	802	-	9,13,13	0.95	0	10,18,18	1.39	2 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	QUS	C	803	-	-	2/6/8/8	0/1/1/1
3	QUS	A	801	-	-	2/6/8/8	0/1/1/1
3	QUS	B	802	-	-	2/6/8/8	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	801	QUS	O18-C04	-2.12	1.19	1.23

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	802	QUS	O20-N14-C03	2.29	118.38	113.83
3	C	803	QUS	O20-C05-N15	2.26	108.18	106.14
3	A	801	QUS	O20-N14-C03	2.15	118.10	113.83
3	C	803	QUS	O18-C04-N15	2.10	130.40	126.47
3	B	802	QUS	O20-C05-N15	2.08	108.02	106.14

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	801	QUS	C01-C02-C03-N14
3	B	802	QUS	C01-C02-C03-N14
3	C	803	QUS	C01-C02-C03-N14
3	A	801	QUS	NP3-C02-C03-N14
3	B	802	QUS	NP3-C02-C03-N14
3	C	803	QUS	NP3-C02-C03-N14

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.