



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 03:02 PM UTC

PDB ID : 4MMK / pdb_00004mmk
Title : Q8A Hfq from Pseudomonas aeruginosa
Authors : Murina, V.N.; Filimonov, V.V.; Melnik, B.S.; Uhlein, M.; Mueller, U.; Weiss, M.; Nikulin, A.D.
Deposited on : 2013-09-09
Resolution : 2.16 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	82	62% 20% • 16%
1	B	82	59% 23% • 16%
1	C	82	56% 26% •• 16%
1	D	82	56% 26% • 17%
1	E	82	59% 24% • 16%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	82	
1	G	82	
1	H	82	
1	I	82	
1	J	82	
1	K	82	
1	L	82	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6928 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein hfq.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	69	Total	C	N	O	S	0	0	0
			549	354	97	97	1			
1	B	69	Total	C	N	O	S	0	0	0
			549	354	97	97	1			
1	C	69	Total	C	N	O	S	0	0	0
			549	354	97	97	1			
1	D	68	Total	C	N	O	S	0	0	0
			541	350	95	95	1			
1	E	69	Total	C	N	O	S	0	0	0
			549	354	97	97	1			
1	F	68	Total	C	N	O	S	0	0	0
			541	348	96	96	1			
1	G	66	Total	C	N	O	S	0	0	0
			526	339	93	93	1			
1	H	69	Total	C	N	O	S	0	0	0
			549	354	97	97	1			
1	I	70	Total	C	N	O	S	0	0	0
			556	359	98	98	1			
1	J	69	Total	C	N	O	S	0	0	0
			550	356	97	96	1			
1	K	70	Total	C	N	O	S	0	0	0
			556	359	98	98	1			
1	L	70	Total	C	N	O	S	0	0	0
			556	359	98	98	1			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	8	ALA	GLN	engineered mutation	UNP Q9HUM0
B	8	ALA	GLN	engineered mutation	UNP Q9HUM0
C	8	ALA	GLN	engineered mutation	UNP Q9HUM0
D	8	ALA	GLN	engineered mutation	UNP Q9HUM0
E	8	ALA	GLN	engineered mutation	UNP Q9HUM0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	8	ALA	GLN	engineered mutation	UNP Q9HUM0
G	8	ALA	GLN	engineered mutation	UNP Q9HUM0
H	8	ALA	GLN	engineered mutation	UNP Q9HUM0
I	8	ALA	GLN	engineered mutation	UNP Q9HUM0
J	8	ALA	GLN	engineered mutation	UNP Q9HUM0
K	8	ALA	GLN	engineered mutation	UNP Q9HUM0
L	8	ALA	GLN	engineered mutation	UNP Q9HUM0

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Zn 2 2	0	0
2	B	2	Total Zn 2 2	0	0
2	C	2	Total Zn 2 2	0	0
2	D	2	Total Zn 2 2	0	0
2	E	2	Total Zn 2 2	0	0
2	F	2	Total Zn 2 2	0	0
2	G	1	Total Zn 1 1	0	0
2	H	2	Total Zn 2 2	0	0
2	I	3	Total Zn 3 3	0	0
2	J	1	Total Zn 1 1	0	0
2	K	2	Total Zn 2 2	0	0
2	L	2	Total Zn 2 2	0	0

- Molecule 3 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total K 1 1	0	0
3	B	1	Total K 1 1	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	2	Total K 2 2	0	0
3	D	1	Total K 1 1	0	0
3	F	1	Total K 1 1	0	0
3	I	1	Total K 1 1	0	0
3	L	1	Total K 1 1	0	0

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	G	1	Total Na 1 1	0	0
4	J	1	Total Na 1 1	0	0
4	K	1	Total Na 1 1	0	0

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	H	1	Total O S 5 4 1	0	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	33	Total 33	O 33	0	0
6	B	28	Total 28	O 28	0	0
6	C	30	Total 30	O 30	0	0
6	D	26	Total 26	O 26	0	0
6	E	34	Total 34	O 34	0	0
6	F	33	Total 33	O 33	0	0
6	G	11	Total 11	O 11	0	0
6	H	18	Total 18	O 18	0	0
6	I	22	Total 22	O 22	0	0
6	J	22	Total 22	O 22	0	0
6	K	37	Total 37	O 37	0	0
6	L	24	Total 24	O 24	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Protein hfq

Chain A: 



- Molecule 1: Protein hfq

Chain B: 



- Molecule 1: Protein hfq

Chain C: 



- Molecule 1: Protein hfq

Chain D: 




- Molecule 1: Protein hfq

Chain E: 



- Molecule 1: Protein hfq

Chain F: 



• Molecule 1: Protein hfq



• Molecule 1: Protein hfq




• Molecule 1: Protein hfq




• Molecule 1: Protein hfq



• Molecule 1: Protein hfq



• Molecule 1: Protein hfq



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.59Å 116.54Å 66.62Å 90.00° 119.98° 90.00°	Depositor
Resolution (Å)	33.31 – 2.16 33.31 – 2.16	Depositor EDS
% Data completeness (in resolution range)	96.0 (33.31-2.16) 91.9 (33.31-2.16)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.38 (at 2.16Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1418)	Depositor
R, R_{free}	0.216 , 0.256 0.187 , 0.194	Depositor DCC
R_{free} test set	2279 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	18.5	Xtriage
Anisotropy	1.080	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 32.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	0.226 for -h-l,k,h 0.226 for l,k,-h-l 0.309 for h,-k,-h-l 0.226 for -h-l,-k,l 0.229 for l,-k,h	Xtriage
F_o , F_c correlation	0.94	EDS
Total number of atoms	6928	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.48 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.6154e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SO4, NA, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	0/559	0.89	0/757
1	B	0.55	0/559	0.88	2/757 (0.3%)
1	C	0.59	0/559	0.90	0/757
1	D	0.53	0/552	0.93	1/750 (0.1%)
1	E	0.52	0/559	0.87	0/757
1	F	0.52	0/551	0.94	2/746 (0.3%)
1	G	0.50	0/536	1.01	2/727 (0.3%)
1	H	0.58	0/559	0.98	2/757 (0.3%)
1	I	0.53	0/567	0.95	0/769
1	J	0.50	0/561	0.87	0/761
1	K	0.51	0/567	0.89	2/769 (0.3%)
1	L	0.58	1/567 (0.2%)	0.98	4/769 (0.5%)
All	All	0.54	1/6696 (0.0%)	0.92	15/9076 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	L	63	VAL	CA-CB	5.38	1.59	1.53

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	63	VAL	CA-C-N	6.51	127.98	119.84
1	F	63	VAL	C-N-CA	6.51	127.98	119.84
1	L	66	ARG	CA-C-N	6.43	127.88	119.84
1	L	66	ARG	C-N-CA	6.43	127.88	119.84
1	L	5	HIS	N-CA-C	-6.31	106.28	112.97

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	549	0	581	13	0
1	B	549	0	581	15	0
1	C	549	0	581	19	0
1	D	541	0	570	18	1
1	E	549	0	581	16	0
1	F	541	0	570	27	0
1	G	526	0	552	21	0
1	H	549	0	581	24	0
1	I	556	0	588	27	1
1	J	550	0	583	16	0
1	K	556	0	588	23	0
1	L	556	0	588	24	1
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
2	E	2	0	0	0	0
2	F	2	0	0	0	0
2	G	1	0	0	0	0
2	H	2	0	0	0	0
2	I	3	0	0	0	0
2	J	1	0	0	0	0
2	K	2	0	0	0	0
2	L	2	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	2	0	0	0	0
3	D	1	0	0	0	0
3	F	1	0	0	0	0
3	I	1	0	0	0	0
3	L	1	0	0	0	0
4	G	1	0	0	0	0
4	J	1	0	0	0	0
4	K	1	0	0	0	0
5	H	5	0	0	0	0
6	A	33	0	0	1	1
6	B	28	0	0	2	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	C	30	0	0	4	0
6	D	26	0	0	1	0
6	E	34	0	0	3	0
6	F	33	0	0	1	0
6	G	11	0	0	1	0
6	H	18	0	0	2	0
6	I	22	0	0	1	0
6	J	22	0	0	1	0
6	K	37	0	0	4	0
6	L	24	0	0	2	1
All	All	6928	0	6944	208	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 208 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:69:ARG:HD3	1:D:70:LEU:H	1.33	0.90
1:D:69:ARG:HD3	1:D:70:LEU:N	1.93	0.83
1:E:9:ASP:OD1	6:E:230:HOH:O	1.98	0.81
1:K:9:ASP:OD1	6:K:237:HOH:O	2.00	0.80
1:C:18:GLU:OE1	1:C:66:ARG:NH2	2.12	0.80

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:3:LYS:NZ	6:B:227:HOH:O[2_556]	1.96	0.24
1:D:18:GLU:OE2	6:A:224:HOH:O[1_554]	2.10	0.10
1:I:66:ARG:NH2	6:L:217:HOH:O[1_554]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	67/82 (82%)	65 (97%)	0	2 (3%)	3	0
1	B	67/82 (82%)	63 (94%)	4 (6%)	0	100	100
1	C	67/82 (82%)	62 (92%)	4 (6%)	1 (2%)	8	4
1	D	66/82 (80%)	59 (89%)	7 (11%)	0	100	100
1	E	67/82 (82%)	59 (88%)	8 (12%)	0	100	100
1	F	66/82 (80%)	59 (89%)	6 (9%)	1 (2%)	8	4
1	G	64/82 (78%)	60 (94%)	3 (5%)	1 (2%)	7	3
1	H	67/82 (82%)	59 (88%)	8 (12%)	0	100	100
1	I	68/82 (83%)	61 (90%)	6 (9%)	1 (2%)	8	4
1	J	67/82 (82%)	58 (87%)	9 (13%)	0	100	100
1	K	68/82 (83%)	66 (97%)	1 (2%)	1 (2%)	8	4
1	L	68/82 (83%)	59 (87%)	7 (10%)	2 (3%)	3	1
All	All	802/984 (82%)	730 (91%)	63 (8%)	9 (1%)	11	7

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	48	ASN
1	A	49	THR
1	I	6	SER
1	C	49	THR
1	K	48	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	64/73 (88%)	64 (100%)	0	100	100
1	B	64/73 (88%)	59 (92%)	5 (8%)	11	7

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	64/73 (88%)	62 (97%)	2 (3%)	35	37
1	D	63/73 (86%)	62 (98%)	1 (2%)	55	61
1	E	64/73 (88%)	61 (95%)	3 (5%)	23	21
1	F	63/73 (86%)	59 (94%)	4 (6%)	16	11
1	G	61/73 (84%)	58 (95%)	3 (5%)	22	19
1	H	64/73 (88%)	60 (94%)	4 (6%)	16	11
1	I	65/73 (89%)	62 (95%)	3 (5%)	24	22
1	J	64/73 (88%)	60 (94%)	4 (6%)	16	11
1	K	65/73 (89%)	63 (97%)	2 (3%)	35	37
1	L	65/73 (89%)	62 (95%)	3 (5%)	24	22
All	All	766/876 (87%)	732 (96%)	34 (4%)	25	23

5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	J	70	LEU
1	K	51	SER
1	L	68	VAL
1	F	14	THR
1	F	2	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	5	HIS
1	L	33	GLN
1	F	52	GLN
1	L	52	GLN
1	J	52	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 35 ligands modelled in this entry, 34 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	SO4	H	103	-	4,4,4	0.26	0	6,6,6	0.17	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	69/82 (84%)	-1.14	0 100 100	10, 18, 30, 35	0
1	B	69/82 (84%)	-1.12	0 100 100	8, 20, 33, 36	0
1	C	69/82 (84%)	-1.05	0 100 100	11, 22, 33, 41	0
1	D	68/82 (82%)	-1.11	0 100 100	11, 23, 37, 47	0
1	E	69/82 (84%)	-1.11	0 100 100	12, 24, 35, 37	0
1	F	68/82 (82%)	-1.10	0 100 100	11, 22, 32, 34	0
1	G	66/82 (80%)	-0.97	0 100 100	21, 27, 38, 45	0
1	H	69/82 (84%)	-0.96	0 100 100	16, 28, 40, 42	0
1	I	70/82 (85%)	-0.96	0 100 100	15, 25, 36, 43	0
1	J	69/82 (84%)	-1.09	0 100 100	13, 23, 37, 40	0
1	K	70/82 (85%)	-1.02	0 100 100	15, 24, 36, 51	0
1	L	70/82 (85%)	-0.96	0 100 100	18, 29, 40, 48	0
All	All	826/984 (83%)	-1.05	0 100 100	8, 24, 36, 51	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	A	102	1/1	0.98	0.03	45,45,45,45	0
3	K	C	103	1/1	0.98	0.05	44,44,44,44	0
3	K	F	103	1/1	0.98	0.02	29,29,29,29	0
4	NA	G	102	1/1	0.98	0.05	30,30,30,30	0
4	NA	J	102	1/1	0.98	0.07	39,39,39,39	0
5	SO4	H	103	5/5	0.98	0.06	36,39,45,46	0
2	ZN	I	101	1/1	0.99	0.02	30,30,30,30	0
2	ZN	I	103	1/1	0.99	0.03	49,49,49,49	0
2	ZN	J	101	1/1	0.99	0.02	37,37,37,37	0
2	ZN	L	102	1/1	0.99	0.03	54,54,54,54	0
2	ZN	B	101	1/1	0.99	0.02	28,28,28,28	0
3	K	C	104	1/1	0.99	0.05	31,31,31,31	0
3	K	D	103	1/1	0.99	0.03	26,26,26,26	0
2	ZN	C	101	1/1	0.99	0.01	25,25,25,25	0
3	K	I	104	1/1	0.99	0.02	46,46,46,46	0
3	K	L	103	1/1	0.99	0.05	38,38,38,38	0
2	ZN	C	102	1/1	0.99	0.03	36,36,36,36	0
2	ZN	D	101	1/1	0.99	0.02	30,30,30,30	0
4	NA	K	103	1/1	0.99	0.05	30,30,30,30	0
2	ZN	D	102	1/1	0.99	0.03	41,41,41,41	0
2	ZN	K	102	1/1	1.00	0.01	45,45,45,45	0
2	ZN	L	101	1/1	1.00	0.03	35,35,35,35	0
2	ZN	E	102	1/1	1.00	0.02	38,38,38,38	0
3	K	A	103	1/1	1.00	0.02	31,31,31,31	0
3	K	B	103	1/1	1.00	0.05	26,26,26,26	0
2	ZN	F	101	1/1	1.00	0.02	33,33,33,33	0
2	ZN	F	102	1/1	1.00	0.03	34,34,34,34	0
2	ZN	G	101	1/1	1.00	0.02	34,34,34,34	0
2	ZN	H	101	1/1	1.00	0.01	35,35,35,35	0
2	ZN	H	102	1/1	1.00	0.03	43,43,43,43	0
2	ZN	A	101	1/1	1.00	0.01	27,27,27,27	0
2	ZN	I	102	1/1	1.00	0.01	33,33,33,33	0
2	ZN	B	102	1/1	1.00	0.01	32,32,32,32	0
2	ZN	E	101	1/1	1.00	0.01	26,26,26,26	0
2	ZN	K	101	1/1	1.00	0.01	35,35,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.