



# wwPDB X-ray Structure Validation Summary Report

Mar 5, 2026 – 11:42 AM UTC

PDB ID : 4MMS / pdb\_00004mms  
Title : Crystal Structure of Prefusion-stabilized RSV F Variant Cav1 at pH 5.5  
Authors : Mclellan, J.S.; Joyce, M.G.; Stewart-Jones, G.B.E.; Sastry, M.; Yang, Y.;  
Graham, B.S.; Kwong, P.D.  
Deposited on : 2013-09-09  
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	82	 13% 88% 10% ..
1	C	82	 15% 77% 17% 6%
1	E	82	 18% 76% 22% ..
2	B	414	 7% 78% 11% 11%
2	D	414	 7% 76% 13% 11%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	414	 <p>6% 76% 12% • 11%</p>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11013 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fusion glycoprotein F2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	81	Total 635	C 397	N 107	O 128	S 3	0	0	0
1	C	77	Total 603	C 380	N 98	O 122	S 3	0	0	0
1	E	81	Total 635	C 397	N 107	O 128	S 3	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	102	ALA	PRO	engineered mutation	UNP P03420
C	102	ALA	PRO	engineered mutation	UNP P03420
E	102	ALA	PRO	engineered mutation	UNP P03420

- Molecule 2 is a protein called Fusion glycoprotein F1 fused with Fibrin trimerization domain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	370	Total 2852	C 1805	N 469	O 560	S 18	0	0	0
2	D	370	Total 2852	C 1805	N 469	O 560	S 18	0	0	0
2	F	369	Total 2844	C 1799	N 468	O 559	S 18	0	0	0

There are 45 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	190	PHE	SER	engineered mutation	UNP P03420
B	207	LEU	VAL	engineered mutation	UNP P03420
B	379	VAL	ILE	engineered mutation	UNP P03420
B	447	VAL	MET	engineered mutation	UNP P03420

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	514	SER	-	linker	UNP P03420
B	515	ALA	-	linker	UNP P03420
B	516	ILE	-	linker	UNP P03420
B	517	GLY	-	linker	UNP P03420
B	539	LEU	PHE	variant	UNP P10104
B	545	GLY	-	expression tag	UNP P10104
B	546	GLY	-	expression tag	UNP P10104
B	547	LEU	-	expression tag	UNP P10104
B	548	VAL	-	expression tag	UNP P10104
B	549	PRO	-	expression tag	UNP P10104
B	550	ARG	-	expression tag	UNP P10104
D	190	PHE	SER	engineered mutation	UNP P03420
D	207	LEU	VAL	engineered mutation	UNP P03420
D	379	VAL	ILE	engineered mutation	UNP P03420
D	447	VAL	MET	engineered mutation	UNP P03420
D	514	SER	-	linker	UNP P03420
D	515	ALA	-	linker	UNP P03420
D	516	ILE	-	linker	UNP P03420
D	517	GLY	-	linker	UNP P03420
D	539	LEU	PHE	variant	UNP P10104
D	545	GLY	-	expression tag	UNP P10104
D	546	GLY	-	expression tag	UNP P10104
D	547	LEU	-	expression tag	UNP P10104
D	548	VAL	-	expression tag	UNP P10104
D	549	PRO	-	expression tag	UNP P10104
D	550	ARG	-	expression tag	UNP P10104
F	190	PHE	SER	engineered mutation	UNP P03420
F	207	LEU	VAL	engineered mutation	UNP P03420
F	379	VAL	ILE	engineered mutation	UNP P03420
F	447	VAL	MET	engineered mutation	UNP P03420
F	514	SER	-	linker	UNP P03420
F	515	ALA	-	linker	UNP P03420
F	516	ILE	-	linker	UNP P03420
F	517	GLY	-	linker	UNP P03420
F	539	LEU	PHE	variant	UNP P10104
F	545	GLY	-	expression tag	UNP P10104
F	546	GLY	-	expression tag	UNP P10104
F	547	LEU	-	expression tag	UNP P10104
F	548	VAL	-	expression tag	UNP P10104
F	549	PRO	-	expression tag	UNP P10104
F	550	ARG	-	expression tag	UNP P10104

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		
3	F	1	Total	O	S	0	0
			5	4	1		
3	F	1	Total	O	S	0	0
			5	4	1		
3	F	1	Total	O	S	0	0
			5	4	1		



- Molecule 4 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	A	21	Total 21	O 21	0	0
4	B	157	Total 157	O 157	0	0
4	C	11	Total 11	O 11	0	0
4	D	161	Total 161	O 161	0	0
4	E	20	Total 20	O 20	0	0
4	F	152	Total 152	O 152	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Fusion glycoprotein F2





- Molecule 1: Fusion glycoprotein F2




- Molecule 1: Fusion glycoprotein F2




- Molecule 2: Fusion glycoprotein F1 fused with Fibrin trimerization domain





- Molecule 2: Fusion glycoprotein F1 fused with Fibrin trimerization domain





• Molecule 2: Fusion glycoprotein F1 fused with Fibrin trimerization domain



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	170.75Å 170.75Å 163.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.86 – 2.40 37.86 – 2.40	Depositor EDS
% Data completeness (in resolution range)	95.5 (37.86-2.40) 95.5 (37.86-2.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.73 (at 2.39Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, $R_{free}$	0.187 , 0.214 0.188 , 0.214	Depositor DCC
$R_{free}$ test set	4591 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	48.8	Xtrriage
Anisotropy	0.122	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 54.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.012 for -h,-l,-k 0.005 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	11013	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.70% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.28	0/642	0.77	1/867 (0.1%)
1	C	0.30	0/610	0.81	2/824 (0.2%)
1	E	0.29	0/642	0.82	1/867 (0.1%)
2	B	0.30	0/2896	0.73	0/3929
2	D	0.29	0/2896	0.73	3/3929 (0.1%)
2	F	0.29	0/2888	0.76	3/3918 (0.1%)
All	All	0.29	0/10574	0.75	10/14334 (0.1%)

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	427	LYS	N-CA-C	6.61	118.48	111.28
1	C	100	THR	CA-C-N	6.56	126.32	119.76
1	C	100	THR	C-N-CA	6.56	126.32	119.76
2	F	209	LYS	CA-C-N	5.74	132.03	121.70
2	F	209	LYS	C-N-CA	5.74	132.03	121.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	635	0	642	7	0
1	C	603	0	612	9	0
1	E	635	0	642	19	0
2	B	2852	0	2884	38	0
2	D	2852	0	2884	44	0
2	F	2844	0	2873	51	0
3	B	25	0	0	2	0
3	D	25	0	0	0	0
3	F	20	0	0	0	0
4	A	21	0	0	2	0
4	B	157	0	0	7	1
4	C	11	0	0	0	0
4	D	161	0	0	10	0
4	E	20	0	0	2	0
4	F	152	0	0	6	1
All	All	11013	0	10537	132	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 132 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:216:ASN:ND2	2:F:218:GLU:OE2	1.95	1.00
2:F:264:MET:HE1	2:F:303:LEU:HD12	1.52	0.92
2:D:184:GLY:O	4:D:854:HOH:O	1.95	0.82
2:B:246:PRO:HB3	2:B:283:GLN:HA	1.62	0.81
2:F:246:PRO:HB3	2:F:283:GLN:HA	1.62	0.81

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:779:HOH:O	4:F:762:HOH:O[7_555]	2.12	0.08

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	79/82 (96%)	72 (91%)	6 (8%)	1 (1%)	9	14
1	C	75/82 (92%)	69 (92%)	6 (8%)	0	100	100
1	E	79/82 (96%)	71 (90%)	7 (9%)	1 (1%)	9	14
2	B	368/414 (89%)	352 (96%)	15 (4%)	1 (0%)	36	50
2	D	368/414 (89%)	354 (96%)	12 (3%)	2 (0%)	24	37
2	F	367/414 (89%)	353 (96%)	14 (4%)	0	100	100
All	All	1336/1488 (90%)	1271 (95%)	60 (4%)	5 (0%)	30	43

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	66	GLU
2	D	216	ASN
1	E	103	THR
2	B	295	GLU
2	D	294	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	72/73 (99%)	71 (99%)	1 (1%)	59	79
1	C	69/73 (94%)	67 (97%)	2 (3%)	37	60
1	E	72/73 (99%)	71 (99%)	1 (1%)	59	79

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	338/373 (91%)	336 (99%)	2 (1%)	78	89
2	D	338/373 (91%)	336 (99%)	2 (1%)	78	89
2	F	337/373 (90%)	335 (99%)	2 (1%)	78	89
All	All	1226/1338 (92%)	1216 (99%)	10 (1%)	73	86

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	67	ASN
2	F	216	ASN
2	F	220	VAL
1	C	64	ILE
1	C	69	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	276	ASN
2	D	494	GLN
2	F	494	GLN
1	E	67	ASN
2	B	462	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	B	601	-	4,4,4	0.24	0	6,6,6	0.05	0
3	SO4	F	601	-	4,4,4	0.23	0	6,6,6	0.08	0
3	SO4	D	603	-	4,4,4	0.24	0	6,6,6	0.07	0
3	SO4	D	605	-	4,4,4	0.24	0	6,6,6	0.08	0
3	SO4	D	602	-	4,4,4	0.23	0	6,6,6	0.08	0
3	SO4	B	604	-	4,4,4	0.24	0	6,6,6	0.19	0
3	SO4	D	601	-	4,4,4	0.23	0	6,6,6	0.07	0
3	SO4	B	603	-	4,4,4	0.24	0	6,6,6	0.07	0
3	SO4	B	605	-	4,4,4	0.24	0	6,6,6	0.13	0
3	SO4	B	602	-	4,4,4	0.24	0	6,6,6	0.08	0
3	SO4	D	604	-	4,4,4	0.24	0	6,6,6	0.08	0
3	SO4	F	603	-	4,4,4	0.23	0	6,6,6	0.08	0
3	SO4	F	602	-	4,4,4	0.23	0	6,6,6	0.08	0
3	SO4	F	604	-	4,4,4	0.24	0	6,6,6	0.08	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	604	SO4	1	0
3	B	605	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	81/82 (98%)	0.60	11 (13%) 7 5	24, 56, 139, 171	0
1	C	77/82 (93%)	0.65	12 (15%) 5 4	23, 58, 157, 173	0
1	E	81/82 (98%)	0.71	15 (18%) 3 3	23, 52, 130, 151	0
2	B	370/414 (89%)	0.20	29 (7%) 19 15	22, 42, 108, 156	0
2	D	370/414 (89%)	0.34	30 (8%) 18 14	21, 46, 124, 152	0
2	F	369/414 (89%)	0.19	24 (6%) 25 21	22, 43, 97, 143	0
All	All	1348/1488 (90%)	0.32	121 (8%) 15 12	21, 46, 121, 173	0

The worst 5 of 121 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	505	PHE	8.2
1	A	103	THR	5.9
1	E	100	THR	5.9
1	C	103	THR	5.7
1	A	102	ALA	5.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	D	605	5/5	0.51	0.16	138,138,142,144	0
3	SO4	B	605	5/5	0.54	0.13	144,151,158,161	0
3	SO4	D	604	5/5	0.69	0.12	92,114,119,122	0
3	SO4	F	604	5/5	0.80	0.10	93,115,119,122	0
3	SO4	F	603	5/5	0.82	0.11	126,131,132,133	0
3	SO4	D	603	5/5	0.82	0.12	113,116,116,119	0
3	SO4	D	602	5/5	0.86	0.11	76,84,97,104	0
3	SO4	F	602	5/5	0.86	0.11	90,100,103,109	0
3	SO4	B	604	5/5	0.86	0.12	62,68,100,104	0
3	SO4	B	603	5/5	0.86	0.12	104,104,109,109	0
3	SO4	F	601	5/5	0.88	0.11	79,90,99,102	0
3	SO4	B	602	5/5	0.93	0.22	72,73,77,97	0
3	SO4	D	601	5/5	0.97	0.14	48,51,56,63	0
3	SO4	B	601	5/5	0.98	0.06	36,38,42,43	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.