



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 01:48 PM UTC

PDB ID : 7MMO / pdb_00007mmo
Title : LY-CoV1404 neutralizing antibody against SARS-CoV-2
Authors : Hendle, J.; Pustilnik, A.; Sauder, J.M.; Coleman, K.A.; Boyles, J.S.; Dickinson, C.D.
Deposited on : 2021-04-30
Resolution : 2.43 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6062 (2.44-2.40)
Clashscore	190562	6562 (2.44-2.40)
Sidechain outliers	187428	6482 (2.44-2.40)
RSRZ outliers	180081	6066 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	223	5% 91% 5%
1	D	223	4% 90% 7%
2	B	215	% 94% 5%
2	E	215	5% 94% 6%
3	C	205	18% 90% 5%
3	F	205	45% 80% 5% 15%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9153 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LY-CoV1404 Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	212	Total	C	N	O	S	0	1	0
			1580	1012	259	302	7			
1	D	208	Total	C	N	O	S	0	1	0
			1558	998	254	299	7			

- Molecule 2 is a protein called LY-CoV1404 Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	214	Total	C	N	O	S	0	1	0
			1585	981	262	335	7			
2	E	214	Total	C	N	O	S	0	1	0
			1584	983	262	332	7			

- Molecule 3 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	194	Total	C	N	O	S	0	2	0
			1486	951	252	275	8			
3	F	175	Total	C	N	O	S	0	0	0
			1273	810	221	236	6			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	528	HIS	-	expression tag	UNP P0DTC2
C	529	HIS	-	expression tag	UNP P0DTC2
C	530	HIS	-	expression tag	UNP P0DTC2
C	531	HIS	-	expression tag	UNP P0DTC2
C	532	HIS	-	expression tag	UNP P0DTC2
C	533	HIS	-	expression tag	UNP P0DTC2
F	528	HIS	-	expression tag	UNP P0DTC2
F	529	HIS	-	expression tag	UNP P0DTC2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	530	HIS	-	expression tag	UNP P0DTC2
F	531	HIS	-	expression tag	UNP P0DTC2
F	532	HIS	-	expression tag	UNP P0DTC2
F	533	HIS	-	expression tag	UNP P0DTC2

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	N	O	
			14	8	1	5	
						0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	12	Total	O		
			12	12	0	0
5	B	24	Total	O		
			24	24	0	0
5	C	7	Total	O		
			8	8	0	1
5	D	8	Total	O		
			8	8	0	0
5	E	19	Total	O		
			19	19	0	0
5	F	2	Total	O		
			2	2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: LY-CoV1404 Fab heavy chain




- Molecule 1: LY-CoV1404 Fab heavy chain




- Molecule 2: LY-CoV1404 Fab light chain




- Molecule 2: LY-CoV1404 Fab light chain





- Molecule 3: Spike protein S1





● Molecule 3: Spike protein S1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	73.09Å 107.69Å 190.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	93.75 – 2.43 93.75 – 2.43	Depositor EDS
% Data completeness (in resolution range)	99.5 (93.75-2.43) 99.5 (93.75-2.43)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.21 (at 2.42Å)	Xtriage
Refinement program	BUSTER 2.11.7 (18-SEP-2020)	Depositor
R, R_{free}	0.226 , 0.253 (Not available) , 0.267	Depositor DCC
R_{free} test set	2788 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å ²)	55.6	Xtriage
Anisotropy	0.003	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 46.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	9153	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 23.23 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.8631e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PCA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.68	0/1617	0.98	2/2219 (0.1%)
1	D	0.67	0/1593	0.95	0/2183
2	B	0.69	0/1619	0.94	1/2203 (0.0%)
2	E	0.68	0/1618	0.95	2/2201 (0.1%)
3	C	0.65	0/1534	0.91	2/2093 (0.1%)
3	F	0.59	0/1306	0.88	0/1784
All	All	0.66	0/9287	0.94	7/12683 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	98	SER	N-CA-C	5.80	117.60	108.96
2	B	98	SER	N-CA-C	5.34	117.19	109.07
3	C	365	TYR	CA-C-N	5.20	127.77	120.28
3	C	365	TYR	C-N-CA	5.20	127.77	120.28
1	A	102	ILE	CA-C-N	5.09	127.05	120.44
1	A	102	ILE	C-N-CA	5.09	127.05	120.44
2	E	118	SER	N-CA-C	-5.03	100.97	108.96

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1580	0	1522	2	0
1	D	1558	0	1515	2	0
2	B	1585	0	1484	4	0
2	E	1584	0	1479	4	0
3	C	1486	0	1346	7	0
3	F	1273	0	1074	7	0
4	C	14	0	13	0	0
5	A	12	0	0	0	0
5	B	24	0	0	0	0
5	C	8	0	0	0	0
5	D	8	0	0	0	0
5	E	19	0	0	0	0
5	F	2	0	0	0	0
All	All	9153	0	8433	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:370:ASN:HA	3:F:384:PRO:HG2	1.34	1.04
3:C:370:ASN:HA	3:F:384:PRO:CG	2.23	0.57
3:C:374:PHE:HA	3:C:436:TRP:HB3	1.89	0.54
3:F:433:VAL:HG22	3:F:512:VAL:HG13	1.88	0.54
3:F:455:LEU:HD22	3:F:493:GLN:HG3	1.90	0.53
2:E:19:ILE:HD11	2:E:78:ILE:HD12	1.90	0.53
2:B:124:PRO:HD3	2:B:136:VAL:HG22	1.93	0.51
2:E:124:PRO:HD3	2:E:136:VAL:HG22	1.93	0.49
3:C:430:THR:HG22	3:C:515:PHE:HB2	1.95	0.49
2:B:19:ILE:HD11	2:B:78:ILE:HD12	1.96	0.48
3:C:431:GLY:HA3	3:C:513:LEU:O	2.14	0.47
2:B:167:VAL:HG22	2:B:179:LEU:HD12	1.97	0.45
1:D:129:PRO:HD3	1:D:215:LYS:HE2	1.99	0.45
2:E:167:VAL:HG22	2:E:179:LEU:HD12	1.99	0.45
1:A:129:PRO:HD3	1:A:215:LYS:HE2	1.97	0.44
3:F:374:PHE:HA	3:F:436:TRP:HB3	2.02	0.42
2:E:86:GLU:HG3	2:E:107:LEU:O	2.20	0.42
2:B:86:GLU:HG3	2:B:107:LEU:O	2.20	0.41
3:F:401:VAL:HG22	3:F:509:ARG:HG2	2.02	0.41
3:C:392:PHE:CE2	3:C:515:PHE:HB3	2.55	0.41
1:A:22:CYS:O	1:A:79:GLN:HA	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:22:CYS:O	1:D:79:GLN:HA	2.21	0.40
3:C:370:ASN:CA	3:F:384:PRO:HG2	2.26	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	173/198 (87%)	170 (98%)	3 (2%)	53	73
1	D	174/198 (88%)	170 (98%)	4 (2%)	44	64
2	B	176/184 (96%)	174 (99%)	2 (1%)	65	81
2	E	173/184 (94%)	171 (99%)	2 (1%)	63	79
3	C	149/178 (84%)	149 (100%)	0	100	100
3	F	112/178 (63%)	111 (99%)	1 (1%)	70	84
All	All	957/1120 (85%)	945 (99%)	12 (1%)	61	78

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	87	ILE
1	A	149	LYS
1	A	199	THR
2	B	32	ASP
2	B	156	ASN
1	D	58	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	87	ILE
1	D	149	LYS
1	D	167	SER
2	E	32	ASP
2	E	215	GLU
3	F	424	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	GLN
2	B	34	ASN
2	B	40	GLN
2	B	41	GLN
2	B	142	ASN
2	E	34	ASN
2	E	142	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PCA	A	1	1	7,8,9	1.21	1 (14%)	9,10,12	1.28	0
1	PCA	D	1	1	7,8,9	0.75	0	9,10,12	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	A	1	1	-	0/0/11/13	0/1/1/1
1	PCA	D	1	1	-	0/0/11/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	PCA	CD-N	2.19	1.39	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	C	601	3	14,14,15	0.27	0	17,19,21	0.61	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	601	3	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	601	NAG	O5-C1-C2	-2.40	107.58	111.29

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	211/223 (94%)	0.48	12 (5%)	29 25	28, 52, 70, 89	1 (0%)
1	D	207/223 (92%)	0.62	10 (4%)	35 32	29, 54, 71, 85	1 (0%)
2	B	214/215 (99%)	0.36	3 (1%)	73 70	28, 48, 60, 74	1 (0%)
2	E	214/215 (99%)	0.44	10 (4%)	36 32	29, 50, 63, 79	1 (0%)
3	C	194/205 (94%)	1.13	36 (18%)	3 2	29, 64, 103, 111	2 (1%)
3	F	175/205 (85%)	2.24	92 (52%)	0 0	57, 92, 119, 121	0
All	All	1215/1286 (94%)	0.83	163 (13%)	7 5	28, 54, 105, 121	6 (0%)

All (163) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	389	ASP	6.0
3	F	481	ASN	5.9
3	C	373	SER	5.5
3	C	371	SER	5.4
3	C	366	SER	5.1
3	F	382	VAL	5.0
3	F	391	CYS	4.9
3	F	491	PRO	4.8
3	F	341	VAL	4.7
3	F	383	SER	4.7
3	F	520	ALA	4.7
3	F	517	LEU	4.7
3	F	363	ALA	4.6
3	F	334	ASN	4.6
3	F	342	PHE	4.5
3	C	369	TYR	4.5
3	C	481	ASN	4.5
1	D	192	SER	4.5
3	C	482	GLY	4.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	F	384	PRO	4.3
3	F	392	PHE	4.2
3	F	505	TYR	4.2
3	F	390	LEU	4.2
3	F	372	ALA	4.2
3	F	430	THR	4.0
3	C	364	ASP	4.0
3	F	346	ARG	4.0
3	F	373	SER	4.0
3	C	372	ALA	4.0
3	F	455	LEU	4.0
3	C	367	VAL	4.0
3	F	518	LEU	3.9
3	F	377	PHE	3.8
2	B	216	CYS	3.8
3	C	527	PRO	3.8
3	F	515	PHE	3.8
3	F	374	PHE	3.7
3	F	337	PRO	3.7
3	F	345	THR	3.6
3	F	347	PHE	3.6
1	D	219	HIS	3.5
1	D	141	THR	3.5
3	F	395	VAL	3.5
3	F	381	GLY	3.5
3	F	397	ALA	3.5
2	E	27	SER	3.4
3	F	338	PHE	3.4
3	C	368	LEU	3.4
3	F	351	TYR	3.4
3	F	483	VAL	3.4
3	F	472	ILE	3.4
3	F	396	TYR	3.3
3	F	398	ASP	3.3
3	F	362	VAL	3.3
3	F	497	PHE	3.3
3	F	405	ASP	3.2
3	F	393	THR	3.2
1	A	219	HIS	3.2
3	F	519	HIS	3.2
3	C	478	THR	3.2
3	F	400	PHE	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	496	GLY	3.1
3	F	380	TYR	3.1
1	A	197	THR	3.1
1	A	192	SER	3.1
3	F	356	LYS	3.1
3	F	492	LEU	3.1
1	A	141	THR	3.0
3	F	477	SER	3.0
2	E	26	THR	3.0
3	F	335	LEU	3.0
3	F	487	ASN	3.0
1	A	193	SER	3.0
3	F	478	THR	3.0
3	C	381	GLY	3.0
1	D	134	SER	2.9
3	C	370	ASN	2.9
3	F	358	ILE	2.9
3	C	518	LEU	2.9
1	A	194	SER	2.9
3	F	339	GLY	2.9
3	F	399	SER	2.8
3	F	359	SER	2.8
3	F	511	VAL	2.8
3	C	517	LEU	2.8
3	F	355	ARG	2.8
3	C	475	ALA	2.8
3	F	495	TYR	2.8
3	F	375	SER	2.8
3	F	461	LEU	2.8
3	F	403	ARG	2.8
2	E	29	ASP	2.8
3	C	345	THR	2.7
2	B	56	ASP	2.7
3	F	401	VAL	2.7
3	C	455	LEU	2.7
1	A	195	LEU	2.6
3	F	340	GLU	2.6
1	D	218	HIS	2.6
3	C	505	TYR	2.6
3	C	382	VAL	2.6
1	A	75	THR	2.6
3	F	357	ARG	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	F	486	PHE	2.6
1	A	218	HIS	2.6
1	D	199	THR	2.6
3	C	377	PHE	2.5
3	F	471	GLU	2.5
3	C	334	ASN	2.5
3	F	460	ASN	2.5
3	F	490	PHE	2.5
3	F	514	SER	2.5
3	C	450	ASN	2.5
3	F	502	GLY	2.4
3	F	453	TYR	2.4
3	C	486	PHE	2.4
3	F	464	PHE	2.4
1	D	133	CYS	2.4
3	F	463	PRO	2.4
2	E	54	VAL	2.4
3	C	483	VAL	2.4
2	B	204	THR	2.4
1	A	133	CYS	2.4
3	F	361	CYS	2.4
1	D	198	LYS	2.4
3	C	384	PRO	2.3
3	F	350	VAL	2.3
3	F	484	GLU	2.3
3	C	477	SER	2.3
2	E	216	CYS	2.3
3	F	479	PRO	2.3
3	F	376	THR	2.3
3	F	353	TRP	2.3
3	C	374	PHE	2.3
3	F	429	PHE	2.3
2	E	28	SER	2.2
3	F	480	CYS	2.2
3	F	510	VAL	2.2
3	C	335	LEU	2.2
1	A	189	THR	2.2
3	F	482	GLY	2.2
3	F	440	ASN	2.1
3	F	513	LEU	2.1
2	E	70	SER	2.1
3	F	408	ARG	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	491	PRO	2.1
3	C	522	ALA	2.1
2	E	24	THR	2.1
3	C	365	TYR	2.1
3	F	458	LYS	2.1
2	E	34	ASN	2.1
3	C	473	TYR	2.1
1	D	18	LEU	2.1
3	F	468	ILE	2.1
3	F	421	TYR	2.1
3	F	509	ARG	2.0
3	F	425	LEU	2.0
3	F	476	GLY	2.0
2	E	133	THR	2.0
1	A	198	LYS	2.0
3	C	376	THR	2.0
3	F	427	ASP	2.0
1	D	28	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PCA	A	1	8/9	0.80	0.15	64,66,68,73	0
1	PCA	D	1	8/9	0.87	0.11	62,63,65,67	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	C	601	14/15	0.82	0.27	71,72,72,72	14

6.5 Other polymers [i](#)

There are no such residues in this entry.