



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 06:47 PM UTC

PDB ID : 4MN4 / pdb_00004mn4
Title : Structural Basis for the MukB-topoisomerase IV Interaction
Authors : Vos, S.M.; Stewart, N.K.; Oakley, M.G.; Berger, J.M.
Deposited on : 2013-09-09
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	259	
1	B	259	
2	C	163	
2	D	163	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6542 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA topoisomerase 4 subunit A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	246	Total	C	N	O	S	Se	0	3	0
			1875	1179	334	351	1	10			
1	B	245	Total	C	N	O	S	Se	0	7	0
			1881	1182	332	355	1	11			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	494	SER	-	expression tag	UNP P0AFI2
A	495	ASN	-	expression tag	UNP P0AFI2
A	496	ALA	-	expression tag	UNP P0AFI2
B	494	SER	-	expression tag	UNP P0AFI2
B	495	ASN	-	expression tag	UNP P0AFI2
B	496	ALA	-	expression tag	UNP P0AFI2

- Molecule 2 is a protein called Chromosome partition protein MukB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	159	Total	C	N	O	S	0	4	0
			1287	804	222	260	1			
2	D	154	Total	C	N	O	S	0	4	0
			1248	782	213	252	1			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	642	SER	-	expression tag	UNP P22523
C	643	ASN	-	expression tag	UNP P22523
C	644	ALA	-	expression tag	UNP P22523
D	642	SER	-	expression tag	UNP P22523
D	643	ASN	-	expression tag	UNP P22523
D	644	ALA	-	expression tag	UNP P22523


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	71	Total 71	O 71	0	0
3	C	58	Total 58	O 58	0	0
3	D	46	Total 46	O 46	0	0
3	B	76	Total 76	O 76	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA topoisomerase 4 subunit A




- Molecule 1: DNA topoisomerase 4 subunit A



- Molecule 2: Chromosome partition protein MukB



- Molecule 2: Chromosome partition protein MukB





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.77Å 103.74Å 186.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.97 – 2.30 49.97 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.1 (49.97-2.30) 97.7 (49.97-2.30)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.06 (at 2.29Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.204 , 0.246 0.204 , 0.244	Depositor DCC
R_{free} test set	2170 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	35.3	Xtriage
Anisotropy	0.123	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 35.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6542	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/1904	0.71	0/2560
1	B	0.39	0/1925	0.70	2/2589 (0.1%)
2	C	0.52	2/1322 (0.2%)	0.72	2/1790 (0.1%)
2	D	0.26	0/1283	0.67	0/1737
All	All	0.39	2/6434 (0.0%)	0.70	4/8676 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	692[A]	ASP	CA-C	8.32	1.64	1.52
2	C	692[B]	ASP	CA-C	8.32	1.64	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	692[A]	ASP	N-CA-C	6.30	120.57	113.01
2	C	692[B]	ASP	N-CA-C	6.30	120.57	113.01
1	B	692	LEU	CA-C-N	5.26	123.50	119.66
1	B	692	LEU	C-N-CA	5.26	123.50	119.66

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1875	0	1940	36	0
1	B	1881	0	1945	30	0
2	C	1287	0	1239	18	0
2	D	1248	0	1202	23	0
3	A	71	0	0	0	0
3	B	76	0	0	0	0
3	C	58	0	0	1	0
3	D	46	0	0	0	0
All	All	6542	0	6326	107	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (107) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:652:MSE:HB2	1:B:728[A]:MSE:HE3	1.51	0.91
1:A:538:LYS:H	1:A:541:GLN:HE21	1.19	0.85
2:C:803:THR:O	2:C:804:LEU:HB2	1.83	0.79
2:C:751:VAL:HG22	2:C:760:VAL:HG22	1.65	0.78
1:A:720:GLU:HB2	1:A:723:ARG:HE	1.51	0.74
1:B:724:ARG:HB3	1:B:725:GLY:HA2	1.70	0.74
1:A:699:THR:HB	1:A:738:GLU:HB2	1.71	0.72
1:A:570:LEU:HD23	1:A:582:VAL:HG21	1.75	0.68
2:D:756:LYS:HA	2:D:772:PHE:CD2	2.31	0.66
2:D:795:GLU:HG2	2:D:796:VAL:N	2.08	0.66
1:B:654:MSE:HE2	1:B:718:THR:HG22	1.77	0.66
1:A:710:ARG:N	1:A:713:GLU:OE1	2.28	0.65
2:D:717:LEU:HG	2:D:738:GLU:HB2	1.79	0.64
1:A:664:SER:O	1:A:665:LYS:HG3	1.97	0.63
1:A:611:LEU:HG	1:A:621:LEU:HD22	1.81	0.63
1:A:538:LYS:N	1:A:541:GLN:HE21	1.94	0.63
1:B:652:MSE:HB2	1:B:728[A]:MSE:CE	2.26	0.63
2:C:654:ASN:O	2:C:658:GLU:HG2	1.99	0.62
1:B:654:MSE:HE2	1:B:718:THR:CG2	2.30	0.61
1:B:662:GLN:C	1:B:663:LEU:HD12	2.29	0.58
1:B:507[A]:MSE:SE	1:B:526:LYS:HE3	2.55	0.57
1:A:664:SER:C	1:A:665:LYS:HG3	2.29	0.57
2:D:680:ARG:HD3	2:D:723:HIS:CD2	2.39	0.57
1:A:654:MSE:HE2	1:A:718:THR:OG1	2.03	0.57
2:C:646:ARG:HG3	2:C:804:LEU:HD13	1.87	0.57
1:B:724:ARG:HB3	1:B:725:GLY:CA	2.35	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:776:PRO:HG2	2:D:783:ARG:HH21	1.71	0.56
1:A:720:GLU:HB3	1:A:723:ARG:HH21	1.70	0.56
1:A:538:LYS:H	1:A:541:GLN:NE2	1.98	0.55
2:D:645:GLU:HG2	2:D:646:ARG:H	1.72	0.55
1:A:622:ILE:HA	1:A:666:GLY:O	2.08	0.54
2:D:698:ALA:HB3	2:D:699:PRO:HD3	1.88	0.54
1:B:644:LEU:HD11	1:B:652:MSE:HE3	1.88	0.54
1:B:709:LEU:HD12	1:B:709:LEU:N	2.22	0.54
2:D:652:ARG:HG3	2:D:797:LEU:HD11	1.89	0.54
1:A:520:ALA:HB3	1:A:521:PRO:HD3	1.90	0.54
2:C:751:VAL:HG13	2:C:758:VAL:HG13	1.91	0.53
1:B:511:ARG:HD2	1:B:563:ALA:HA	1.90	0.53
1:A:720:GLU:CB	1:A:723:ARG:HH21	2.22	0.53
1:A:644:LEU:HD11	1:A:652:MSE:HE2	1.91	0.52
1:A:541:GLN:HG3	1:A:587:MSE:HE2	1.91	0.52
1:B:611:LEU:HG	1:B:621:LEU:HD22	1.90	0.52
1:B:646:ILE:HG12	1:B:652:MSE:HG2	1.91	0.52
2:C:717:LEU:HG	2:C:738:GLU:HB2	1.90	0.52
2:C:756:LYS:HB3	2:C:772:PHE:CD2	2.44	0.51
1:B:605:VAL:O	1:B:662:GLN:HG3	2.11	0.51
2:D:677:LEU:HD23	2:D:680:ARG:NH2	2.25	0.51
1:B:520:ALA:HB3	1:B:521:PRO:HD3	1.92	0.51
1:A:570:LEU:HG	1:A:574:LEU:HD12	1.92	0.51
2:C:720:VAL:HG11	2:C:736:LEU:HD13	1.92	0.50
1:B:607:THR:HG22	1:B:609:ASN:H	1.77	0.50
2:C:698:ALA:HB3	2:C:699:PRO:HD3	1.94	0.50
1:B:660:LEU:HD12	1:B:661:PRO:HD2	1.95	0.49
2:C:749:PHE:HB3	2:C:751:VAL:HG23	1.95	0.49
2:D:662:ARG:O	2:D:665:GLN:HB3	2.12	0.49
2:D:661:GLU:O	2:D:665:GLN:HB2	2.14	0.48
2:D:730:CYS:HB2	2:D:731:PRO:HD2	1.96	0.48
1:A:710:ARG:HB2	1:A:712:GLU:HG2	1.95	0.48
2:C:690:TYR:O	2:C:693:VAL:HG22	2.13	0.47
1:B:643:LEU:C	1:B:643:LEU:HD23	2.39	0.47
2:D:796:VAL:O	2:D:799:GLU:HG3	2.14	0.47
1:B:726:THR:O	1:B:728[A]:MSE:HE2	2.15	0.47
1:B:607:THR:HG22	1:B:609:ASN:N	2.30	0.46
1:B:707:ILE:N	1:B:707:ILE:HD12	2.30	0.46
1:A:525:TYR:OH	1:A:531:PHE:HB2	2.16	0.45
1:A:643:LEU:C	1:A:643:LEU:HD23	2.41	0.45
2:C:730:CYS:HB2	2:C:731:PRO:HD2	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:595:LEU:HB3	1:A:634:VAL:HB	1.98	0.45
1:A:599:ASP:HA	1:A:686[B]:LEU:HB2	1.97	0.45
2:D:783:ARG:O	2:D:787:ILE:HG13	2.16	0.45
2:D:693[A]:VAL:CG2	2:D:697:ASP:HB2	2.47	0.45
1:B:607:THR:CG2	1:B:609:ASN:H	2.31	0.44
1:B:714:LEU:O	1:B:718:THR:HG23	2.17	0.44
2:D:666:PRO:HA	2:D:667:GLY:HA2	1.74	0.44
2:D:780:ARG:HA	2:D:783:ARG:NH1	2.33	0.43
1:A:638:ASP:HB3	1:A:641:ASP:CG	2.43	0.43
2:D:683:GLY:HA3	2:D:714:VAL:HG12	1.99	0.43
1:A:638:ASP:HB3	1:A:641:ASP:OD1	2.19	0.43
2:C:780:ARG:HD2	3:C:936:HOH:O	2.18	0.43
1:B:644:LEU:HD11	1:B:652:MSE:CE	2.48	0.43
2:C:704:LEU:CD1	2:C:751:VAL:HG21	2.48	0.43
1:A:570:LEU:HD12	1:A:570:LEU:HA	1.89	0.42
2:C:684:VAL:CG2	2:C:741:PRO:HB3	2.50	0.42
1:B:693:PRO:HG2	1:B:741:SER:HB3	2.02	0.42
1:B:700:ILE:HG21	1:B:731:LEU:HD11	2.02	0.42
1:B:724:ARG:CB	1:B:725:GLY:HA2	2.38	0.42
2:D:720:VAL:HG11	2:D:736:LEU:HD13	2.02	0.42
1:A:509:TRP:CZ2	1:A:569:PRO:HG3	2.54	0.42
1:A:700:ILE:HG21	1:A:731:LEU:HD11	2.02	0.42
1:A:660:LEU:HD12	1:A:661:PRO:HD2	2.01	0.42
1:A:568:GLU:OE2	1:A:573:LYS:HE2	2.19	0.42
2:C:680[B]:ARG:NH1	2:C:727:LEU:HD13	2.34	0.41
2:C:652:ARG:O	2:C:656:VAL:HG23	2.20	0.41
2:D:714:VAL:O	2:D:738:GLU:HA	2.20	0.41
1:B:675:PRO:HB2	1:B:678:GLU:HG3	2.03	0.41
1:B:647:THR:HG23	1:B:672:ILE:HD13	2.02	0.41
1:A:520:ALA:O	1:A:523:LEU:HB2	2.21	0.41
2:C:695:LEU:HD11	2:C:786:ARG:HG2	2.02	0.41
2:D:652:ARG:HG3	2:D:797:LEU:CD1	2.50	0.41
2:D:793:GLU:O	2:D:797:LEU:HG	2.20	0.41
1:B:547:ASP:OD1	1:B:547:ASP:C	2.64	0.41
1:A:508:GLY:O	1:A:570:LEU:HB2	2.21	0.40
2:D:690:TYR:O	2:D:693[B]:VAL:HG22	2.22	0.40
1:A:523:LEU:HD12	1:A:523:LEU:HA	1.91	0.40
1:A:621:LEU:O	1:A:666:GLY:HA2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/259 (95%)	242 (98%)	4 (2%)	1 (0%)	30	38
1	B	250/259 (96%)	242 (97%)	8 (3%)	0	100	100
2	C	159/163 (98%)	157 (99%)	2 (1%)	0	100	100
2	D	154/163 (94%)	149 (97%)	5 (3%)	0	100	100
All	All	810/844 (96%)	790 (98%)	19 (2%)	1 (0%)	48	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	666	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	205/203 (101%)	199 (97%)	6 (3%)	37	55
1	B	208/203 (102%)	206 (99%)	2 (1%)	68	82
2	C	141/140 (101%)	139 (99%)	2 (1%)	59	76
2	D	138/140 (99%)	133 (96%)	5 (4%)	31	47
All	All	692/686 (101%)	677 (98%)	15 (2%)	45	65

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	564	ARG
1	A	665	LYS
1	A	712	GLU
1	A	716	LYS
1	A	717	VAL
1	A	723	ARG
2	C	652	ARG
2	C	758	VAL
2	D	680	ARG
2	D	685	LEU
2	D	725	GLU
2	D	795	GLU
2	D	798	SER
1	B	607	THR
1	B	717	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	541	GLN
1	A	566	GLN
1	A	584	HIS
1	A	673	ASN
1	A	695	GLN
1	A	732	GLN
2	C	665	GLN
2	D	665	GLN
2	D	672	GLN
2	D	723	HIS
1	B	566	GLN
1	B	688	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	237/259 (91%)	0.12	6 (2%) 58 60	14, 32, 68, 78	2 (0%)
1	B	236/259 (91%)	-0.07	5 (2%) 63 65	14, 30, 55, 81	5 (2%)
2	C	159/163 (97%)	0.15	6 (3%) 44 46	16, 32, 66, 96	4 (2%)
2	D	154/163 (94%)	0.58	20 (12%) 7 8	15, 37, 93, 106	4 (2%)
All	All	786/844 (93%)	0.16	37 (4%) 36 38	14, 32, 72, 106	15 (1%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	667	GLY	6.0
2	D	666	PRO	4.6
2	C	667	GLY	4.5
2	D	801	PHE	3.8
2	D	646	ARG	3.7
2	D	795	GLU	3.5
2	C	755	GLU	3.4
1	B	524[A]	ASN	3.3
2	C	666	PRO	3.3
2	D	650	GLY	3.2
2	D	653	LYS	3.2
1	A	497	SER	3.1
2	D	656	VAL	3.0
2	D	797	LEU	2.8
1	A	742	PRO	2.7
2	D	649	VAL	2.7
2	D	658	GLU	2.6
1	A	523	LEU	2.6
1	A	515	GLY	2.6
2	C	658	GLU	2.5
2	D	672	GLN	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	564	ARG	2.4
2	D	655	ALA	2.4
1	A	695	GLN	2.4
2	D	755	GLU	2.4
1	B	517	ASP	2.3
1	B	558	ILE	2.3
2	D	671	ASP	2.3
2	D	798	SER	2.3
1	B	518	ILE	2.2
2	D	746	ASP	2.1
2	D	796	VAL	2.1
1	A	518	ILE	2.1
2	C	672	GLN	2.0
2	D	748	VAL	2.0
2	D	800	ARG	2.0
2	C	756	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.