



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:51 PM UTC

PDB ID : 5MN2 / pdb_00005mn2
Title : Cocrystal structure of Fc gamma receptor IIIa interacting with Affimer G3, a specific binding protein which blocks IgG binding to the receptor.
Authors : Robinson, J.I.; Owen, R.L.; Tomlinson, D.C.; Baxter, E.W.; Nettleship, J.E.; Waterhouse, M.P.; Harris, S.A.; Owens, R.J.; McPherson, M.J.; Morgan, A.W.; Tiede, C.; Goldman, A.; Thomsen, M.
Deposited on : 2016-12-12
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1596 (2.36-2.36)
Clashscore	190562	1663 (2.36-2.36)
Ramachandran outliers	187476	1646 (2.36-2.36)
Sidechain outliers	187428	1646 (2.36-2.36)
RSRZ outliers	180081	1598 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	175	3% 97% ..
1	B	175	21% 94% ..
2	C	101	4% 87% 13%
2	D	101	4% 89% 11%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 8222 atoms, of which 3877 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Low affinity immunoglobulin gamma Fc region receptor III-A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	172	Total	C	H	N	O	S	0	7	0
			2719	902	1303	241	269	4			
1	B	168	Total	C	H	N	O	S	0	1	0
			2354	792	1100	212	246	4			

- Molecule 2 is a protein called Affimer G3.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	C	88	Total	C	H	N	O	S	0	0	0
			1395	454	690	119	129	3			
2	D	90	Total	C	H	N	O	S	0	0	0
			1346	448	649	116	130	3			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
3	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
3	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
3	A	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
3	A	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
3	B	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
3	B	1	Total	C	H	N	O	0	0
			27	8	13	1	5		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			17	4	10	3		
4	B	1	Total	C	H	O	0	0
			9	2	5	2		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 6 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	D	1	Total	C	H	O	0	0
			29	8	17	4		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	77	Total 77	O 77	0	2
7	B	10	Total 10	O 10	0	0
7	C	32	Total 32	O 32	0	0
7	D	27	Total 27	O 27	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Low affinity immunoglobulin gamma Fc region receptor III-A





- Molecule 1: Low affinity immunoglobulin gamma Fc region receptor III-A



- Molecule 2: Affimer G3



- Molecule 2: Affimer G3



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	64.98Å 59.92Å 100.00Å 90.00° 102.09° 90.00°	Depositor
Resolution (Å)	43.39 – 2.35 43.39 – 2.35	Depositor EDS
% Data completeness (in resolution range)	95.5 (43.39-2.35) 95.5 (43.39-2.35)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 2.34Å)	Xtriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, R_{free}	0.205 , 0.244 0.211 , 0.248	Depositor DCC
R_{free} test set	1511 reflections (4.77%)	wwPDB-VP
Wilson B-factor (Å ²)	58.5	Xtriage
Anisotropy	0.128	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 57.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8222	wwPDB-VP
Average B, all atoms (Å ²)	80.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, GOL, NAG, PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.11	0/1461	0.30	0/1995
1	B	0.09	0/1288	0.23	0/1765
2	C	0.10	0/719	0.27	0/969
2	D	0.10	0/711	0.27	0/966
All	All	0.10	0/4179	0.27	0/5695

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1416	1303	1301	1	0
1	B	1254	1100	1096	2	0
2	C	705	690	690	0	0
2	D	697	649	645	0	0
3	A	70	68	65	0	0
3	B	28	27	26	0	0
4	A	7	10	10	0	0
4	B	4	5	5	0	0
5	A	6	8	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	D	12	17	15	0	0
7	A	77	0	0	0	0
7	B	10	0	0	0	0
7	C	32	0	0	0	0
7	D	27	0	0	0	0
All	All	4345	3877	3861	3	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (3) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:94:GLN:OE1	1:A:109:ARG:NH2	2.32	0.63
1:B:83:GLN:NE2	1:B:85:GLU:OE2	2.37	0.57
1:B:94:GLN:OE1	1:B:109:ARG:NH1	2.39	0.55

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	177/175 (101%)	174 (98%)	3 (2%)	0	100	100
1	B	163/175 (93%)	160 (98%)	3 (2%)	0	100	100
2	C	86/101 (85%)	83 (96%)	3 (4%)	0	100	100
2	D	88/101 (87%)	84 (96%)	4 (4%)	0	100	100
All	All	514/552 (93%)	501 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	152/158 (96%)	152 (100%)	0	100	100
1	B	126/158 (80%)	126 (100%)	0	100	100
2	C	71/84 (84%)	71 (100%)	0	100	100
2	D	66/84 (79%)	66 (100%)	0	100	100
All	All	415/484 (86%)	415 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	72	GLN
1	A	135	HIS
1	B	72	GLN
1	B	111	HIS
1	B	115	ASN
2	D	90	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	403	1	14,14,15	0.24	0	17,19,21	0.40	0
4	PEG	A	406	-	6,6,6	0.42	0	5,5,5	0.28	0
4	PEG	B	403	-	3,3,6	0.39	0	2,2,5	0.39	0
3	NAG	B	402	1	14,14,15	0.23	0	17,19,21	0.44	0
5	GOL	A	407	-	5,5,5	0.99	0	5,5,5	1.00	0
3	NAG	B	401	1	14,14,15	0.21	0	17,19,21	0.53	0
3	NAG	A	405	1	14,14,15	0.21	0	17,19,21	0.43	0
6	PG4	D	501	-	11,11,12	0.50	0	10,10,11	0.28	0
3	NAG	A	404	1	14,14,15	0.18	0	17,19,21	0.49	0
3	NAG	A	402	1	14,14,15	0.29	0	17,19,21	0.55	0
3	NAG	A	401	1	14,14,15	0.15	0	17,19,21	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	403	1	-	2/6/23/26	0/1/1/1
4	PEG	A	406	-	-	0/4/4/4	-
4	PEG	B	403	-	-	0/1/1/4	-
3	NAG	B	402	1	-	2/6/23/26	0/1/1/1
5	GOL	A	407	-	-	2/4/4/4	-
3	NAG	B	401	1	-	2/6/23/26	0/1/1/1
3	NAG	A	405	1	-	2/6/23/26	0/1/1/1
6	PG4	D	501	-	-	1/9/9/10	-
3	NAG	A	404	1	-	0/6/23/26	0/1/1/1
3	NAG	A	402	1	-	3/6/23/26	0/1/1/1
3	NAG	A	401	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	407	GOL	O1-C1-C2-C3
3	B	402	NAG	C4-C5-C6-O6
3	A	401	NAG	C4-C5-C6-O6
3	A	405	NAG	O5-C5-C6-O6
3	B	401	NAG	O5-C5-C6-O6
3	A	401	NAG	O5-C5-C6-O6
3	B	402	NAG	O5-C5-C6-O6
3	A	405	NAG	C4-C5-C6-O6
6	D	501	PG4	O2-C3-C4-O3
3	B	401	NAG	C4-C5-C6-O6
5	A	407	GOL	O1-C1-C2-O2
3	A	403	NAG	C4-C5-C6-O6
3	A	403	NAG	O5-C5-C6-O6
3	A	402	NAG	O5-C5-C6-O6
3	A	402	NAG	C3-C2-N2-C7
3	A	402	NAG	C1-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	172/175 (98%)	0.35	6 (3%)	47	53	26, 62, 85, 111	7 (4%)
1	B	168/175 (96%)	1.48	37 (22%)	2	2	70, 105, 121, 128	1 (0%)
2	C	88/101 (87%)	0.43	4 (4%)	38	44	45, 74, 96, 105	0
2	D	90/101 (89%)	0.51	4 (4%)	39	45	46, 76, 104, 114	0
All	All	518/552 (93%)	0.76	51 (9%)	13	15	26, 79, 115, 128	8 (1%)

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	174	GLN	4.0
1	B	32	ALA	3.8
1	B	132	TYR	3.6
1	B	75	LEU	3.5
1	B	134	HIS	3.5
1	B	92	LEU	3.5
1	B	11	PHE	3.4
1	A	163	VAL	3.4
2	C	14	LEU	3.2
2	D	67	GLY	3.2
1	B	88	ILE	3.2
1	B	57	PHE	3.1
1	A	175	GLY	3.1
1	B	53	ALA	3.0
1	B	103	GLU	3.0
1	B	121	VAL	2.9
1	B	93	LEU	2.9
1	B	157	LEU	2.8
1	A	159	GLY	2.8
2	D	72	LEU	2.8
1	B	139	PHE	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	123	TYR	2.7
1	A	158	PHE	2.6
2	C	100	VAL	2.6
1	B	140	TYR	2.6
1	B	64	ASP	2.4
1	B	144	ALA	2.4
1	B	51	SER	2.4
1	B	141	ILE	2.4
2	C	13	SER	2.3
1	B	136	ASN	2.3
1	B	16	TRP	2.3
1	B	25	VAL	2.3
1	A	140	TYR	2.2
1	B	33	TYR	2.2
1	B	116	THR	2.2
2	C	86	ASN	2.2
1	B	145	THR	2.2
1	B	90	TRP	2.2
1	B	133	PHE	2.2
2	D	11	GLU	2.2
1	A	76[A]	SER	2.2
1	B	84	LEU	2.1
1	B	142	PRO	2.1
2	D	99	PRO	2.1
1	B	165	SER	2.1
1	B	122	THR	2.1
1	B	115	ASN	2.1
1	B	27	LEU	2.0
1	B	163	VAL	2.0
1	B	13	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	B	402	14/15	0.50	0.15	97,130,159,165	0
3	NAG	A	404	14/15	0.68	0.17	88,108,133,152	0
3	NAG	A	405	14/15	0.74	0.14	93,112,135,138	0
4	PEG	B	403	4/7	0.75	0.14	74,88,103,105	0
3	NAG	A	403	14/15	0.79	0.10	91,107,124,128	0
6	PG4	D	501	12/13	0.85	0.18	67,94,117,129	0
5	GOL	A	407	6/6	0.86	0.14	72,90,105,109	0
3	NAG	A	402	14/15	0.89	0.10	56,75,96,100	0
4	PEG	A	406	7/7	0.90	0.14	65,90,109,112	0
3	NAG	B	401	14/15	0.90	0.13	81,97,109,118	0
3	NAG	A	401	14/15	0.94	0.09	45,69,93,95	0

6.5 Other polymers [i](#)

There are no such residues in this entry.