



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:34 PM UTC

PDB ID : 1MOR / pdb\_00001mor  
Title : ISOMERASE DOMAIN OF GLUCOSAMINE 6-PHOSPHATE SYNTHASE  
COMPLEXED WITH GLUCOSE 6-PHOSPHATE  
Authors : Teplyakov, A.  
Deposited on : 1997-04-12  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	368	

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3064 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLUCOSAMINE 6-PHOSPHATE SYNTHASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	366	2825	1784	483	546	12	0	2	0

- Molecule 2 is 6-O-phosphono-alpha-D-glucopyranose (CCD ID: G6P) (formula: C<sub>6</sub>H<sub>13</sub>O<sub>9</sub>P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
2	A	1	16	6	9	1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	223	Total	O	0	0
			223	223		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GLUCOSAMINE 6-PHOSPHATE SYNTHASE



## 4 Data and refinement statistics i

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	146.10Å 146.10Å 173.90Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	10.00 – 1.90 10.00 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.5 (10.00-1.90) 98.5 (10.00-1.90)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.68 (at 1.90Å)	Xtriage
Refinement program	PROLSQ	Depositor
R, $R_{free}$	0.191 , (Not available) (Not available) , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.8	Xtriage
Anisotropy	0.253	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 108.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.017 for $-2/3^*h-1/3^*k+2/3^*l,-1/3^*h-2/3^*k-2/3^*l,2/3^*h-2/3^*k+1/3^*l$ 0.006 for $-h,1/3^*h-1/3^*k+2/3^*l,2/3^*h+4/3^*k+1/3^*l$ 0.004 for $-1/3^*h+1/3^*k-2/3^*l,-k,-4/3^*h-2/3^*k+1/3^*l$	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3064	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: G6P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.94	1/2881 (0.0%)	1.66	44/3899 (1.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	520	ALA	CA-C	-5.77	1.49	1.53

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	244	ASP	CA-CB-CG	9.18	121.78	112.60
1	A	460	ASP	CA-CB-CG	7.44	120.04	112.60
1	A	554	VAL	O-C-N	7.04	130.78	123.18
1	A	560	HIS	CA-CB-CG	-6.94	106.86	113.80
1	A	389	LEU	CA-C-O	-6.88	113.17	120.40
1	A	554	VAL	CB-CA-C	-6.73	100.60	110.62
1	A	445	GLU	N-CA-C	-6.63	104.12	111.82
1	A	586	HIS	CA-CB-CG	-6.38	107.42	113.80
1	A	320	ILE	CB-CA-C	-6.35	103.80	110.53
1	A	445	GLU	CA-CB-CG	6.33	126.76	114.10
1	A	320	ILE	CA-C-N	6.29	126.63	119.83
1	A	320	ILE	C-N-CA	6.29	126.63	119.83
1	A	523	ASN	CA-CB-CG	-6.14	106.46	112.60
1	A	312	TYR	CA-C-N	6.05	128.67	120.44
1	A	312	TYR	C-N-CA	6.05	128.67	120.44
1	A	506	PRO	N-CA-CB	6.01	109.49	103.48
1	A	517	ILE	CA-C-O	-5.93	114.19	120.48
1	A	339	ARG	NE-CZ-NH1	-5.92	115.58	121.50
1	A	558	ASN	CA-CB-CG	-5.87	106.73	112.60
1	A	504	HIS	CA-CB-CG	-5.80	108.00	113.80

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	575	PHE	CA-CB-CG	5.79	119.59	113.80
1	A	562	ILE	N-CA-CB	5.61	120.49	111.23
1	A	367	TYR	CA-C-O	-5.58	114.69	121.05
1	A	374	CYS	N-CA-CB	-5.55	102.39	111.66
1	A	376	VAL	CA-C-N	5.49	125.41	119.76
1	A	376	VAL	C-N-CA	5.49	125.41	119.76
1	A	259	GLN	CA-C-N	5.48	124.67	118.97
1	A	259	GLN	C-N-CA	5.48	124.67	118.97
1	A	337	VAL	N-CA-CB	5.47	117.75	110.26
1	A	296	GLN	OE1-CD-NE2	-5.46	117.14	122.60
1	A	578	VAL	CA-C-N	5.45	125.10	119.05
1	A	578	VAL	C-N-CA	5.45	125.10	119.05
1	A	365	LEU	N-CA-C	5.41	119.99	113.17
1	A	605	VAL	N-CA-CB	-5.39	106.28	111.89
1	A	306	SER	CA-CB-OG	5.38	121.86	111.10
1	A	598	PRO	N-CA-CB	5.33	107.98	103.35
1	A	520	ALA	CB-CA-C	-5.25	106.43	111.00
1	A	554	VAL	CA-C-O	-5.22	114.56	120.67
1	A	280	GLU	N-CA-C	-5.19	107.33	113.97
1	A	423	LYS	N-CA-C	-5.11	106.83	113.16
1	A	314	PHE	N-CA-CB	5.08	117.58	110.12
1	A	441	PRO	N-CA-CB	5.06	109.03	103.26
1	A	465	HIS	CA-CB-CG	-5.06	108.74	113.80
1	A	438	GLN	OE1-CD-NE2	5.04	127.64	122.60

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2825	0	2846	66	0
2	A	16	0	11	0	0
3	A	223	0	0	4	0
All	All	3064	0	2857	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:558:ASN:HD22	1:A:558:ASN:H	1.04	0.95
1:A:340:ASN:HA	1:A:368:LEU:HD13	1.49	0.94
1:A:434:VAL:O	1:A:438:GLN:HG3	1.70	0.91
1:A:558:ASN:HD22	1:A:558:ASN:N	1.62	0.86
1:A:558:ASN:H	1:A:558:ASN:ND2	1.70	0.86
1:A:337:VAL:HG21	1:A:365:LEU:O	1.80	0.82
1:A:420:SER:HA	1:A:425:LEU:HD12	1.70	0.73
1:A:412:LEU:O	1:A:416:VAL:HG23	1.93	0.67
1:A:537:ARG:NH1	1:A:537:ARG:HG3	2.10	0.65
1:A:270:ARG:HD2	1:A:414:MET:HE1	1.80	0.62
1:A:377:PRO:HA	1:A:390:MET:CE	2.29	0.62
1:A:377:PRO:HA	1:A:390:MET:HE2	1.80	0.62
1:A:420:SER:CA	1:A:425:LEU:HD12	2.29	0.61
1:A:447:MET:HE2	1:A:575:PHE:CZ	2.36	0.61
1:A:337:VAL:HG23	3:A:221:HOH:O	2.00	0.61
1:A:250:HIS:HD2	1:A:585:TYR:OH	1.84	0.60
1:A:320:ILE:CD1	1:A:419:LEU:HD13	2.34	0.56
1:A:440:LEU:HB3	1:A:441:PRO:HD3	1.85	0.56
1:A:537:ARG:HG3	1:A:537:ARG:HH11	1.69	0.56
1:A:264:LYS:O	1:A:268:THR:HG23	2.05	0.56
1:A:419:LEU:O	1:A:423:LYS:HG2	2.08	0.54
1:A:377:PRO:CA	1:A:390:MET:HE2	2.37	0.54
1:A:578:VAL:N	1:A:579:PRO:HD2	2.22	0.54
1:A:310:SER:HB3	1:A:412:LEU:HD13	1.89	0.54
1:A:271:ILE:HD12	1:A:434:VAL:HG13	1.90	0.53
1:A:293:GLU:OE2	1:A:340:ASN:ND2	2.44	0.51
1:A:346:LEU:HD23	1:A:373:ILE:HB	1.92	0.51
1:A:562:ILE:HG22	1:A:564:MET:HE2	1.93	0.50
1:A:320:ILE:HD12	1:A:419:LEU:CD1	2.42	0.50
1:A:260:PRO:HG3	1:A:444:ILE:HG22	1.94	0.49
1:A:558:ASN:N	1:A:558:ASN:ND2	2.32	0.49
1:A:288:LEU:HD23	1:A:387:LEU:HD11	1.94	0.48
1:A:421:ARG:O	1:A:424:GLY:N	2.39	0.48
1:A:558:ASN:ND2	1:A:558:ASN:C	2.71	0.48
1:A:377:PRO:CA	1:A:390:MET:CE	2.91	0.48
1:A:518:VAL:CG2	1:A:543:LEU:HD11	2.44	0.48
1:A:376:VAL:HA	1:A:377:PRO:HD2	1.72	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:480:LEU:HD23	1:A:496:ALA:HB3	1.98	0.46
1:A:461:PHE:CZ	1:A:517:ILE:HD11	2.52	0.45
1:A:356:LEU:HD13	1:A:381:LEU:HG	1.99	0.45
1:A:276:VAL:HG21	1:A:417:ALA:HB3	1.98	0.45
1:A:294:HIS:HD1	1:A:341:SER:HG	1.65	0.44
1:A:340:ASN:OD1	1:A:368:LEU:HD11	2.17	0.44
1:A:472:ARG:HG2	1:A:473:GLY:N	2.31	0.44
1:A:537:ARG:HH11	1:A:537:ARG:CG	2.30	0.43
1:A:461:PHE:CE2	1:A:517:ILE:HD11	2.53	0.43
1:A:320:ILE:HD12	1:A:419:LEU:HD13	2.01	0.43
1:A:376:VAL:C	1:A:390:MET:HE1	2.44	0.43
1:A:363:LYS:HE2	3:A:109:HOH:O	2.19	0.43
1:A:430:GLU:O	1:A:434:VAL:HG23	2.19	0.42
1:A:447:MET:HE2	1:A:575:PHE:CE2	2.54	0.42
1:A:476:TYR:HB3	1:A:477:PRO:HD3	2.00	0.42
1:A:320:ILE:HA	1:A:321:PRO:HD3	1.89	0.42
1:A:380:SER:O	1:A:384:GLU:HG3	2.19	0.42
1:A:530:LYS:HD3	1:A:530:LYS:HA	1.81	0.42
1:A:572:ALA:N	1:A:573:PRO:CD	2.82	0.42
1:A:476:TYR:N	1:A:477:PRO:CD	2.82	0.42
1:A:399:VAL:HB	1:A:598:PRO:HD2	2.02	0.41
1:A:294:HIS:ND1	1:A:341:SER:OG	2.50	0.41
1:A:359:LEU:HD12	1:A:381:LEU:HD23	2.01	0.41
1:A:549:GLN:HG3	1:A:550:ASP:N	2.35	0.41
1:A:341:SER:HB3	1:A:367:TYR:CD1	2.55	0.41
1:A:254:LYS:HE2	3:A:49:HOH:O	2.20	0.40
1:A:278:LEU:HD12	1:A:418:LYS:HG2	2.03	0.40
1:A:260:PRO:HD2	3:A:105:HOH:O	2.22	0.40
1:A:558:ASN:HD22	1:A:558:ASN:C	2.28	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	366/368 (100%)	356 (97%)	9 (2%)	1 (0%)	36 29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	505	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	307/306 (100%)	294 (96%)	13 (4%)	26 19

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	244	ASP
1	A	286	ASP
1	A	341	SER
1	A	356	LEU
1	A	359	LEU
1	A	368	LEU
1	A	401	SER
1	A	421	ARG
1	A	530	LYS
1	A	554	VAL
1	A	558	ASN
1	A	601	LEU
1	A	608	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	250	HIS
1	A	259	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	296	GLN
1	A	475	GLN
1	A	549	GLN
1	A	558	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	G6P	A	609	-	16,16,16	1.07	0	23,24,24	1.52	4 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	G6P	A	609	-	-	1/6/26/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	A	609	G6P	C1-O5-C5	4.32	122.01	113.65
2	A	609	G6P	O4-C4-C3	2.65	116.61	110.38
2	A	609	G6P	O2P-P-O1P	2.64	117.69	107.80
2	A	609	G6P	O1-C1-O5	-2.13	104.08	110.41

There are no chirality outliers.


All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	609	G6P	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	366/368 (99%)	-0.38	8 (2%) 62 66	11, 25, 62, 100	2 (0%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	283	PRO	3.6
1	A	243	GLY	3.0
1	A	607	VAL	2.9
1	A	244	ASP	2.6
1	A	272	SER	2.6
1	A	282	GLY	2.4
1	A	284	ASN	2.1
1	A	247	ILE	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	G6P	A	609	16/16	0.98	0.04	16,21,24,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.