



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 5, 2026 – 01:52 PM UTC

PDB ID : 1MOV / pdb_00001mov
Title : Crystal structure of Coral protein mutant
Authors : Prescott, M.; Ling, M.; Beddoe, T.; Oakley, A.J.; Dove, S.; Hoegh-Guldberg, O.; Devenish, R.J.; Rossjohn, J.
Deposited on : 2002-09-10
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	219	 74% 24% .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1930 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GFP-like non-fluorescent chromoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	219	1742	1115	289	326	12	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	5	GLY	-	expression tag	UNP P83690
A	6	GLY	-	expression tag	UNP P83690
A	66	CRQ	GLN	chromophore	UNP P83690
A	66	CRQ	TYR	chromophore	UNP P83690
A	66	CRQ	GLY	chromophore	UNP P83690
A	146	SER	HIS	engineered mutation	UNP P83690

- Molecule 2 is IODIDE ION (CCD ID: IOD) (formula: I).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	5	Total	I	0	0
			5	5		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	183	Total	O	0	0
			183	183		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: GFP-like non-fluorescent chromoprotein



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	131.39Å 131.39Å 150.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.40	Depositor
% Data completeness (in resolution range)	99.7 (20.00-2.40)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.225 , 0.249	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1930	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IOD, CRQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/1766	0.89	7/2387 (0.3%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	209	THR	N-CA-C	-6.23	105.68	113.28
1	A	206	LYS	N-CA-C	6.16	118.78	111.33
1	A	133	GLY	CA-C-N	6.01	126.44	119.47
1	A	133	GLY	C-N-CA	6.01	126.44	119.47
1	A	122	VAL	N-CA-C	5.43	116.54	108.46
1	A	83	VAL	N-CA-C	5.39	115.59	110.53
1	A	174	LEU	N-CA-C	5.32	118.35	110.48

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1742	0	1676	39	0
2	A	5	0	0	0	0
3	A	183	0	0	1	0
All	All	1930	0	1676	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:42:GLN:HE22	1:A:69:SER:HB3	1.42	0.85
1:A:42:GLN:HE22	1:A:69:SER:CB	1.90	0.84
1:A:80:PRO:HG2	1:A:190:PRO:HA	1.75	0.65
1:A:85:GLN:HE22	1:A:188:LYS:HB3	1.62	0.63
1:A:62:SER:OG	1:A:63:PRO:HD3	2.01	0.60
1:A:80:PRO:HG2	1:A:190:PRO:CA	2.32	0.60
1:A:28:GLU:HB2	1:A:50:LYS:HB3	1.85	0.58
1:A:148:GLU:OE1	1:A:197:ARG:NH2	2.37	0.58
1:A:5:GLY:HA3	1:A:38:TYR:OH	2.07	0.55
1:A:109:ASN:C	1:A:109:ASN:ND2	2.64	0.55
1:A:14:TYR:HB3	1:A:118:PHE:HB2	1.91	0.53
1:A:198:LYS:HE2	1:A:200:ASP:OD2	2.10	0.52
1:A:42:GLN:NE2	1:A:69:SER:CB	2.68	0.52
1:A:79:ILE:HD11	1:A:221:LYS:HG3	1.93	0.51
1:A:149:ARG:HD2	1:A:149:ARG:C	2.35	0.51
1:A:42:GLN:NE2	1:A:69:SER:HB3	2.19	0.50
1:A:37:PRO:HA	1:A:72:PHE:HA	1.94	0.49
1:A:93:TRP:CE2	1:A:109:ASN:HB3	2.47	0.49
1:A:8:ILE:HG23	1:A:12:MET:HE2	1.94	0.49
1:A:85:GLN:NE2	1:A:188:LYS:H	2.12	0.47
1:A:109:ASN:ND2	1:A:110:ASP:N	2.61	0.47
1:A:205:ASN:ND2	1:A:210:SER:HB2	2.28	0.47
1:A:149:ARG:HH21	1:A:160:ASN:HD22	1.62	0.47
1:A:12:MET:HE1	1:A:113:ILE:HD11	1.97	0.46
1:A:149:ARG:HH21	1:A:160:ASN:ND2	2.14	0.46
1:A:85:GLN:HE22	1:A:188:LYS:H	1.63	0.45
1:A:80:PRO:CG	1:A:190:PRO:HA	2.46	0.45
1:A:16:VAL:O	1:A:30:GLU:HA	2.17	0.45
1:A:89:GLU:CD	1:A:185:LYS:HD3	2.42	0.45
1:A:85:GLN:HE22	1:A:188:LYS:CB	2.29	0.44
1:A:14:TYR:OH	1:A:44:VAL:HB	2.19	0.43
1:A:87:PHE:HB3	1:A:88:PRO:HA	2.01	0.43
1:A:27:PHE:C	1:A:27:PHE:CD1	2.97	0.42
1:A:66:CRQ:HB11	1:A:215:GLU:OE1	2.19	0.42
1:A:121:HIS:HD2	3:A:443:HOH:O	2.02	0.42
1:A:70:ILE:N	1:A:71:PRO:CD	2.84	0.41
1:A:109:ASN:C	1:A:109:ASN:HD22	2.27	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:97:MET:HB2	1:A:105:CYS:HB2	2.03	0.41
1:A:127:LEU:O	1:A:128:ASN:HB2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/219 (98%)	210 (98%)	4 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	187/187 (100%)	184 (98%)	3 (2%)	55	76

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	109	ASN
1	A	199	LEU
1	A	210	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	GLN
1	A	85	GLN
1	A	109	ASN
1	A	121	HIS
1	A	160	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CRQ	A	66	1	25,25,26	2.52	9 (36%)	28,34,36	2.69	6 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRQ	A	66	1	-	3/10/32/33	0/2/2/2

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66	CRQ	CA3-C3	-8.88	1.17	1.49
1	A	66	CRQ	CA3-N3	3.98	1.54	1.47
1	A	66	CRQ	C1-N3	3.83	1.44	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66	CRQ	CB2-CA2	2.61	1.37	1.35
1	A	66	CRQ	C1-CA1	-2.30	1.44	1.48
1	A	66	CRQ	CE1-CD1	2.23	1.42	1.38
1	A	66	CRQ	O3-C3	2.16	1.32	1.20
1	A	66	CRQ	CE2-CD2	2.08	1.42	1.38
1	A	66	CRQ	CG2-CB2	2.04	1.50	1.46

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	CRQ	CG2-CB2-CA2	8.47	139.94	129.87
1	A	66	CRQ	CB2-CA2-N2	-7.52	118.55	128.76
1	A	66	CRQ	CB2-CA2-C2	6.97	130.81	122.36
1	A	66	CRQ	C3-CA3-N3	2.91	119.04	112.43
1	A	66	CRQ	C2-CA2-N2	2.36	110.64	108.95
1	A	66	CRQ	CA2-N2-C1	2.05	107.87	104.09

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	66	CRQ	C1-CA1-CB1-CG1
1	A	66	CRQ	C2-CA2-CB2-CG2
1	A	66	CRQ	N2-CA2-CB2-CG2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	66	CRQ	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.