



wwPDB X-ray Structure Validation Summary Report

Mar 9, 2026 – 08:15 AM UTC

PDB ID : 4MOK / pdb_00004mok
Title : Pyranose 2-oxidase H167A mutant soaked with 3-fluorinated galactose (not bound)
Authors : Tan, T.C.; Spadiut, O.; Gandini, R.; Haltrich, D.; Divne, C.
Deposited on : 2013-09-12
Resolution : 1.90 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	633	 2% 80% 10% • 9%
1	B	633	 3% 81% 9% • 9%
1	C	633	 2% 83% 7% 9%
1	D	633	 2% 80% 9% • 10%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 20179 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyranose 2-oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	576	4537	2865	775	872	25	0	0	0
1	B	575	4529	2860	774	871	24	0	0	0
1	C	574	4521	2856	773	868	24	0	0	0
1	D	572	4502	2841	771	866	24	0	0	0

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	ALA	SER	SEE REMARK 999	UNP Q7ZA32
A	167	ALA	HIS	engineered mutation	UNP Q7ZA32
A	623	ALA	-	expression tag	UNP Q7ZA32
A	624	ALA	-	expression tag	UNP Q7ZA32
A	625	ALA	-	expression tag	UNP Q7ZA32
A	626	LEU	-	expression tag	UNP Q7ZA32
A	627	GLU	-	expression tag	UNP Q7ZA32
A	628	HIS	-	expression tag	UNP Q7ZA32
A	629	HIS	-	expression tag	UNP Q7ZA32
A	630	HIS	-	expression tag	UNP Q7ZA32
A	631	HIS	-	expression tag	UNP Q7ZA32
A	632	HIS	-	expression tag	UNP Q7ZA32
A	633	HIS	-	expression tag	UNP Q7ZA32
B	2	ALA	SER	SEE REMARK 999	UNP Q7ZA32
B	167	ALA	HIS	engineered mutation	UNP Q7ZA32
B	623	ALA	-	expression tag	UNP Q7ZA32
B	624	ALA	-	expression tag	UNP Q7ZA32
B	625	ALA	-	expression tag	UNP Q7ZA32
B	626	LEU	-	expression tag	UNP Q7ZA32
B	627	GLU	-	expression tag	UNP Q7ZA32
B	628	HIS	-	expression tag	UNP Q7ZA32

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	629	HIS	-	expression tag	UNP Q7ZA32
B	630	HIS	-	expression tag	UNP Q7ZA32
B	631	HIS	-	expression tag	UNP Q7ZA32
B	632	HIS	-	expression tag	UNP Q7ZA32
B	633	HIS	-	expression tag	UNP Q7ZA32
C	2	ALA	SER	SEE REMARK 999	UNP Q7ZA32
C	167	ALA	HIS	engineered mutation	UNP Q7ZA32
C	623	ALA	-	expression tag	UNP Q7ZA32
C	624	ALA	-	expression tag	UNP Q7ZA32
C	625	ALA	-	expression tag	UNP Q7ZA32
C	626	LEU	-	expression tag	UNP Q7ZA32
C	627	GLU	-	expression tag	UNP Q7ZA32
C	628	HIS	-	expression tag	UNP Q7ZA32
C	629	HIS	-	expression tag	UNP Q7ZA32
C	630	HIS	-	expression tag	UNP Q7ZA32
C	631	HIS	-	expression tag	UNP Q7ZA32
C	632	HIS	-	expression tag	UNP Q7ZA32
C	633	HIS	-	expression tag	UNP Q7ZA32
D	2	ALA	SER	SEE REMARK 999	UNP Q7ZA32
D	167	ALA	HIS	engineered mutation	UNP Q7ZA32
D	623	ALA	-	expression tag	UNP Q7ZA32
D	624	ALA	-	expression tag	UNP Q7ZA32
D	625	ALA	-	expression tag	UNP Q7ZA32
D	626	LEU	-	expression tag	UNP Q7ZA32
D	627	GLU	-	expression tag	UNP Q7ZA32
D	628	HIS	-	expression tag	UNP Q7ZA32
D	629	HIS	-	expression tag	UNP Q7ZA32
D	630	HIS	-	expression tag	UNP Q7ZA32
D	631	HIS	-	expression tag	UNP Q7ZA32
D	632	HIS	-	expression tag	UNP Q7ZA32
D	633	HIS	-	expression tag	UNP Q7ZA32

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	53	27	9	15	2	0	0
2	B	1	53	27	9	15	2	0	0
2	C	1	53	27	9	15	2	0	0
2	D	1	53	27	9	15	2	0	0

- Molecule 3 is DODECAETHYLENE GLYCOL (CCD ID: 12P) (formula: $C_{24}H_{50}O_{13}$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			16	10	6		
3	B	1	Total	C	O	0	0
			16	10	6		
3	C	1	Total	C	O	0	0
			16	10	6		
3	D	1	Total	C	O	0	0
			13	8	5		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	C	1	Total	C	N	O	S	0	0
			12	6	1	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	536	Total	O	0	0
			536	536		
5	B	498	Total	O	0	0
			498	498		
5	C	449	Total	O	0	0
			449	449		



Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	310	Total 310	O 310	0	0



● Molecule 1: Pyranose 2-oxidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	99.88Å 102.30Å 136.96Å 90.00° 90.78° 90.00°	Depositor
Resolution (Å)	47.92 – 1.90 47.92 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (47.92-1.90) 99.8 (47.92-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.40 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.170 , 0.214 0.177 , 0.222	Depositor DCC
R_{free} test set	1066 reflections (0.49%)	wwPDB-VP
Wilson B-factor (Å ²)	24.3	Xtriage
Anisotropy	0.173	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 33.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.011 for -k,-h,-l 0.008 for k,h,-l 0.018 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	20179	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, MES, 12P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.39	21/4652 (0.5%)	1.14	9/6325 (0.1%)
1	B	1.31	15/4644 (0.3%)	1.12	6/6315 (0.1%)
1	C	1.25	3/4636 (0.1%)	1.10	13/6304 (0.2%)
1	D	1.10	4/4616 (0.1%)	1.05	6/6277 (0.1%)
All	All	1.27	43/18548 (0.2%)	1.10	34/25221 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

The worst 5 of 43 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	443	HIS	CG-CD2	8.79	1.45	1.35
1	B	299	HIS	CG-CD2	8.70	1.45	1.35
1	A	349	LEU	C-O	-7.73	1.20	1.23
1	B	299	HIS	ND1-CE1	7.27	1.39	1.32
1	B	349	LEU	C-O	6.91	1.27	1.23

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	345	PRO	CA-C-N	-8.07	112.40	120.31
1	C	345	PRO	C-N-CA	-8.07	112.40	120.31
1	D	458	ALA	N-CA-C	-7.87	100.01	110.24
1	B	385	THR	CA-C-N	7.24	127.29	119.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	385	THR	C-N-CA	7.24	127.29	119.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	230	TYR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4537	0	4386	36	0
1	B	4529	0	4377	36	0
1	C	4521	0	4373	27	0
1	D	4502	0	4353	35	0
2	A	53	0	31	1	0
2	B	53	0	30	1	0
2	C	53	0	30	0	0
2	D	53	0	29	1	0
3	A	16	0	21	0	0
3	B	16	0	21	0	0
3	C	16	0	21	0	0
3	D	13	0	16	0	0
4	A	12	0	13	4	0
4	C	12	0	13	0	0
5	A	536	0	0	2	0
5	B	498	0	0	10	0
5	C	449	0	0	3	0
5	D	310	0	0	3	0
All	All	20179	0	17714	133	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 133 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:133:ALA:HB3	4:A:803:MES:H71	1.35	1.03
1:B:380:MET:HE1	1:B:413:LYS:HB2	1.47	0.95
1:A:133:ALA:CB	4:A:803:MES:H71	2.03	0.89
1:A:133:ALA:HB2	4:A:803:MES:O1S	1.74	0.88
1:B:417:MET:SD	5:B:1356:HOH:O	2.38	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	574/633 (91%)	562 (98%)	12 (2%)	0	100	100
1	B	573/633 (90%)	553 (96%)	18 (3%)	2 (0%)	36	29
1	C	572/633 (90%)	550 (96%)	21 (4%)	1 (0%)	43	36
1	D	570/633 (90%)	549 (96%)	21 (4%)	0	100	100
All	All	2289/2532 (90%)	2214 (97%)	72 (3%)	3 (0%)	48	40

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	82	SER
1	C	82	SER
1	B	459	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	503/547 (92%)	500 (99%)	3 (1%)	78	81
1	B	502/547 (92%)	496 (99%)	6 (1%)	63	63
1	C	501/547 (92%)	495 (99%)	6 (1%)	63	63
1	D	499/547 (91%)	489 (98%)	10 (2%)	48	46
All	All	2005/2188 (92%)	1980 (99%)	25 (1%)	63	63

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	611	GLN
1	D	128	PRO
1	D	560	LYS
1	D	100	ILE
1	D	272	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	499	GLN
1	C	344	ASN
1	D	499	GLN
1	D	419	HIS
1	D	450	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	12P	D	802	-	12,12,36	0.52	0	11,11,35	0.47	0
3	12P	C	802	-	15,15,36	0.56	0	14,14,35	0.50	0
2	FAD	D	801	-	58,58,58	1.52	9 (15%)	85,89,89	2.26	31 (36%)
4	MES	C	803	-	12,12,12	2.07	1 (8%)	15,16,16	1.59	4 (26%)
2	FAD	A	801	-	58,58,58	1.41	7 (12%)	85,89,89	2.24	23 (27%)
2	FAD	C	801	-	58,58,58	1.42	9 (15%)	85,89,89	2.26	27 (31%)
2	FAD	B	801	-	58,58,58	1.31	8 (13%)	85,89,89	2.18	24 (28%)
3	12P	B	802	-	15,15,36	0.64	0	14,14,35	0.90	1 (7%)
4	MES	A	803	-	12,12,12	1.94	1 (8%)	15,16,16	1.80	4 (26%)
3	12P	A	802	-	15,15,36	0.65	0	14,14,35	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	12P	D	802	-	-	1/10/10/34	-
3	12P	C	802	-	-	3/13/13/34	-
2	FAD	D	801	-	-	3/34/50/50	0/6/6/6
4	MES	C	803	-	-	0/6/14/14	0/1/1/1
2	FAD	A	801	-	-	0/34/50/50	0/6/6/6
2	FAD	C	801	-	-	0/34/50/50	0/6/6/6
2	FAD	B	801	-	-	5/34/50/50	0/6/6/6
3	12P	B	802	-	-	3/13/13/34	-
4	MES	A	803	-	-	0/6/14/14	0/1/1/1
3	12P	A	802	-	-	1/13/13/34	-

The worst 5 of 35 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	803	MES	C8-S	-6.46	1.68	1.77
2	D	801	FAD	P-O3P	5.70	1.65	1.59
4	A	803	MES	C8-S	-5.53	1.69	1.77
2	A	801	FAD	C6-C5X	5.06	1.47	1.40
2	A	801	FAD	P-O3P	3.70	1.63	1.59

The worst 5 of 114 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	FAD	C4A-N9A-C8A	8.13	114.27	105.74
2	C	801	FAD	C4A-N9A-C8A	7.24	113.34	105.74
2	B	801	FAD	C4A-N9A-C8A	7.24	113.34	105.74
2	D	801	FAD	C4A-N9A-C8A	6.96	113.04	105.74
2	C	801	FAD	N9A-C8A-N7A	-6.53	104.67	113.94

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	801	FAD	PA-O3P-P-O5'
2	D	801	FAD	PA-O3P-P-O5'
3	B	802	12P	O13-C14-C15-O16
3	C	802	12P	O13-C14-C15-O16
2	B	801	FAD	P-O3P-PA-O1A


There are no ring outliers.


4 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	801	FAD	1	0
2	A	801	FAD	1	0
2	B	801	FAD	1	0
4	A	803	MES	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	576/633 (90%)	-0.31	13 (2%) 61 65	17, 23, 45, 83	0
1	B	575/633 (90%)	-0.21	18 (3%) 51 55	18, 25, 51, 90	0
1	C	574/633 (90%)	-0.14	13 (2%) 61 65	19, 29, 53, 89	0
1	D	572/633 (90%)	0.25	12 (2%) 63 67	22, 39, 65, 97	0
All	All	2297/2532 (90%)	-0.10	56 (2%) 59 63	17, 29, 58, 97	0

The worst 5 of 56 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	45	ILE	7.6
1	D	458	ALA	7.0
1	B	458	ALA	6.7
1	A	458	ALA	6.6
1	A	45	ILE	6.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	MES	A	803	12/12	0.85	0.14	45,58,60,62	0
3	12P	D	802	13/37	0.91	0.10	39,41,51,57	0
3	12P	B	802	16/37	0.91	0.11	31,37,50,59	0
3	12P	A	802	16/37	0.92	0.10	35,38,50,51	0
3	12P	C	802	16/37	0.92	0.10	28,36,52,55	0
4	MES	C	803	12/12	0.92	0.10	42,44,49,50	0
2	FAD	D	801	53/53	0.96	0.06	26,32,36,38	0
2	FAD	B	801	53/53	0.98	0.04	16,19,23,24	0
2	FAD	C	801	53/53	0.98	0.05	19,24,28,29	0
2	FAD	A	801	53/53	0.98	0.04	14,18,21,22	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around 12P B 802:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 12P A 802:

$2mF_o-DF_c$ (at 0.7 rnsd) in gray
 mF_o-DF_c (at 3 rnsd) in purple (negative)
and green (positive)





Electron density around 12P C 802:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD D 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD B 801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD C 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD A 801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.