



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:47 PM UTC

PDB ID : 5MOL / pdb_00005mol
Title : Human IgE-Fc crystal structure
Authors : Dore, K.A.; Davies, A.M.; Drinkwater, N.; Beavil, A.J.; McDonnell, J.M.; Sutton, B.J.
Deposited on : 2016-12-14
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	327	6% 89% 8% .
1	B	327	8% 91% 8% .
2	C	5	20% 80%
3	D	7	14% 86%

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 5750 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ig epsilon chain C region.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	318	Total	C	N	O	S	0	6	0
			2432	1529	409	483	11			
1	B	322	Total	C	N	O	S	0	9	0
			2501	1565	434	491	11			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	222	ASP	-	expression tag	UNP P01854
A	223	ILE	-	expression tag	UNP P01854
A	225	ALA	CYS	conflict	UNP P01854
A	265	GLN	ASN	conflict	UNP P01854
A	371	GLN	ASN	conflict	UNP P01854
B	222	ASP	-	expression tag	UNP P01854
B	223	ILE	-	expression tag	UNP P01854
B	225	ALA	CYS	conflict	UNP P01854
B	265	GLN	ASN	conflict	UNP P01854
B	371	GLN	ASN	conflict	UNP P01854


- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	5	Total	C	N	O	0	0	0
			61	34	2	25			


- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyran

ose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	7	Total	C	N	O	0	0	0
			83	46	2	35			

- Molecule 4 is 2-(2-METHOXYETHOXY)ETHANOL (CCD ID: PG0) (formula: C₅H₁₂O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			8	5	3		
4	B	1	Total	C	O	0	0
			8	5	3		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	1
			8	4	4		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			7	4	3		
6	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 7 is 2-(2-ETHOXYETHOXY)ETHANOL (CCD ID: AE3) (formula: C₆H₁₄O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	1
			18	12	6		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	295	Total	O	0	0
			295	295		
8	B	306	Total	O	0	0
			306	306		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ig epsilon chain C region



- Molecule 1: Ig epsilon chain C region



- Molecule 2: alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	130.67Å 75.78Å 79.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	79.82 – 1.75 79.82 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.5 (79.82-1.75) 99.5 (79.82-1.75)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 1.75Å)	Xtriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, R_{free}	0.197 , 0.232 0.203 , 0.233	Depositor DCC
R_{free} test set	4016 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	23.8	Xtriage
Anisotropy	0.184	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 49.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.016 for -h,l,k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5750	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PG0, BMA, MAN, PEG, AE3, EDO, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/2509	0.53	0/3433
1	B	0.40	0/2589	0.54	0/3540
All	All	0.40	0/5098	0.54	0/6973

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2432	0	2311	19	0
1	B	2501	0	2408	15	0
2	C	61	0	52	1	0
3	D	83	0	70	0	0
4	A	8	0	12	0	0
4	B	8	0	12	1	0
5	A	16	0	24	1	0
5	B	8	0	12	0	0
6	A	7	0	10	0	0
6	B	7	0	10	0	0
7	B	18	0	28	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	A	295	0	0	2	0
8	B	306	0	0	2	0
All	All	5750	0	4949	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:422:HIS:HB3	1:B:425:LEU:HD13	1.71	0.72
1:A:477:GLN:NE2	1:A:484:GLN:OE1	2.27	0.68
1:A:307:ASP:OD2	8:A:701:HOH:O	2.12	0.68
1:A:439:PRO:HG3	1:B:278:ASP:HB3	1.80	0.64
1:B:238:GLN:NE2	8:B:704:HOH:O	2.30	0.64
1:A:363:LEU:HD13	1:A:370:VAL:HG21	1.82	0.61
1:B:459:LYS:O	1:B:460[A]:ARG:HG3	2.04	0.58
1:A:420:VAL:HG13	1:A:429:LEU:HB3	1.85	0.57
4:B:608:PG0:H22	8:B:748:HOH:O	2.04	0.57
1:A:390:GLU:HG3	1:A:392:GLN:HE22	1.69	0.56
1:A:373:THR:HG23	1:A:419:ARG:HB3	1.89	0.54
1:B:475:SER:HA	7:B:609[A]:AE3:H3C2	1.90	0.53
1:A:342:ARG:O	5:A:607:EDO:H11	2.12	0.49
1:A:390:GLU:HG3	1:A:392:GLN:NE2	2.27	0.49
1:A:527:VAL:HG22	1:A:536[B]:THR:HG22	1.95	0.47
1:B:235:LYS:HE3	1:B:255:LEU:HD12	1.96	0.47
1:A:392:GLN:OE1	2:C:1:NAG:O6	2.33	0.46
1:A:337:SER:HB2	1:A:361:VAL:HG22	1.98	0.45
1:A:428:ALA:O	8:A:702:HOH:O	2.21	0.45
1:B:476:VAL:H	7:B:609[A]:AE3:H4C1	1.83	0.44
1:A:443:PRO:HG2	1:A:537[A]:VAL:HG11	1.99	0.44
1:B:470:MET:HG3	1:B:502:GLY:HA2	2.00	0.43
1:B:269:LEU:HB2	1:B:311:THR:HB	2.00	0.43
1:B:516:TRP:CD2	1:B:543[B]:VAL:HG11	2.54	0.43
1:A:436:THR:HG21	1:A:471:PRO:HG3	1.99	0.43
1:B:520:ASP:OD1	1:B:543[B]:VAL:HG12	2.19	0.43
1:A:368:GLY:HA3	1:A:423:PRO:HG2	2.02	0.42
1:B:258:GLY:HA2	1:B:291:ALA:O	2.20	0.42
1:A:372:LEU:HD21	1:A:418:CYS:SG	2.60	0.42
1:B:419:ARG:HD2	1:B:430:MET:HE3	2.02	0.42
1:A:493:THR:HG21	1:B:493:THR:HG21	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:452:GLU:HB2	1:B:460[B]:ARG:CZ	2.50	0.41
1:A:477:GLN:HE21	1:A:484:GLN:CD	2.29	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	322/327 (98%)	316 (98%)	6 (2%)	0	100	100
1	B	329/327 (101%)	322 (98%)	7 (2%)	0	100	100
All	All	651/654 (100%)	638 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	267/289 (92%)	265 (99%)	2 (1%)	76	67
1	B	280/289 (97%)	276 (99%)	4 (1%)	59	44
All	All	547/578 (95%)	541 (99%)	6 (1%)	65	52

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	369	THR
1	A	373	THR
1	B	369	THR
1	B	470	MET
1	B	477	GLN
1	B	518	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	332	ASN
1	A	392	GLN
1	A	477	GLN
1	B	238	GLN
1	B	424	HIS
1	B	481	ASN
1	B	484	GLN
1	B	518	GLN
1	B	535	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	1,2	14,14,15	0.24	0	17,19,21	0.54	0
2	NAG	C	2	2	14,14,15	0.29	0	17,19,21	0.58	0
2	BMA	C	3	2	11,11,12	0.81	0	15,15,17	1.10	1 (6%)
2	MAN	C	4	2	11,11,12	0.90	0	15,15,17	0.87	1 (6%)
2	MAN	C	5	2	11,11,12	1.26	2 (18%)	15,15,17	1.80	5 (33%)
3	NAG	D	1	1,3	14,14,15	0.28	0	17,19,21	0.83	1 (5%)
3	NAG	D	2	3	14,14,15	0.65	1 (7%)	17,19,21	0.63	0
3	BMA	D	3	3	11,11,12	0.54	0	15,15,17	0.70	0
3	MAN	D	4	3	11,11,12	0.83	0	15,15,17	1.03	2 (13%)
3	MAN	D	5	3	11,11,12	0.78	0	15,15,17	1.03	1 (6%)
3	MAN	D	6	3	11,11,12	1.07	0	15,15,17	1.04	1 (6%)
3	MAN	D	7	3	11,11,12	0.83	0	15,15,17	0.93	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	MAN	C	4	2	-	0/2/19/22	0/1/1/1
2	MAN	C	5	2	-	1/2/19/22	0/1/1/1
3	NAG	D	1	1,3	-	1/6/23/26	0/1/1/1
3	NAG	D	2	3	-	4/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1
3	MAN	D	4	3	-	0/2/19/22	0/1/1/1
3	MAN	D	5	3	-	2/2/19/22	0/1/1/1
3	MAN	D	6	3	-	0/2/19/22	0/1/1/1
3	MAN	D	7	3	-	2/2/19/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	5	MAN	C1-C2	2.61	1.58	1.52
2	C	5	MAN	C2-C3	2.40	1.56	1.52
3	D	2	NAG	O5-C1	-2.09	1.40	1.43

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	5	MAN	C2-C3-C4	3.44	116.91	110.86
2	C	5	MAN	C1-C2-C3	3.32	114.48	109.64
2	C	5	MAN	C1-O5-C5	3.05	116.27	112.19
3	D	6	MAN	C1-O5-C5	2.82	115.97	112.19
3	D	5	MAN	C1-O5-C5	2.71	115.82	112.19
2	C	5	MAN	C3-C4-C5	2.58	114.91	110.23
2	C	4	MAN	O2-C2-C3	-2.52	104.94	110.15
3	D	7	MAN	O2-C2-C3	-2.49	104.98	110.15
3	D	4	MAN	O2-C2-C3	-2.44	105.10	110.15
2	C	5	MAN	O2-C2-C3	-2.32	105.35	110.15
3	D	1	NAG	C1-O5-C5	2.31	115.28	112.19
3	D	4	MAN	C1-O5-C5	2.23	115.18	112.19
2	C	3	BMA	O2-C2-C3	-2.11	105.78	110.15

There are no chirality outliers.

All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	D	2	NAG	O5-C5-C6-O6
3	D	7	MAN	O5-C5-C6-O6
2	C	1	NAG	C4-C5-C6-O6
3	D	2	NAG	C8-C7-N2-C2
3	D	2	NAG	O7-C7-N2-C2
3	D	2	NAG	C4-C5-C6-O6
3	D	7	MAN	C4-C5-C6-O6
2	C	1	NAG	O5-C5-C6-O6
3	D	5	MAN	O5-C5-C6-O6
3	D	5	MAN	C4-C5-C6-O6
3	D	1	NAG	O5-C5-C6-O6
2	C	5	MAN	C4-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	B	611	-	3,3,3	0.45	0	2,2,2	0.24	0
6	PEG	A	610	-	6,6,6	0.86	0	5,5,5	0.26	0
4	PG0	B	608	-	7,7,7	0.47	0	6,6,6	0.36	0
5	EDO	A	607	-	3,3,3	0.43	0	2,2,2	0.39	0
5	EDO	A	608	-	3,3,3	0.49	0	2,2,2	0.22	0
6	PEG	B	612	-	6,6,6	0.84	0	5,5,5	0.21	0
4	PG0	A	606	-	7,7,7	0.50	0	6,6,6	0.84	0
7	AE3	B	609[A]	-	8,8,8	0.53	0	7,7,7	0.35	0
7	AE3	B	609[B]	-	8,8,8	0.51	0	7,7,7	0.38	0
5	EDO	A	609[A]	-	3,3,3	0.46	0	2,2,2	0.36	0
5	EDO	A	609[B]	-	3,3,3	0.42	0	2,2,2	0.39	0
5	EDO	B	610	-	3,3,3	0.42	0	2,2,2	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	B	611	-	-	1/1/1/1	-
6	PEG	A	610	-	-	1/4/4/4	-
4	PG0	B	608	-	-	2/5/5/5	-
5	EDO	A	607	-	-	1/1/1/1	-
5	EDO	A	608	-	-	1/1/1/1	-
6	PEG	B	612	-	-	1/4/4/4	-
4	PG0	A	606	-	-	3/5/5/5	-
7	AE3	B	609[A]	-	-	4/6/6/6	-
7	AE3	B	609[B]	-	-	2/6/6/6	-
5	EDO	A	609[A]	-	-	0/1/1/1	-
5	EDO	A	609[B]	-	-	1/1/1/1	-
5	EDO	B	610	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	606	PG0	O1-C3-C4-O2
5	A	607	EDO	O1-C1-C2-O2
5	B	610	EDO	O1-C1-C2-O2
5	B	611	EDO	O1-C1-C2-O2
7	B	609[B]	AE3	O2-C3-C4-O3
4	A	606	PG0	OTT-C1-C2-O1
4	A	606	PG0	C3-C4-O2-C5
7	B	609[B]	AE3	C6-C5-O3-C4
7	B	609[A]	AE3	C4-C3-O2-C2
6	A	610	PEG	C4-C3-O2-C2
4	B	608	PG0	O1-C3-C4-O2
7	B	609[A]	AE3	C1-C2-O2-C3
4	B	608	PG0	OTT-C1-C2-O1
5	A	608	EDO	O1-C1-C2-O2
5	A	609[B]	EDO	O1-C1-C2-O2
7	B	609[A]	AE3	O2-C3-C4-O3
6	B	612	PEG	O1-C1-C2-O2
7	B	609[A]	AE3	C6-C5-O3-C4

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	608	PG0	1	0
5	A	607	EDO	1	0
7	B	609[A]	AE3	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	318/327 (97%)	0.26	21 (6%) 24 28	13, 31, 72, 102	6 (1%)
1	B	322/327 (98%)	0.30	25 (7%) 19 22	13, 29, 69, 98	9 (2%)
All	All	640/654 (97%)	0.28	46 (7%) 21 25	13, 30, 70, 102	15 (2%)

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	321	PHE	4.9
1	B	225	ALA	4.8
1	A	365	PRO	4.1
1	B	320	THR	4.0
1	B	226	SER	3.9
1	B	367	LYS	3.8
1	B	501	SER	3.7
1	B	366	SER	3.7
1	A	423	PRO	3.5
1	B	257	SER	3.5
1	A	424	HIS	3.5
1	B	500	GLY	3.4
1	A	364	ALA	3.2
1	A	281	THR	3.2
1	A	422	HIS	3.2
1	B	319	HIS	3.2
1	A	421	THR	3.2
1	B	289	GLU	3.2
1	B	369	THR	3.2
1	B	545	PRO	3.1
1	B	426	PRO	3.0
1	A	372	LEU	2.9
1	A	425	LEU	2.9
1	B	227	ARG	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	426	PRO	2.8
1	A	367	LYS	2.8
1	B	318	GLY	2.7
1	B	277	VAL	2.7
1	A	398	THR	2.6
1	A	501	SER	2.6
1	A	397	LEU	2.5
1	A	389	GLU	2.5
1	B	368	GLY	2.4
1	A	393	ARG	2.4
1	B	425	LEU	2.4
1	B	229	PHE	2.3
1	A	437	SER	2.3
1	A	369	THR	2.2
1	B	258	GLY	2.2
1	B	424	HIS	2.2
1	B	264	ILE	2.2
1	B	263	THR	2.2
1	A	370	VAL	2.2
1	A	363	LEU	2.1
1	B	317	GLN	2.1
1	A	368	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MAN	D	5	11/12	0.57	0.15	77,82,84,85	0
3	MAN	D	7	11/12	0.60	0.17	67,75,82,83	0
2	MAN	C	4	11/12	0.74	0.14	64,71,76,80	0
2	MAN	C	5	11/12	0.77	0.20	65,79,82,84	0
3	MAN	D	6	11/12	0.82	0.16	48,62,65,68	0
3	MAN	D	4	11/12	0.84	0.12	59,63,65,68	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BMA	C	3	11/12	0.84	0.13	60,62,66,68	0
2	NAG	C	2	14/15	0.85	0.13	55,62,70,74	0
2	NAG	C	1	14/15	0.85	0.12	50,62,69,70	0
3	NAG	D	1	14/15	0.91	0.10	27,33,39,45	0
3	NAG	D	2	14/15	0.91	0.11	29,34,49,53	0
3	BMA	D	3	11/12	0.93	0.08	37,43,52,53	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	EDO	A	609[A]	4/4	0.54	0.24	46,50,52,53	4
5	EDO	A	609[B]	4/4	0.54	0.24	40,49,52,54	4
7	AE3	B	609[A]	9/9	0.56	0.28	40,47,52,52	9
7	AE3	B	609[B]	9/9	0.56	0.28	37,47,52,52	9
4	PG0	A	606	8/8	0.73	0.20	45,46,53,54	0
6	PEG	A	610	7/7	0.75	0.17	66,67,68,71	0
5	EDO	A	608	4/4	0.76	0.18	61,61,61,62	0
5	EDO	A	607	4/4	0.78	0.21	64,64,64,65	0
4	PG0	B	608	8/8	0.80	0.18	53,59,62,63	0
6	PEG	B	612	7/7	0.83	0.15	48,52,62,62	0
5	EDO	B	611	4/4	0.85	0.17	45,50,51,52	0
5	EDO	B	610	4/4	0.86	0.17	53,54,56,58	0

6.5 Other polymers [i](#)

There are no such residues in this entry.