



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 14, 2026 – 02:34 PM UTC

PDB ID : 7MPL / pdb\_00007mpl  
Title : Bartonella henselae NrnC bound to pGG  
Authors : Lormand, J.D.; Sondermann, H.  
Deposited on : 2021-05-04  
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)
RNA backbone	3983	1011 (2.20-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	207	97%
1	C	207	96%
1	E	207	96%
1	G	207	95%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	I	207	 93% 6%
1	K	207	 97%
1	M	207	 96%
1	O	207	 95%
2	B	2	 50% 50%
2	D	2	 50% 50%
2	F	2	 50% 50%
2	H	2	 50% 50%
2	J	2	 50% 50%
2	L	2	 50% 50%
2	N	2	 50% 50%
2	P	2	 50% 50%

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 15456 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NanoRNase C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	206	Total 1667	C 1049	N 295	O 316	S 7	0	6	0
1	C	205	Total 1660	C 1045	N 294	O 314	S 7	0	5	0
1	E	205	Total 1653	C 1040	N 296	O 311	S 6	0	3	0
1	G	205	Total 1655	C 1041	N 293	O 315	S 6	0	4	0
1	I	205	Total 1650	C 1038	N 294	O 312	S 6	0	3	0
1	K	205	Total 1652	C 1039	N 297	O 310	S 6	0	3	0
1	M	204	Total 1646	C 1036	N 295	O 309	S 6	0	3	0
1	O	204	Total 1631	C 1026	N 291	O 308	S 6	0	1	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP X5MEI1
C	0	SER	-	expression tag	UNP X5MEI1
E	0	SER	-	expression tag	UNP X5MEI1
G	0	SER	-	expression tag	UNP X5MEI1
I	0	SER	-	expression tag	UNP X5MEI1
K	0	SER	-	expression tag	UNP X5MEI1
M	0	SER	-	expression tag	UNP X5MEI1
O	0	SER	-	expression tag	UNP X5MEI1

- Molecule 2 is a RNA chain called 5'-phosphorylated GG.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
2	B	2	Total	C	N	O	P	0	0	0
			47	20	10	15	2			
2	D	2	Total	C	N	O	P	0	0	0
			47	20	10	15	2			
2	F	2	Total	C	N	O	P	0	0	0
			47	20	10	15	2			
2	H	2	Total	C	N	O	P	0	0	0
			47	20	10	15	2			
2	J	2	Total	C	N	O	P	0	0	0
			47	20	10	15	2			
2	L	2	Total	C	N	O	P	0	0	0
			47	20	10	15	2			
2	N	2	Total	C	N	O	P	0	0	0
			47	20	10	15	2			
2	P	2	Total	C	N	O	P	0	0	0
			47	20	10	15	2			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	263	Total	O	0	0
			263	263		
3	B	15	Total	O	0	0
			15	15		
3	C	265	Total	O	0	0
			265	265		
3	D	11	Total	O	0	0
			11	11		
3	E	237	Total	O	0	0
			237	237		
3	F	11	Total	O	0	0
			11	11		
3	G	249	Total	O	0	0
			249	249		
3	H	12	Total	O	0	0
			12	12		
3	I	223	Total	O	0	0
			223	223		
3	J	10	Total	O	0	0
			10	10		
3	K	213	Total	O	0	0
			213	213		
3	L	12	Total	O	0	0
			12	12		

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	M	179	Total 179	O 179	0	0
3	N	10	Total 10	O 10	0	0
3	O	143	Total 143	O 143	0	0
3	P	13	Total 13	O 13	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NanoRNase C




- Molecule 1: NanoRNase C



- Molecule 1: NanoRNase C



- Molecule 1: NanoRNase C




- Molecule 1: NanoRNase C



- Molecule 1: NanoRNase C



- Molecule 1: NanoRNase C



- Molecule 1: NanoRNase C



- Molecule 2: 5'-phosphorylated GG



- Molecule 2: 5'-phosphorylated GG



- Molecule 2: 5'-phosphorylated GG



- Molecule 2: 5'-phosphorylated GG



- Molecule 2: 5'-phosphorylated GG



G603  
G604

- Molecule 2: 5'-phosphorylated GG

Chain L:  50% 50%G603  
G604

- Molecule 2: 5'-phosphorylated GG

Chain N:  50% 50%G603  
G604

- Molecule 2: 5'-phosphorylated GG

Chain P:  50% 50%G603  
G604

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.16Å 128.26Å 129.28Å 90.00° 94.68° 90.00°	Depositor
Resolution (Å)	64.39 – 1.80 64.39 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.2 (64.39-1.80) 99.3 (64.39-1.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.37 (at 1.80Å)	Xtrriage
Refinement program	PHENIX 1.15.2_3472	Depositor
R, $R_{free}$	0.151 , 0.171 0.156 , 0.175	Depositor DCC
$R_{free}$ test set	10613 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.5	Xtrriage
Anisotropy	0.227	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 42.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	15456	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.34	0/1713	0.48	0/2314
1	C	0.30	0/1703	0.45	0/2300
1	E	0.29	0/1690	0.46	0/2282
1	G	0.29	0/1695	0.46	0/2290
1	I	0.27	0/1687	0.45	0/2279
1	K	0.25	0/1689	0.43	0/2281
1	M	0.25	0/1683	0.43	0/2272
1	O	0.21	0/1662	0.40	0/2245
2	B	0.33	0/52	0.76	0/78
2	D	0.41	0/52	0.72	0/78
2	F	0.36	0/52	0.73	0/78
2	H	0.33	0/52	0.71	0/78
2	J	0.31	0/52	0.67	0/78
2	L	0.33	0/52	0.65	0/78
2	N	0.35	0/52	0.74	0/78
2	P	0.29	0/52	0.62	0/78
All	All	0.28	0/13938	0.46	0/18887

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1667	0	1672	6	0
1	C	1660	0	1664	5	0
1	E	1653	0	1658	5	0
1	G	1655	0	1653	7	0
1	I	1650	0	1651	8	0
1	K	1652	0	1658	4	0
1	M	1646	0	1651	3	0
1	O	1631	0	1628	5	0
2	B	47	0	23	1	0
2	D	47	0	23	1	0
2	F	47	0	23	1	0
2	H	47	0	23	1	0
2	J	47	0	23	1	0
2	L	47	0	23	1	0
2	N	47	0	23	1	0
2	P	47	0	23	1	0
3	A	263	0	0	2	1
3	B	15	0	0	1	0
3	C	265	0	0	1	0
3	D	11	0	0	1	0
3	E	237	0	0	3	1
3	F	11	0	0	1	0
3	G	249	0	0	3	0
3	H	12	0	0	1	0
3	I	223	0	0	3	0
3	J	10	0	0	1	0
3	K	213	0	0	1	0
3	L	12	0	0	1	0
3	M	179	0	0	1	0
3	N	10	0	0	1	0
3	O	143	0	0	1	0
3	P	13	0	0	1	0
All	All	15456	0	13419	40	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 40 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:178:VAL:HG22	1:O:188:MET:HE1	1.67	0.77
1:A:188[B]:MET:HE1	1:G:178:VAL:HG22	1.75	0.69

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:164:ASP:OD1	3:G:301:HOH:O	2.09	0.69
1:C:188[B]:MET:HE1	1:K:178:VAL:HG22	1.73	0.68
2:H:604:G:OP1	3:H:701:HOH:O	2.10	0.68

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1350:HOH:O	3:E:351:HOH:O[2_655]	2.14	0.06

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	210/207 (101%)	209 (100%)	1 (0%)	0	100	100
1	C	208/207 (100%)	207 (100%)	1 (0%)	0	100	100
1	E	206/207 (100%)	205 (100%)	1 (0%)	0	100	100
1	G	207/207 (100%)	206 (100%)	1 (0%)	0	100	100
1	I	206/207 (100%)	205 (100%)	1 (0%)	0	100	100
1	K	206/207 (100%)	205 (100%)	1 (0%)	0	100	100
1	M	205/207 (99%)	204 (100%)	1 (0%)	0	100	100
1	O	203/207 (98%)	202 (100%)	1 (0%)	0	100	100
All	All	1651/1656 (100%)	1643 (100%)	8 (0%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/180 (102%)	183 (100%)	1 (0%)	81	80
1	C	183/180 (102%)	182 (100%)	1 (0%)	81	80
1	E	181/180 (101%)	180 (99%)	1 (1%)	78	77
1	G	182/180 (101%)	180 (99%)	2 (1%)	65	60
1	I	181/180 (101%)	180 (99%)	1 (1%)	78	77
1	K	181/180 (101%)	180 (99%)	1 (1%)	78	77
1	M	180/180 (100%)	178 (99%)	2 (1%)	65	60
1	O	178/180 (99%)	177 (99%)	1 (1%)	78	77
All	All	1450/1440 (101%)	1440 (99%)	10 (1%)	76	73

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	M	42	GLN
1	M	127	ASN
1	O	42	GLN
1	G	42	GLN
1	G	68	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	42	GLN
1	M	134	GLN
1	G	135	GLN
1	O	8	GLN
1	M	8	GLN

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	1/2 (50%)	0	0
2	D	1/2 (50%)	0	0
2	F	1/2 (50%)	0	0
2	H	1/2 (50%)	0	0
2	J	1/2 (50%)	0	0
2	L	1/2 (50%)	0	0
2	N	1/2 (50%)	0	0
2	P	1/2 (50%)	0	0
All	All	8/16 (50%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

#### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

#### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

#### 5.7 Other polymers [i](#)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	206/207 (99%)	-0.49	2 (0%) 79 80	16, 27, 45, 77	6 (2%)
1	C	205/207 (99%)	-0.44	2 (0%) 79 80	17, 29, 45, 74	5 (2%)
1	E	205/207 (99%)	-0.37	0 100 100	19, 32, 47, 60	3 (1%)
1	G	205/207 (99%)	-0.37	1 (0%) 87 88	17, 32, 47, 59	4 (1%)
1	I	205/207 (99%)	-0.28	1 (0%) 87 88	22, 34, 48, 88	3 (1%)
1	K	205/207 (99%)	-0.18	2 (0%) 79 80	21, 38, 54, 71	3 (1%)
1	M	204/207 (98%)	-0.15	1 (0%) 87 88	23, 39, 59, 74	3 (1%)
1	O	204/207 (98%)	0.03	1 (0%) 87 88	28, 45, 66, 81	1 (0%)
2	B	2/2 (100%)	-1.15	0 100 100	26, 26, 26, 27	0
2	D	2/2 (100%)	-1.06	0 100 100	28, 28, 28, 28	0
2	F	2/2 (100%)	-0.87	0 100 100	31, 31, 31, 31	0
2	H	2/2 (100%)	-0.94	0 100 100	29, 29, 29, 30	0
2	J	2/2 (100%)	-0.94	0 100 100	32, 32, 32, 32	0
2	L	2/2 (100%)	-0.67	0 100 100	34, 34, 34, 36	0
2	N	2/2 (100%)	-0.69	0 100 100	31, 31, 31, 32	0
2	P	2/2 (100%)	-0.46	0 100 100	38, 38, 38, 38	0
All	All	1655/1672 (98%)	-0.29	10 (0%) 85 86	16, 34, 54, 88	28 (1%)

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	2	THR	7.0
1	C	2	THR	4.9
1	A	1	MET	3.4
1	A	146	ARG	3.3
1	C	3	GLU	3.2

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.