



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 18, 2026 – 05:22 PM UTC

PDB ID : 4MQF / pdb_00004mqf
Title : Crystal structure of the extracellular domain of human GABA(B) receptor bound to the antagonist 2-hydroxysaclofen
Authors : Geng, Y.; Bush, M.; Mosyak, L.; Wang, F.; Fan, Q.R.
Deposited on : 2013-09-16
Resolution : 2.22 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7682 (2.24-2.20)
Clashscore	190562	8402 (2.24-2.20)
Ramachandran outliers	187476	8303 (2.24-2.20)
Sidechain outliers	187428	8304 (2.24-2.20)
RSRZ outliers	180081	7683 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	420	12% 79% 17% ..
2	B	433	5% 84% 8% 7%
3	C	6	100%
4	D	4	25% 75%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7070 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gamma-aminobutyric acid type B receptor subunit 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	406	3262	2084	549	615	14	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	460	ASP	-	expression tag	UNP Q9UBS5
A	461	TYR	-	expression tag	UNP Q9UBS5
A	462	LYS	-	expression tag	UNP Q9UBS5
A	463	ASP	-	expression tag	UNP Q9UBS5
A	464	ASP	-	expression tag	UNP Q9UBS5
A	465	ASP	-	expression tag	UNP Q9UBS5
A	466	ASP	-	expression tag	UNP Q9UBS5
A	467	LYS	-	expression tag	UNP Q9UBS5


- Molecule 2 is a protein called Gamma-aminobutyric acid type B receptor subunit 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	403	3214	2052	543	604	15	0	0	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	467	ASP	-	expression tag	UNP O75899
B	468	TYR	-	expression tag	UNP O75899
B	469	LYS	-	expression tag	UNP O75899
B	470	ASP	-	expression tag	UNP O75899
B	471	ASP	-	expression tag	UNP O75899
B	472	ASP	-	expression tag	UNP O75899
B	473	ASP	-	expression tag	UNP O75899
B	474	LYS	-	expression tag	UNP O75899

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	6	Total	C	N	O	0	0	0
			71	40	2	29			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	4	Total	C	N	O	0	0	0
			49	28	2	19			

- Molecule 5 is 2-hydroxysaclofen (CCD ID: 2BQ) (formula: C₉H₁₂ClNO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	Cl	N	O	S	
			16	9	1	1	4	1	
								0	0

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O		
			14	8	1	5		
							0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	201	Total	O		
			201	201	0	0
7	B	243	Total	O		
			243	243	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Gamma-aminobutyric acid type B receptor subunit 1



- Molecule 2: Gamma-aminobutyric acid type B receptor subunit 2



- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



MAG1
MAG2
MAN3
MAN4
MAN5
FUC6

- Molecule 4: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  25% 75%

MAG1
MAG2
MAN3
FUC4

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	70.28Å 112.63Å 73.22Å 90.00° 97.79° 90.00°	Depositor
Resolution (Å)	32.50 – 2.22 32.50 – 2.22	Depositor EDS
% Data completeness (in resolution range)	90.5 (32.50-2.22) 88.6 (32.50-2.22)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.04 (at 2.22Å)	Xtriage
Refinement program	BUSTER-TNT, BUSTER	Depositor
R, R_{free}	0.207 , 0.234 0.206 , 0.237	Depositor DCC
R_{free} test set	2538 reflections (4.56%)	wwPDB-VP
Wilson B-factor (Å ²)	27.6	Xtriage
Anisotropy	0.271	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 51.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.018 for l,-k,h	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7070	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, MAN, 2BQ, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	1/3345 (0.0%)	1.29	26/4536 (0.6%)
2	B	0.84	1/3287 (0.0%)	1.27	8/4449 (0.2%)
All	All	0.84	2/6632 (0.0%)	1.28	34/8985 (0.4%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	61	MET	SD-CE	-8.10	1.59	1.79
2	B	129	MET	SD-CE	-7.29	1.61	1.79

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	389	ASN	N-CA-C	-8.58	96.68	109.81
1	A	258	PHE	CA-CB-CG	-7.06	106.74	113.80
2	B	349	VAL	N-CA-CB	6.41	118.19	110.31
2	B	390	TYR	CA-C-N	6.05	130.92	120.58
2	B	390	TYR	C-N-CA	6.05	130.92	120.58
1	A	454	GLY	N-CA-C	-6.04	106.76	115.27
1	A	197	GLN	N-CA-C	-6.01	101.59	110.48
1	A	326	SER	CA-C-N	5.99	128.30	120.28
1	A	326	SER	C-N-CA	5.99	128.30	120.28
1	A	411	ASP	CA-CB-CG	5.98	118.58	112.60
1	A	385	ASN	CA-C-N	5.80	131.29	121.14
1	A	385	ASN	C-N-CA	5.80	131.29	121.14
1	A	458	ALA	CA-C-N	5.76	130.34	120.72
1	A	458	ALA	C-N-CA	5.76	130.34	120.72
1	A	434	GLY	N-CA-C	5.59	118.31	110.38
1	A	201	VAL	N-CA-CB	5.43	116.90	110.55
1	A	106	GLY	CA-C-N	5.41	127.53	120.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	106	GLY	C-N-CA	5.41	127.53	120.28
1	A	337	LEU	CA-C-N	5.40	127.52	120.28
1	A	337	LEU	C-N-CA	5.40	127.52	120.28
2	B	256	ASP	CA-CB-CG	5.37	117.97	112.60
1	A	159	ASN	CA-CB-CG	5.31	117.91	112.60
1	A	139	ALA	CA-C-N	5.30	127.64	120.38
1	A	139	ALA	C-N-CA	5.30	127.64	120.38
2	B	105	ASP	CA-CB-CG	5.28	117.88	112.60
1	A	299	THR	N-CA-C	-5.24	105.75	111.82
1	A	259	CYS	N-CA-C	-5.21	105.51	111.14
1	A	54	TYR	N-CA-C	5.21	117.98	109.59
2	B	360	ASP	CA-C-N	5.15	125.69	119.98
2	B	360	ASP	C-N-CA	5.15	125.69	119.98
1	A	299	THR	CA-C-N	5.06	127.33	120.44
1	A	299	THR	C-N-CA	5.06	127.33	120.44
1	A	158	SER	CA-C-N	5.04	130.74	123.13
1	A	158	SER	C-N-CA	5.04	130.74	123.13

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3262	0	3180	35	0
2	B	3214	0	3161	21	0
3	C	71	0	61	0	0
4	D	49	0	43	2	0
5	A	16	0	11	0	0
6	A	14	0	13	0	0
7	A	201	0	0	4	0
7	B	243	0	0	1	0
All	All	7070	0	6469	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:234:VAL:HG21	1:A:260:GLU:HB3	1.65	0.78
1:A:141:ARG:HH11	1:A:142:MET:HE3	1.52	0.75
2:B:86:GLU:HA	2:B:89:ARG:HH21	1.57	0.69
2:B:373:MET:HG2	2:B:385:ILE:HD11	1.75	0.68
2:B:388:PHE:HB2	2:B:395:LEU:HD23	1.80	0.64
2:B:404:ASN:HD22	4:D:1:NAG:H83	1.63	0.63
1:A:238:LYS:HD2	1:A:264:GLU:HB3	1.82	0.62
1:A:53:VAL:HG13	1:A:123:ILE:HG12	1.82	0.61
2:B:320:GLU:HG2	2:B:427:LYS:HB2	1.83	0.61
2:B:126:ASN:HD21	2:B:388:PHE:HD2	1.49	0.59
1:A:269:LYS:HE2	1:A:452:ILE:HG21	1.85	0.58
1:A:283:TRP:HA	1:A:286:ILE:HD12	1.87	0.56
2:B:206:GLN:HE21	2:B:210:GLU:HG2	1.71	0.56
2:B:373:MET:HG2	2:B:385:ILE:CD1	2.35	0.56
1:A:191:LYS:HB2	1:A:459:ASP:HB2	1.87	0.56
1:A:302:VAL:O	1:A:305:HIS:HD2	1.90	0.55
2:B:89:ARG:HG3	7:B:650:HOH:O	2.06	0.55
2:B:346:ARG:O	2:B:349:VAL:HG22	2.07	0.54
2:B:129:MET:HB2	2:B:152:VAL:O	2.07	0.53
1:A:269:LYS:HE2	1:A:452:ILE:CG2	2.39	0.52
2:B:86:GLU:CA	2:B:89:ARG:HH21	2.21	0.52
4:D:2:NAG:O3	4:D:3:MAN:H2	2.08	0.52
1:A:305:HIS:CE1	1:A:430:TYR:OH	2.62	0.51
2:B:376:LEU:HD12	2:B:385:ILE:HG12	1.91	0.51
1:A:238:LYS:HE3	1:A:264:GLU:O	2.13	0.49
1:A:61:MET:HE2	7:A:729:HOH:O	2.12	0.49
1:A:48:SER:HB2	7:A:725:HOH:O	2.13	0.48
1:A:305:HIS:HE1	1:A:430:TYR:OH	1.95	0.48
2:B:150:ASN:ND2	2:B:393:HIS:HA	2.29	0.48
1:A:85:ARG:HB3	1:A:88:ILE:HD12	1.94	0.48
1:A:141:ARG:NH1	1:A:142:MET:HE3	2.25	0.47
1:A:243:ARG:HB2	1:A:459:ASP:HA	1.95	0.47
1:A:255:ARG:HB3	1:A:291:ILE:HD13	1.95	0.47
1:A:409:VAL:HG23	1:A:417:MET:HB2	1.95	0.47
1:A:351:PRO:HB3	1:A:404:VAL:HG23	1.97	0.47
1:A:224:GLN:HB3	1:A:233:PRO:HB3	1.97	0.46
1:A:98:HIS:HD2	7:A:729:HOH:O	1.99	0.46
1:A:302:VAL:O	1:A:305:HIS:CD2	2.67	0.46
2:B:389:ASN:C	2:B:391:THR:H	2.23	0.46
1:A:132:VAL:O	1:A:136:VAL:HG13	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:327:GLN:H	1:A:327:GLN:CD	2.24	0.44
1:A:269:LYS:HG2	1:A:452:ILE:HB	1.99	0.44
2:B:86:GLU:HG3	2:B:89:ARG:NH2	2.33	0.44
2:B:61:LEU:HD13	2:B:142:ILE:HD13	2.00	0.44
1:A:85:ARG:CB	1:A:88:ILE:HD12	2.48	0.43
1:A:226:PHE:HB3	1:A:233:PRO:HG2	2.00	0.43
1:A:297:GLU:O	1:A:300:GLU:HB3	2.19	0.43
1:A:175:LEU:HD23	1:A:310:ILE:HG12	2.00	0.43
2:B:160:THR:HA	2:B:161:PRO:HD3	1.94	0.43
2:B:371:ARG:HD2	2:B:405:GLU:OE1	2.19	0.43
2:B:146:LEU:HB3	2:B:151:LEU:O	2.19	0.42
1:A:181:VAL:HG21	7:A:723:HOH:O	2.19	0.42
1:A:230:PRO:C	1:A:233:PRO:HD2	2.45	0.41
1:A:321:ILE:HD13	1:A:362:LEU:HD11	2.02	0.41
1:A:124:ILE:CD1	1:A:360:LEU:HD13	2.50	0.41
2:B:325:LYS:HB3	2:B:327:ILE:HD12	2.03	0.41
1:A:255:ARG:HB3	1:A:291:ILE:CD1	2.51	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	402/420 (96%)	390 (97%)	11 (3%)	1 (0%)	43	50
2	B	397/433 (92%)	384 (97%)	13 (3%)	0	100	100
All	All	799/853 (94%)	774 (97%)	24 (3%)	1 (0%)	48	56

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	457	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	353/362 (98%)	338 (96%)	15 (4%)	26	34
2	B	349/375 (93%)	343 (98%)	6 (2%)	53	68
All	All	702/737 (95%)	681 (97%)	21 (3%)	36	47

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	ARG
1	A	82	VAL
1	A	85	ARG
1	A	114	GLU
1	A	136	VAL
1	A	232	VAL
1	A	239	ARG
1	A	245	ILE
1	A	249	PHE
1	A	276	ILE
1	A	336	ARG
1	A	368	SER
1	A	425	LEU
1	A	431	LYS
1	A	440	LYS
2	B	66	LYS
2	B	188	LYS
2	B	204	ASP
2	B	320	GLU
2	B	437	VAL
2	B	451	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	72	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	98	HIS
1	A	144	ASN
1	A	305	HIS
1	A	407	HIS
2	B	126	ASN
2	B	150	ASN
2	B	206	GLN
2	B	306	ASN
2	B	337	GLN
2	B	414	GLN
2	B	432	GLN
2	B	459	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	C	1	1,3	14,14,15	0.29	0	17,19,21	2.54	2 (11%)
3	NAG	C	2	3	14,14,15	0.32	0	17,19,21	2.52	4 (23%)
3	MAN	C	3	3	11,11,12	0.75	0	15,15,17	1.84	2 (13%)
3	MAN	C	4	3	11,11,12	0.51	0	15,15,17	1.52	2 (13%)
3	MAN	C	5	3	11,11,12	0.73	0	15,15,17	2.04	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	FUC	C	6	3	10,10,11	0.66	0	14,14,16	1.54	2 (14%)
4	NAG	D	1	2,4	14,14,15	0.35	0	17,19,21	1.05	1 (5%)
4	NAG	D	2	4	14,14,15	0.34	0	17,19,21	3.19	6 (35%)
4	MAN	D	3	4	11,11,12	0.59	0	15,15,17	1.42	3 (20%)
4	FUC	D	4	4	10,10,11	0.48	0	14,14,16	0.69	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	3/6/23/26	0/1/1/1
3	NAG	C	2	3	-	2/6/23/26	0/1/1/1
3	MAN	C	3	3	-	2/2/19/22	0/1/1/1
3	MAN	C	4	3	-	1/2/19/22	0/1/1/1
3	MAN	C	5	3	-	0/2/19/22	0/1/1/1
3	FUC	C	6	3	-	-	0/1/1/1
4	NAG	D	1	2,4	-	2/6/23/26	0/1/1/1
4	NAG	D	2	4	-	2/6/23/26	0/1/1/1
4	MAN	D	3	4	-	1/2/19/22	0/1/1/1
4	FUC	D	4	4	-	-	0/1/1/1

There are no bond length outliers.

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1	NAG	C1-O5-C5	9.58	125.03	112.19
4	D	2	NAG	O5-C1-C2	8.32	124.17	111.29
3	C	5	MAN	C1-O5-C5	7.14	121.75	112.19
3	C	2	NAG	C1-O5-C5	6.77	121.26	112.19
3	C	2	NAG	O5-C1-C2	6.38	121.16	111.29
4	D	2	NAG	C1-O5-C5	6.28	120.61	112.19
4	D	2	NAG	C1-C2-N2	-5.99	101.00	110.43
3	C	3	MAN	C1-O5-C5	4.50	118.22	112.19
3	C	4	MAN	C1-C2-C3	-4.30	103.38	109.64
3	C	6	FUC	C1-O5-C5	3.93	122.24	112.97
3	C	6	FUC	C1-C2-C3	3.79	115.17	109.64
3	C	3	MAN	C1-C2-C3	3.59	114.87	109.64
4	D	1	NAG	C1-C2-N2	3.46	115.89	110.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	3	MAN	C1-C2-C3	3.34	114.51	109.64
3	C	4	MAN	C3-C4-C5	2.92	115.53	110.23
4	D	3	MAN	C1-O5-C5	2.57	115.63	112.19
4	D	2	NAG	C3-C4-C5	2.56	114.86	110.23
3	C	5	MAN	C1-C2-C3	2.49	113.27	109.64
3	C	2	NAG	C1-C2-N2	-2.49	106.51	110.43
4	D	2	NAG	O4-C4-C3	-2.35	104.83	110.38
4	D	3	MAN	C2-C3-C4	2.21	114.75	110.86
3	C	1	NAG	O5-C1-C2	2.14	114.60	111.29
4	D	2	NAG	O4-C4-C5	-2.08	104.20	109.32
3	C	2	NAG	O5-C5-C4	2.01	115.73	110.83

There are no chirality outliers.

All (13) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	3	MAN	O5-C5-C6-O6
3	C	3	MAN	C4-C5-C6-O6
3	C	1	NAG	C8-C7-N2-C2
3	C	1	NAG	O7-C7-N2-C2
4	D	1	NAG	C8-C7-N2-C2
4	D	1	NAG	O7-C7-N2-C2
3	C	2	NAG	O5-C5-C6-O6
4	D	2	NAG	C4-C5-C6-O6
3	C	4	MAN	O5-C5-C6-O6
3	C	1	NAG	O5-C5-C6-O6
4	D	3	MAN	O5-C5-C6-O6
3	C	2	NAG	C4-C5-C6-O6
4	D	2	NAG	O5-C5-C6-O6


There are no ring outliers.

3 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	3	MAN	1	0
4	D	1	NAG	1	0
4	D	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	2BQ	A	501	-	14,16,16	2.02	3 (21%)	22,24,24	1.09	1 (4%)
6	NAG	A	508	1	14,14,15	0.28	0	17,19,21	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	2BQ	A	501	-	-	1/13/15/15	0/1/1/1
6	NAG	A	508	1	-	2/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	501	2BQ	O3-S	5.63	1.61	1.45
5	A	501	2BQ	C9-S	-3.41	1.68	1.77
5	A	501	2BQ	C7-C6	-2.32	1.50	1.53

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	501	2BQ	O1-S-C9	3.38	112.62	105.39

There are no chirality outliers.


All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	508	NAG	C4-C5-C6-O6
6	A	508	NAG	O5-C5-C6-O6
5	A	501	2BQ	C1-C6-C7-C8

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	406/420 (96%)	0.57	51 (12%) 8 6	14, 35, 72, 87	0
2	B	403/433 (93%)	0.13	23 (5%) 29 26	14, 31, 64, 96	0
All	All	809/853 (94%)	0.35	74 (9%) 15 12	14, 33, 70, 96	0

All (74) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	461	TYR	6.2
2	B	89	ARG	5.6
2	B	300	SER	5.3
2	B	379	SER	5.1
1	A	368	SER	5.1
2	B	388	PHE	5.1
1	A	377	ARG	4.9
1	A	222	PHE	4.8
2	B	385	ILE	4.0
2	B	378	ALA	3.9
2	B	387	ASP	3.8
1	A	223	ARG	3.4
2	B	326	GLN	3.4
1	A	292	ASN	3.2
1	A	444	SER	3.2
1	A	221	THR	3.1
1	A	445	TRP	3.1
1	A	266	LEU	3.0
1	A	198	THR	3.0
1	A	236	ASN	2.9
1	A	452	ILE	2.8
2	B	377	HIS	2.8
1	A	270	LYS	2.8
2	B	301	ARG	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	227	PHE	2.7
2	B	292	GLN	2.7
1	A	431	LYS	2.7
1	A	450	LYS	2.7
1	A	212	ARG	2.7
1	A	448	THR	2.7
2	B	245	GLY	2.7
2	B	371	ARG	2.7
1	A	224	GLN	2.6
2	B	302	CYS	2.6
1	A	449	ASP	2.6
1	A	235	LYS	2.5
1	A	263	LYS	2.5
1	A	265	ARG	2.4
2	B	68	VAL	2.4
1	A	220	ILE	2.4
1	A	454	GLY	2.4
1	A	158	SER	2.4
2	B	305	LYS	2.3
1	A	458	ALA	2.3
2	B	327	ILE	2.3
1	A	86	ARG	2.3
1	A	262	TYR	2.3
2	B	304	ARG	2.3
1	A	272	VAL	2.3
1	A	261	VAL	2.2
1	A	340	HIS	2.2
1	A	267	PHE	2.2
2	B	223	ASP	2.2
1	A	90	PRO	2.2
1	A	447	LYS	2.2
1	A	231	ALA	2.2
1	A	455	SER	2.2
2	B	53	SER	2.2
2	B	231	SER	2.2
1	A	269	LYS	2.1
2	B	287	PRO	2.1
1	A	240	GLN	2.1
1	A	192	ILE	2.1
1	A	451	TRP	2.1
1	A	129	CYS	2.1
1	A	294	THR	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	271	TYR	2.1
1	A	457	PRO	2.1
1	A	327	GLN	2.1
2	B	234	ASN	2.1
1	A	89	LEU	2.0
1	A	237	LEU	2.0
1	A	239	ARG	2.0
1	A	244	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	MAN	D	3	11/12	0.41	0.16	94,98,99,100	0
3	MAN	C	4	11/12	0.49	0.33	90,91,95,96	0
3	MAN	C	5	11/12	0.62	0.23	83,85,88,89	0
3	FUC	C	6	10/11	0.66	0.28	84,89,90,91	0
4	NAG	D	1	14/15	0.78	0.16	54,61,69,70	0
3	MAN	C	3	11/12	0.79	0.27	80,83,84,88	0
3	NAG	C	2	14/15	0.81	0.21	65,69,72,76	0
3	NAG	C	1	14/15	0.82	0.15	55,59,70,77	0
4	NAG	D	2	14/15	0.84	0.14	75,79,82,88	0
4	FUC	D	4	10/11	0.87	0.16	67,67,69,69	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	NAG	A	508	14/15	0.72	0.16	72,77,79,80	0
5	2BQ	A	501	16/16	0.98	0.06	13,19,23,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.