



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 08:30 AM UTC

PDB ID : 6MQR / pdb_00006mqr
Title : Vaccine-elicited NHP FP-targeting neutralizing antibody 0PV-a.01 in complex with FP (residue 512-519)
Authors : Xu, K.; Wang, Y.; Kwong, P.D.
Deposited on : 2018-10-10
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	228	3% 86% 12%
2	L	214	82% 16%
3	A	8	12% 88% 12%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SO4	H	301	-	-	X	-
4	SO4	L	301	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3562 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called antibody 0PV-A.01 Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	227	Total	C	N	O	S	0	0	0
			1727	1090	288	342	7			


- Molecule 2 is a protein called antibody 0PV-A.01 Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	213	Total	C	N	O	S	0	0	0
			1642	1021	279	336	6			

- Molecule 3 is a protein called HIV fusion peptide residue 512-519.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	A	8	Total	C	N	O	0	0	0
			51	35	8	8			

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	H	1	Total O S 5 4 1	0	0
4	L	1	Total O S 5 4 1	0	0

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	L	1	Total Ca 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	H	58	Total O 58 58	0	0
6	L	70	Total O 70 70	0	0
6	A	3	Total O 3 3	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: antibody 0PV-A.01 Fab heavy chain



- Molecule 2: antibody 0PV-A.01 Fab light chain



- Molecule 3: HIV fusion peptide residue 512-519



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 3	Depositor
Cell constants a, b, c, α , β , γ	159.25Å 159.25Å 159.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.56 – 2.45 42.56 – 2.45	Depositor EDS
% Data completeness (in resolution range)	99.6 (42.56-2.45) 99.6 (42.56-2.45)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 2.45Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.175 , 0.208 0.180 , 0.191	Depositor DCC
R_{free} test set	1148 reflections (4.62%)	wwPDB-VP
Wilson B-factor (Å ²)	50.3	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 21.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.024 for -l,-k,-h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3562	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	H	0.73	3/1770 (0.2%)	0.81	1/2420 (0.0%)
2	L	1.09	12/1676 (0.7%)	0.93	10/2274 (0.4%)
3	A	0.36	0/51	0.87	0/68
All	All	0.92	15/3497 (0.4%)	0.87	11/4762 (0.2%)

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	L	185	GLU	CA-CB	9.01	1.67	1.53
2	L	130	VAL	C-O	-8.05	1.15	1.24
1	H	111	VAL	C-O	-7.95	1.15	1.24
2	L	192	TYR	C-O	-7.59	1.14	1.24
2	L	186	TYR	C-O	-7.44	1.15	1.24
2	L	185	GLU	C-O	-7.02	1.15	1.24
2	L	129	THR	C-O	-6.17	1.16	1.23
2	L	191	VAL	C-O	-6.12	1.17	1.24
2	L	210	ASN	C-O	-5.82	1.17	1.24
2	L	187	GLN	C-O	-5.82	1.16	1.24
1	H	85	THR	C-O	-5.28	1.17	1.24
2	L	209	PHE	C-O	-5.26	1.17	1.23
1	H	83	THR	C-O	-5.13	1.19	1.24
2	L	184	THR	C-O	-5.06	1.18	1.24
2	L	181	LEU	CA-C	-5.02	1.46	1.52

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	L	185	GLU	N-CA-CB	9.37	124.02	110.16
2	L	185	GLU	CB-CG-CD	9.33	128.46	112.60
2	L	185	GLU	OE1-CD-OE2	-9.10	101.06	122.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	85	THR	N-CA-C	7.66	122.21	113.01
2	L	185	GLU	CG-CD-OE2	6.52	133.40	118.40
2	L	184	THR	CA-C-O	-6.33	113.85	120.55
2	L	185	GLU	CA-CB-CG	5.75	125.59	114.10
2	L	184	THR	N-CA-C	5.48	117.25	111.28
2	L	190	LYS	N-CA-C	5.39	119.85	112.68
2	L	130	VAL	N-CA-C	5.18	116.82	108.85
2	L	185	GLU	CB-CA-C	5.01	119.36	110.85

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1727	0	1692	19	1
2	L	1642	0	1587	22	1
3	A	51	0	53	2	0
4	H	5	0	0	3	0
4	L	5	0	0	3	0
5	L	1	0	0	0	0
6	A	3	0	0	0	0
6	H	58	0	0	3	1
6	L	70	0	0	6	1
All	All	3562	0	3332	41	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:39:LYS:NZ	6:L:401:HOH:O	1.98	0.82
1:H:77:HIS:NE2	4:H:301:SO4:S	2.63	0.69
2:L:7:SER:OG	6:L:402:HOH:O	2.15	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:50:HIS:O	2:L:50:HIS:ND1	2.31	0.62
2:L:202:SER:OG	6:L:403:HOH:O	2.16	0.62
1:H:191:THR:O	6:H:401:HOH:O	2.16	0.61
2:L:24:ARG:NH2	6:L:402:HOH:O	2.35	0.59
1:H:143:LYS:NZ	6:H:402:HOH:O	2.19	0.58
2:L:49:HIS:NE2	4:L:301:SO4:S	2.66	0.56
1:H:57:VAL:HG22	1:H:59:TYR:CE2	2.41	0.56
2:L:45:ARG:NH1	6:L:405:HOH:O	2.29	0.55
2:L:4:MET:HE3	2:L:23:CYS:SG	2.46	0.55
2:L:49:HIS:NE2	4:L:301:SO4:O2	2.39	0.54
1:H:131:THR:HG23	1:H:133:GLU:H	1.74	0.53
1:H:77:HIS:NE2	4:H:301:SO4:O3	2.42	0.53
1:H:143:LYS:CE	6:H:402:HOH:O	2.57	0.53
1:H:61:PRO:HA	1:H:64:GLN:OE1	2.10	0.51
2:L:91:TYR:O	3:A:513:VAL:HG12	2.11	0.51
2:L:18:ARG:HG2	2:L:18:ARG:HH11	1.78	0.49
2:L:122:GLU:HG3	2:L:126:LYS:HE2	1.95	0.49
1:H:96:ARG:HD3	1:H:100(D):GLY:O	2.13	0.48
2:L:28:ASP:OD1	2:L:68:GLY:HA2	2.12	0.48
1:H:119:PRO:HB3	1:H:145:TYR:HB3	1.96	0.48
1:H:131:THR:HG22	1:H:134:SER:HB3	1.95	0.47
2:L:49:HIS:NE2	4:L:301:SO4:O1	2.46	0.47
2:L:39:LYS:HE2	2:L:81:GLU:O	2.14	0.47
1:H:77:HIS:NE2	4:H:301:SO4:O4	2.47	0.45
1:H:72:ASP:OD2	1:H:75:LYS:HD2	2.17	0.44
1:H:143:LYS:HB2	1:H:143:LYS:HE2	1.74	0.44
1:H:12:MET:HE3	1:H:12:MET:HB3	1.54	0.44
1:H:63:LEU:O	1:H:67:VAL:HG13	2.17	0.43
1:H:2:VAL:HG12	1:H:102:LEU:HD11	2.01	0.43
2:L:159:SER:HA	2:L:178:THR:O	2.18	0.43
2:L:12:SER:OG	2:L:105:GLU:OE1	2.28	0.42
1:H:29:ILE:HD12	1:H:34:TRP:CH2	2.55	0.42
2:L:24:ARG:HG2	2:L:24:ARG:HH21	1.86	0.41
2:L:94:PHE:CE2	3:A:513:VAL:CG2	3.04	0.41
2:L:212:GLY:C	2:L:213:GLU:HG3	2.46	0.40
1:H:126:PRO:HG2	1:H:213:ILE:HD12	2.03	0.40
2:L:144:ALA:O	6:L:404:HOH:O	2.21	0.40
2:L:79:GLN:HB3	2:L:81:GLU:CD	2.46	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:84:ASP:CB	2:L:185:GLU:OE2[2_655]	1.84	0.36
6:H:456:HOH:O	6:L:428:HOH:O[23_555]	1.95	0.25

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	225/228 (99%)	221 (98%)	4 (2%)	0	100	100
2	L	211/214 (99%)	206 (98%)	5 (2%)	0	100	100
3	A	6/8 (75%)	6 (100%)	0	0	100	100
All	All	442/450 (98%)	433 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	201/201 (100%)	198 (98%)	3 (2%)	57	70
2	L	189/190 (100%)	185 (98%)	4 (2%)	47	62
3	A	4/4 (100%)	4 (100%)	0	100	100
All	All	394/395 (100%)	387 (98%)	7 (2%)	51	65

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	57	VAL
1	H	67	VAL
1	H	212	GLU
2	L	124	GLN
2	L	129	THR
2	L	185	GLU
2	L	190	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	171	GLN
2	L	37	GLN
2	L	138	ASN
2	L	158	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	L	301	-	4,4,4	0.24	0	6,6,6	0.14	0
4	SO4	H	301	-	4,4,4	0.24	0	6,6,6	0.08	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	L	301	SO4	3	0
4	H	301	SO4	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	227/228 (99%)	-0.27	6 (2%) 57 58	32, 46, 73, 111	0
2	L	213/214 (99%)	-0.30	1 (0%) 87 88	34, 48, 65, 78	0
3	A	8/8 (100%)	0.21	1 (12%) 8 6	49, 50, 67, 75	0
All	All	448/450 (99%)	-0.28	8 (1%) 67 69	32, 47, 70, 111	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	84	ASP	2.9
1	H	100(B)	TYR	2.7
1	H	216	CYS	2.7
2	L	1	ASP	2.5
1	H	214	LYS	2.3
1	H	127	SER	2.2
3	A	519	PHE	2.2
1	H	215	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SO4	L	301	5/5	0.67	0.20	60,60,62,65	0
4	SO4	H	301	5/5	0.79	0.19	72,74,88,94	0
5	CA	L	302	1/1	0.82	0.10	51,51,51,51	0

6.5 Other polymers [i](#)

There are no such residues in this entry.