



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 03:46 AM UTC

PDB ID : 3MT1 / pdb_00003mt1
Title : Crystal structure of putative carboxynorspermidine decarboxylase protein from *Sinorhizobium meliloti*
Authors : Chang, C.; Xu, X.; Cui, H.; Chin, S.; Savchenko, A.; Edwards, A.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2010-04-29
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	365	2% 79% 15% • 5%
1	B	365	% 82% 13% 5%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 5523 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative carboxynorspermidine decarboxylase protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	346	Total	C	N	O	S	Se	0	0	0
			2704	1719	464	508	3	10			
1	B	348	Total	C	N	O	S	Se	0	0	0
			2748	1740	475	519	4	10			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	30	Total 30	O 30	0	0
3	B	31	Total 31	O 31	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.77Å 75.77Å 129.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.50 50.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.3 (50.00-2.50) 98.3 (50.00-2.50)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.58 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.190 , 0.256 (Not available) , 0.231	Depositor DCC
R_{free} test set	1224 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	45.4	Xtriage
Anisotropy	0.086	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 34.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5523	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5.1 Standard geometry

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	363	SER	N-CA-C	-5.75	106.27	113.28
1	A	237	THR	CB-CA-C	-5.55	100.76	109.70
1	A	163	ASN	N-CA-C	5.18	116.52	110.41
1	A	236	ILE	N-CA-C	5.14	117.16	112.43

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	345	LEU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2704	0	2623	38	0
1	B	2748	0	2675	29	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	30	0	0	2	0
3	B	31	0	0	2	0
All	All	5523	0	5298	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:305:ARG:HD3	3:A:385:HOH:O	1.71	0.90
1:A:163:ASN:HD21	1:A:176:MSE:HB3	1.37	0.88
1:A:163:ASN:ND2	1:A:176:MSE:HB3	1.89	0.88
1:B:290:ILE:CD1	1:B:304:PHE:HD2	1.92	0.83
1:B:256:LEU:HD23	1:B:289:MSE:HB3	1.71	0.71
1:B:290:ILE:HD13	1:B:304:PHE:HD2	1.55	0.69

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:145:ASP:OD1	1:B:147:PRO:HD2	1.95	0.66
1:A:217:ARG:HH11	1:A:217:ARG:HG2	1.61	0.65
1:A:58:SER:HA	1:A:77:TYR:O	1.97	0.64
1:B:58:SER:HA	1:B:77:TYR:O	1.97	0.64
1:B:162:HIS:CE1	1:B:198:GLY:H	2.17	0.63
1:A:161:ILE:HD12	1:A:194:VAL:HG13	1.82	0.61
1:A:112:ALA:HA	1:A:116:ARG:NH1	2.16	0.60
1:A:146:VAL:HB	1:A:147:PRO:HD3	1.84	0.59
1:B:290:ILE:HD13	1:B:304:PHE:CD2	2.35	0.59
1:A:24:VAL:HG22	1:A:209:VAL:HG13	1.85	0.58
1:A:56:THR:HG21	1:A:230:GLU:HG2	1.86	0.57
1:B:86:ILE:HG21	1:B:110:LYS:HD3	1.88	0.56
1:A:240:THR:HB	1:A:322:ALA:HB2	1.89	0.55
1:A:195:SER:HB3	1:A:228:TYR:HB2	1.87	0.55
1:A:144:TRP:HE3	1:A:144:TRP:H	1.56	0.54
1:B:205:ASP:O	1:B:206:ASP:HB2	2.08	0.53
1:B:240:THR:HB	1:B:322:ALA:HB2	1.91	0.53
1:A:112:ALA:HA	1:A:116:ARG:HH12	1.74	0.52
1:B:244:VAL:O	1:B:315:ARG:HD3	2.11	0.51
1:B:266:HIS:CD2	1:B:338:PRO:HD3	2.45	0.51
1:B:240:THR:CB	1:B:322:ALA:HB2	2.41	0.50
1:B:294:SER:CB	1:B:299:ASP:OD2	2.59	0.50
1:B:74:THR:HG22	3:B:367:HOH:O	2.11	0.50
1:A:75:HIS:CE1	1:A:95:LYS:HD2	2.47	0.49
1:B:77:TYR:CG	1:B:160:MSE:HE1	2.47	0.49
1:A:217:ARG:HH11	1:A:217:ARG:CG	2.25	0.49
1:B:146:VAL:HB	1:B:147:PRO:HD3	1.94	0.49
1:B:155:ARG:HD2	3:B:396:HOH:O	2.14	0.48
1:A:288:TYR:OH	1:A:309:GLU:HG2	2.14	0.47
1:A:150:GLU:HA	1:A:153:MSE:HG2	1.97	0.47
1:B:54:GLY:HA3	1:B:73:GLU:O	2.14	0.47
1:B:163:ASN:OD1	1:B:163:ASN:N	2.44	0.47
1:A:122:PRO:HG3	1:A:161:ILE:HG22	1.96	0.46
1:A:101:ILE:HG23	1:A:152:VAL:HG11	1.97	0.46
1:A:18:MSE:HB3	1:A:51:TYR:CG	2.51	0.46
1:A:236:ILE:HD12	1:A:236:ILE:C	2.41	0.46
1:B:290:ILE:CD1	1:B:304:PHE:CD2	2.84	0.46
1:B:74:THR:OG1	1:B:93:ALA:HA	2.16	0.45
1:A:240:THR:CB	1:A:322:ALA:HB2	2.45	0.45
1:B:158:GLY:HA3	1:B:193:TRP:CE2	2.52	0.45
1:A:87:ASP:HB2	3:A:382:HOH:O	2.17	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:78:SER:HB3	1:A:81:TYR:CE2	2.53	0.44
1:A:280:LEU:HA	1:A:281:PRO:HA	1.78	0.44
1:A:54:GLY:HA3	1:A:73:GLU:O	2.19	0.43
1:A:332:PHE:O	1:A:333:ASN:HB2	2.17	0.43
1:B:149:VAL:O	1:B:152:VAL:HG12	2.19	0.42
1:A:236:ILE:HD13	1:A:322:ALA:HB3	2.02	0.42
1:A:8:LEU:O	1:A:240:THR:HA	2.20	0.42
1:A:95:LYS:HD3	1:A:193:TRP:CZ2	2.55	0.42
1:B:266:HIS:NE2	1:B:338:PRO:HD3	2.34	0.42
1:B:95:LYS:HG2	1:B:115:ALA:HB3	2.02	0.41
1:A:115:ALA:HB1	1:A:157:ASN:HD22	1.86	0.41
1:A:47:LEU:C	1:A:47:LEU:HD23	2.46	0.41
1:B:236:ILE:HG23	1:B:322:ALA:HB3	2.02	0.41
1:A:290:ILE:HD12	1:A:304:PHE:HD2	1.85	0.41
1:A:18:MSE:HB3	1:A:51:TYR:CD2	2.56	0.41
1:B:178:GLY:O	1:B:182:GLU:HG3	2.21	0.41
1:A:187:LEU:HA	1:A:187:LEU:HD23	1.86	0.41
1:A:34:LEU:HB3	1:A:55:THR:HG22	2.03	0.41
1:B:67:ARG:CZ	1:B:74:THR:HG23	2.51	0.40
1:A:77:TYR:CG	1:A:78:SER:N	2.89	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	307/365 (84%)	288 (94%)	17 (6%)	2 (1%)	18	34
1	B	342/365 (94%)	325 (95%)	15 (4%)	2 (1%)	21	38
All	All	649/730 (89%)	613 (94%)	32 (5%)	4 (1%)	21	38

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	346	ASP
1	A	299	ASP
1	A	293	LYS
1	B	157	ASN

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	401	-	4,4,4	0.14	0	6,6,6	0.41	0
2	SO4	A	401	-	4,4,4	0.29	0	6,6,6	0.56	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å²)	Q<0.9
1	A	336/365 (92%)	-0.18	6 (1%) 6764	25, 45, 72, 84	2 (0%)
1	B	337/365 (92%)	-0.21	4 (1%) 7673	24, 45, 73, 88	0
All	All	673/730 (92%)	-0.19	10 (1%) 7268	24, 45, 73, 88	2 (0%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	179	GLU	4.4
1	A	248	ASP	3.2
1	B	165	CYS	3.0
1	B	252	ASN	2.6
1	A	198	GLY	2.4
1	B	123	GLN	2.3
1	A	2	ILE	2.2
1	B	2	ILE	2.1
1	A	168	LYS	2.0
1	A	169	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	B	401	5/5	0.94	0.08	49,50,51,53	0
2	SO4	A	401	5/5	0.95	0.08	51,51,52,53	0

6.5 Other polymers (i)

There are no such residues in this entry.