



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 26, 2026 – 09:48 AM EDT

PDB ID : 5MTW / pdb_00005mtw
Title : Mycobacterium tuberculosis Rv1957 SecB-like chaperone in complex with a ChAD peptide from Rv1956 HigA1 antitoxin
Authors : Guillet, V.; Mourey, L.
Deposited on : 2017-01-10
Resolution : 1.84 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	1296 (1.84-1.84)
Clashscore	190562	1329 (1.84-1.84)
Ramachandran outliers	187476	1318 (1.84-1.84)
Sidechain outliers	187428	1318 (1.84-1.84)
RSRZ outliers	180081	1296 (1.84-1.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	184	9% 68% 5% 25%
1	B	184	15% 63% 10% 25%
1	C	184	11% 62% 11% 25%
1	D	184	15% 62% 10% 26%
2	E	13	31% 62% 31% 8%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	13	23% 85% 8% 8%
2	G	13	23% 46% 46% 8%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4656 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SecB-like chaperone Rv1957.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	138	Total	C	N	O	S	0	0	0
			1060	676	180	203	1			
1	D	136	Total	C	N	O	S	0	2	0
			1069	683	184	201	1			
1	A	138	Total	C	N	O	S	0	0	0
			1072	682	184	205	1			
1	B	138	Total	C	N	O	S	0	0	0
			1077	686	186	204	1			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	GLY	-	expression tag	UNP P95257
C	-1	SER	-	expression tag	UNP P95257
C	0	HIS	-	expression tag	UNP P95257
D	-2	GLY	-	expression tag	UNP P95257
D	-1	SER	-	expression tag	UNP P95257
D	0	HIS	-	expression tag	UNP P95257
A	-2	GLY	-	expression tag	UNP P95257
A	-1	SER	-	expression tag	UNP P95257
A	0	HIS	-	expression tag	UNP P95257
B	-2	GLY	-	expression tag	UNP P95257
B	-1	SER	-	expression tag	UNP P95257
B	0	HIS	-	expression tag	UNP P95257

- Molecule 2 is a protein called Antitoxin HigA1.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	12	Total	C	N	O	0	0	0
			108	69	21	18			
2	F	12	Total	C	N	O	0	0	0
			108	69	21	18			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	G	12	Total	C	N	O	0	0	0
			102	66	18	18			

- Molecule 3 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	C	1	Total	C	O	S	0	0
			4	2	1	1		
3	B	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	1	Total	Ca	0	0
			1	1		
4	A	1	Total	Ca	0	0
			1	1		
4	B	1	Total	Ca	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	13	Total	O	0	0
			13	13		

Continued on next page...

Continued from previous page...



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	16	Total 16	O 16	0	0
5	A	11	Total 11	O 11	0	0
5	B	4	Total 4	O 4	0	0
5	E	1	Total 1	O 1	0	0
5	G	4	Total 4	O 4	0	0



● Molecule 2: Antitoxin HigA1



● Molecule 2: Antitoxin HigA1



● Molecule 2: Antitoxin HigA1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	86.86Å 89.96Å 91.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	64.09 – 1.84 64.09 – 1.84	Depositor EDS
% Data completeness (in resolution range)	99.9 (64.09-1.84) 99.9 (64.09-1.84)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 1.83Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.212 , 0.265 0.217 , 0.267	Depositor DCC
R_{free} test set	3397 reflections (5.37%)	wwPDB-VP
Wilson B-factor (Å ²)	27.6	Xtriage
Anisotropy	0.207	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 44.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.024 for -h,l,k 0.027 for -l,-k,-h 0.025 for k,h,-l 0.009 for k,l,h 0.009 for l,h,k	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4656	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.30	5/1092 (0.5%)	1.16	1/1489 (0.1%)
1	B	1.38	5/1097 (0.5%)	1.18	7/1496 (0.5%)
1	C	1.42	5/1081 (0.5%)	1.21	2/1477 (0.1%)
1	D	1.33	2/1097 (0.2%)	1.15	3/1496 (0.2%)
2	E	1.33	1/112 (0.9%)	0.94	0/152
2	F	1.00	0/112	1.00	1/152 (0.7%)
2	G	1.42	1/106 (0.9%)	0.99	0/145
All	All	1.35	19/4697 (0.4%)	1.16	14/6407 (0.2%)

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	141	ILE	C-O	-6.68	1.16	1.24
1	B	155	PRO	CA-C	6.57	1.58	1.52
1	B	25	ILE	N-CA	6.38	1.53	1.46
1	A	102	ASP	CA-C	6.22	1.59	1.52
1	A	143	GLU	C-O	-6.08	1.16	1.24
2	E	106	PRO	CA-C	6.01	1.58	1.52
1	C	32	ARG	N-CA	-5.89	1.38	1.46
1	C	107	ALA	CA-C	5.60	1.59	1.52
1	C	79	ASN	N-CA	5.47	1.52	1.46
1	C	96	GLN	CA-C	5.39	1.59	1.52
1	D	144	TYR	CA-C	5.33	1.59	1.52
1	A	139	PRO	CA-C	5.32	1.60	1.52
1	B	131	THR	CA-C	5.26	1.55	1.52
1	D	35	ALA	N-CA	5.15	1.52	1.45
1	B	40	ALA	C-N	5.14	1.40	1.33
2	G	109	HIS	C-O	-5.13	1.17	1.23
1	A	26	ARG	C-O	-5.12	1.15	1.23
1	B	140	TYR	C-O	-5.11	1.17	1.24
1	A	117	GLU	C-O	-5.00	1.18	1.23

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	40	ALA	CA-C-N	-7.20	112.56	119.76
1	B	40	ALA	C-N-CA	-7.20	112.56	119.76
1	C	63	THR	CB-CA-C	-6.41	99.40	109.34
1	A	151	ARG	N-CA-C	6.10	117.93	111.28
1	C	47	LEU	CB-CA-C	5.78	119.72	109.72
1	D	155	PRO	CA-C-N	-5.26	114.15	119.83
1	D	155	PRO	C-N-CA	-5.26	114.15	119.83
2	F	114	TYR	N-CA-C	5.23	118.08	109.72
1	B	119	ASP	CA-C-N	-5.22	114.54	119.76
1	B	119	ASP	C-N-CA	-5.22	114.54	119.76
1	B	137	LEU	N-CA-C	5.20	118.78	112.23
1	B	64	ILE	N-CA-C	5.14	115.98	108.53
1	B	60	ASP	N-CA-C	-5.13	98.46	109.81
1	D	44	ALA	N-CA-C	5.13	119.39	113.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1072	0	1029	9	0
1	B	1077	0	1040	9	0
1	C	1060	0	1015	14	0
1	D	1069	0	1040	14	0
2	E	108	0	101	4	0
2	F	108	0	101	0	0
2	G	102	0	90	7	1
3	B	4	0	6	0	0
3	C	4	0	6	1	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	D	1	0	0	0	0
5	A	11	0	0	1	1
5	B	4	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	13	0	0	0	0
5	D	16	0	0	1	0
5	E	1	0	0	0	0
5	G	4	0	0	0	0
All	All	4656	0	4428	45	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:47:LEU:HD21	2:E:114:TYR:CD2	2.06	0.90
1:A:19:LEU:HD11	1:A:120:PRO:HB3	1.58	0.84
1:A:26:ARG:NE	1:A:110:ASP:OD2	2.14	0.81
1:C:47:LEU:HD21	2:E:114:TYR:CE2	2.19	0.77
1:D:44:ALA:HB2	2:G:111:LEU:HD22	1.67	0.77
1:D:147:ASP:OD1	5:D:301:HOH:O	2.07	0.71
1:B:114:GLN:HE21	1:B:114:GLN:HA	1.56	0.70
1:D:161:ILE:H	1:B:24:GLN:HE22	1.40	0.67
1:A:26:ARG:HG3	1:A:110:ASP:OD2	1.95	0.66
1:C:18:ARG:NH2	1:C:118:ASP:O	2.27	0.65
1:C:47:LEU:CD2	2:E:114:TYR:CD2	2.78	0.64
1:B:24:GLN:HE21	1:B:25:ILE:H	1.45	0.63
1:B:50:ASP:OD2	1:B:76:ARG:NH2	2.33	0.61
1:C:28:ILE:HD13	1:C:140:TYR:CE1	2.35	0.60
1:C:47:LEU:C	1:C:47:LEU:HD23	2.26	0.60
1:A:26:ARG:CG	1:A:110:ASP:OD2	2.50	0.60
1:A:116:GLY:O	5:A:301:HOH:O	2.17	0.59
1:B:58:ASP:OD2	1:B:70:ARG:NH1	2.37	0.56
1:D:44:ALA:CB	2:G:111:LEU:HD22	2.34	0.56
1:C:71:ILE:HD11	1:C:134:ARG:HA	1.89	0.53
1:C:28:ILE:HD13	1:C:140:TYR:CZ	2.45	0.51
1:C:113:LEU:C	1:C:117:GLU:HG2	2.35	0.51
1:C:47:LEU:HD21	2:E:114:TYR:CG	2.44	0.50
1:D:22:ARG:NE	1:D:117:GLU:OE2	2.45	0.50
1:D:43:PRO:HB3	2:G:111:LEU:N	2.27	0.50
3:C:201:DMS:H13	1:B:151:ARG:HG2	1.95	0.48
1:D:53:PHE:N	1:D:53:PHE:CD1	2.81	0.47
1:B:24:GLN:HE21	1:B:25:ILE:N	2.13	0.47
1:D:55:PRO:HA	1:D:71:ILE:HD13	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:118:ASP:N	1:D:118:ASP:OD1	2.47	0.46
1:A:147:ASP:O	1:A:151:ARG:HG3	2.16	0.45
1:B:28:ILE:HD13	1:B:140:TYR:CE1	2.51	0.45
1:D:76:ARG:NH1	1:D:97:ASP:OD2	2.51	0.43
1:D:44:ALA:HB3	2:G:111:LEU:HD13	2.01	0.42
1:C:117:GLU:HG3	1:C:118:ASP:N	2.33	0.42
1:D:74:HIS:ND1	1:D:102:ASP:OD1	2.38	0.42
1:D:41:PRO:HG3	1:D:154:LEU:HD23	2.02	0.42
1:A:26:ARG:CD	1:A:110:ASP:OD2	2.68	0.41
1:C:165:PRO:O	1:C:166:MET:C	2.64	0.41
1:B:42:LYS:H	1:B:79:ASN:HD21	1.68	0.41
1:C:138:TYR:CZ	1:C:162:LEU:HG	2.56	0.41
1:A:11:ASP:OD1	1:A:11:ASP:C	2.64	0.41
1:A:47:LEU:HD12	2:G:112:SER:OG	2.21	0.41
1:C:64:ILE:HD11	2:G:105:VAL:HB	2.02	0.40
1:D:43:PRO:HB3	2:G:110:ARG:C	2.46	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:G:104:GLU:OE2	5:A:301:HOH:O[2_575]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	132/184 (72%)	129 (98%)	3 (2%)	0	100	100
1	B	132/184 (72%)	127 (96%)	4 (3%)	1 (1%)	16	5
1	C	132/184 (72%)	129 (98%)	3 (2%)	0	100	100
1	D	132/184 (72%)	130 (98%)	2 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	E	10/13 (77%)	10 (100%)	0	0	100	100
2	F	10/13 (77%)	10 (100%)	0	0	100	100
2	G	10/13 (77%)	10 (100%)	0	0	100	100
All	All	558/775 (72%)	545 (98%)	12 (2%)	1 (0%)	43	34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	114	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	106/149 (71%)	105 (99%)	1 (1%)	70	60
1	B	107/149 (72%)	103 (96%)	4 (4%)	30	13
1	C	104/149 (70%)	98 (94%)	6 (6%)	18	4
1	D	107/149 (72%)	102 (95%)	5 (5%)	23	7
2	E	12/12 (100%)	10 (83%)	2 (17%)	2	0
2	F	12/12 (100%)	12 (100%)	0	100	100
2	G	11/12 (92%)	11 (100%)	0	100	100
All	All	459/632 (73%)	441 (96%)	18 (4%)	28	11

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	37	VAL
1	C	45	GLN
1	C	47	LEU
1	C	63	THR
1	C	117	GLU
1	C	163	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	45	GLN
1	D	53	PHE
1	D	113	LEU
1	D	118	ASP
1	D	122	GLU
1	A	11	ASP
1	B	10	LEU
1	B	63	THR
1	B	114	GLN
1	B	162	LEU
2	E	111	LEU
2	E	115	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	34	GLN
1	D	34	GLN
1	D	79	ASN
1	A	79	ASN
1	B	24	GLN
1	B	74	HIS
1	B	79	ASN
1	B	112	HIS
1	B	114	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	DMS	B	201	-	3,3,3	0.63	0	3,3,3	0.96	0
3	DMS	C	201	-	3,3,3	0.74	0	3,3,3	0.63	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	201	DMS	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	138/184 (75%)	0.69	17 (12%) 8 8	16, 31, 62, 72	7 (5%)
1	B	138/184 (75%)	1.26	27 (19%) 3 3	18, 32, 54, 69	12 (8%)
1	C	138/184 (75%)	0.72	21 (15%) 5 6	15, 30, 61, 75	7 (5%)
1	D	136/184 (73%)	1.09	28 (20%) 2 2	10, 29, 56, 69	16 (11%)
2	E	12/13 (92%)	1.72	4 (33%) 1 1	29, 37, 46, 70	2 (16%)
2	F	12/13 (92%)	1.73	3 (25%) 2 2	24, 39, 53, 65	1 (8%)
2	G	12/13 (92%)	1.55	3 (25%) 2 2	21, 26, 41, 45	3 (25%)
All	All	586/775 (75%)	0.99	103 (17%) 4 3	10, 31, 59, 75	48 (8%)

All (103) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	80	GLN	18.1
1	D	79	ASN	14.1
1	B	7	ALA	13.0
1	B	42	LYS	12.7
1	D	116	GLY	11.0
2	G	115	ARG	10.1
1	B	6	ASP	9.8
1	B	8	ASP	9.7
1	B	112	HIS	9.0
2	F	115	ARG	8.2
2	E	115	ARG	6.9
1	B	22	ARG	5.6
1	A	9	ASP	5.6
1	B	9	ASP	5.5
1	C	13	GLN	5.1
1	B	114	GLN	5.0
2	G	113	SER	4.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	10	LEU	4.9
1	D	45	GLN	4.7
1	C	113	LEU	4.7
1	C	117	GLU	4.6
1	B	115	GLU	4.6
1	B	10	LEU	4.3
1	C	44	ALA	4.3
1	D	117	GLU	4.2
2	E	109	HIS	4.1
1	A	53	PHE	4.1
1	C	118	ASP	4.1
1	C	80	GLN	4.0
1	B	130	ALA	3.8
1	C	166	MET	3.7
1	C	95	THR	3.7
1	C	81	ALA	3.6
1	A	81	ALA	3.6
1	B	39	ARG	3.6
1	C	45	GLN	3.4
1	C	62	ALA	3.4
1	A	42	LYS	3.4
1	D	74	HIS	3.3
1	A	45	GLN	3.3
1	C	47	LEU	3.3
1	A	80	GLN	3.2
1	B	62	ALA	3.2
1	B	64	ILE	3.2
1	B	113	LEU	3.2
1	D	40	ALA	3.1
1	D	14	ARG	3.1
1	D	47	LEU	3.1
1	D	53	PHE	3.1
1	D	46	GLY	3.1
1	D	48	THR	3.1
1	C	46	GLY	3.0
1	B	12	LEU	2.9
2	F	109	HIS	2.9
1	C	165	PRO	2.8
1	B	63	THR	2.8
2	E	111	LEU	2.8
1	D	112	HIS	2.7
1	A	26	ARG	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	G	111	LEU	2.7
1	A	79	ASN	2.7
1	D	118	ASP	2.6
2	E	104	GLU	2.6
1	B	53	PHE	2.6
1	D	43	PRO	2.6
1	B	134	ARG	2.6
1	B	80	GLN	2.6
1	D	18	ARG	2.5
1	A	11	ASP	2.5
1	D	162	LEU	2.5
1	B	116	GLY	2.4
1	C	112	HIS	2.4
1	D	12	LEU	2.4
1	A	47	LEU	2.4
1	C	163	SER	2.4
1	A	163	SER	2.4
1	D	10	LEU	2.4
1	D	50	ASP	2.4
1	A	162	LEU	2.4
1	D	76	ARG	2.4
1	D	113	LEU	2.3
1	B	48	THR	2.3
1	B	162	LEU	2.3
1	C	22	ARG	2.2
1	B	17	ALA	2.2
1	A	97	ASP	2.2
1	B	11	ASP	2.2
1	D	44	ALA	2.2
1	D	13	GLN	2.1
1	D	78	GLN	2.1
1	A	102	ASP	2.1
1	B	61	PRO	2.1
1	C	119	ASP	2.1
2	F	111	LEU	2.1
1	C	63	THR	2.1
1	D	15	VAL	2.1
1	C	162	LEU	2.1
1	D	135	PHE	2.0
1	A	46	GLY	2.0
1	C	102	ASP	2.0
1	D	54	GLU	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	48	THR	2.0
1	B	37	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	CA	B	202	1/1	0.89	0.12	58,58,58,58	0
3	DMS	C	201	4/4	0.90	0.18	58,64,65,69	0
3	DMS	B	201	4/4	0.92	0.17	57,60,65,66	0
4	CA	A	201	1/1	0.95	0.26	41,41,41,41	1
4	CA	D	201	1/1	0.96	0.06	37,37,37,37	0

6.5 Other polymers [i](#)

There are no such residues in this entry.