



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 05:38 PM UTC

PDB ID : 4MUI / pdb_00004mui
Title : Crystal structure of pantothenate synthetase in complex with 2-(5-methoxy-2-(4-methoxyphenylsulfonylcarbamoyl)-1H-indol-1-yl)acetic acid
Authors : Silvestre, H.L.; Blundell, T.L.
Deposited on : 2013-09-22
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	300	5% 78% 14% 6%
1	B	300	4% 76% 13% 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	EOH	A	401	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4524 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pantothenate synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	283	Total	C	N	O	S	0	3	0
			2108	1337	378	387	6			
1	B	272	Total	C	N	O	S	0	2	0
			2016	1277	366	368	5			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	2	ALA	THR	engineered mutation	UNP P0A5R0
A	77	GLY	GLU	engineered mutation	UNP P0A5R0
B	2	ALA	THR	engineered mutation	UNP P0A5R0
B	77	GLY	GLU	engineered mutation	UNP P0A5R0

- Molecule 2 is ETHANOL (CCD ID: EOH) (formula: C₂H₆O).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 3 2 1	0	0
2	A	1	Total C O 3 2 1	0	0
2	A	1	Total C O 3 2 1	0	0
2	A	1	Total C O 3 2 1	0	0
2	B	1	Total C O 3 2 1	0	0

- Molecule 3 is (5-methoxy-2-[(4-methoxyphenyl)sulfonyl]carbamoyl}-1H-indol-1-yl)acetic acid (CCD ID: 2DQ) (formula: C₁₉H₁₈N₂O₇S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O S 29 19 2 7 1	0	0
3	B	1	Total C N O S 29 19 2 7 1	0	0
3	B	1	Total C N O S 29 19 2 7 1	0	0

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	148	Total	O	0	0
			148	148		
5	B	138	Total	O	0	0
			138	138		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pantothenate synthetase



• Molecule 1: Pantothenate synthetase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	48.97Å 70.69Å 81.69Å 90.00° 99.91° 90.00°	Depositor
Resolution (Å)	30.00 – 2.10 30.00 – 2.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (30.00-2.10) 100.0 (30.00-2.10)	Depositor EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.53 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.186 , 0.251 0.186 , 0.249	Depositor DCC
R_{free} test set	1631 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	25.4	Xtriage
Anisotropy	0.152	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 49.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4524	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EOH, 2DQ, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.18	0/2146	1.16	7/2931 (0.2%)
1	B	1.22	3/2053 (0.1%)	1.16	6/2803 (0.2%)
All	All	1.20	3/4199 (0.1%)	1.16	13/5734 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	37	VAL	CA-CB	5.44	1.61	1.54
1	B	118	VAL	CA-CB	5.28	1.60	1.54
1	B	103	THR	CA-CB	5.06	1.59	1.53

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	121	GLY	CA-C-N	6.99	127.57	119.47
1	B	121	GLY	C-N-CA	6.99	127.57	119.47
1	B	51	VAL	N-CA-C	6.84	116.99	110.42
1	A	69	ASN	CA-C-N	-6.82	112.60	119.56
1	A	69	ASN	C-N-CA	-6.82	112.60	119.56
1	A	132	ARG	CA-C-N	5.47	125.12	119.05
1	A	132	ARG	C-N-CA	5.47	125.12	119.05
1	A	85	THR	CA-C-N	5.35	125.02	119.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	85	THR	C-N-CA	5.35	125.02	119.56
1	B	130	GLY	CA-C-N	5.22	124.72	119.19
1	B	130	GLY	C-N-CA	5.22	124.72	119.19
1	A	247	VAL	N-CA-C	5.18	115.78	109.30
1	B	174	ASP	N-CA-C	5.05	116.78	111.28

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	289	THR	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2108	0	2153	37	0
1	B	2016	0	2081	37	0
2	A	12	0	24	2	0
2	B	3	0	6	0	0
3	A	29	0	17	2	0
3	B	58	0	34	6	0
4	A	4	0	6	1	0
4	B	8	0	12	2	0
5	A	148	0	0	5	0
5	B	138	0	0	3	0
All	All	4524	0	4333	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:278:ARG:HG2	1:B:278:ARG:HH11	1.32	0.92

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:115:ARG:O	2:A:401:EOH:H22	1.70	0.92
1:B:278:ARG:HG2	1:B:278:ARG:NH1	1.90	0.83
1:A:114:LEU:HD21	5:A:646:HOH:O	1.83	0.78
1:B:44:HIS:HD2	1:B:46:GLY:H	1.36	0.73
1:B:44:HIS:H	1:B:47:HIS:HD1	1.37	0.72
1:B:221:ALA:HA	1:B:285:ILE:HD13	1.73	0.70
1:B:278:ARG:HH11	1:B:278:ARG:CG	2.05	0.69
1:A:263:ASN:HD22	1:A:263:ASN:H	1.39	0.69
4:B:401:EDO:H22	3:B:402:2DQ:H15	1.75	0.68
1:B:263:ASN:H	1:B:263:ASN:ND2	1.95	0.65
1:A:44:HIS:HD2	1:A:46:GLY:H	1.44	0.65
1:A:286:GLU:HG2	1:A:290:PHE:CD2	2.33	0.63
1:B:187:VAL:HG12	1:B:195:MET:HG3	1.79	0.63
1:A:151:ARG:HH22	1:B:151:ARG:CD	2.12	0.62
1:A:157:PHE:CZ	1:A:168:ILE:HD13	2.36	0.61
1:A:255:ILE:HG13	5:A:648:HOH:O	2.01	0.60
1:A:151:ARG:HH22	1:B:151:ARG:HD2	1.65	0.60
1:B:52:ARG:O	1:B:56:ARG:HG3	2.01	0.59
1:B:3:ILE:HD13	1:B:26:ALA:HB2	1.84	0.58
1:A:3:ILE:HG22	1:A:22:ASP:HB3	1.84	0.58
1:B:217:LEU:HD23	1:B:268:LEU:HD23	1.87	0.57
1:A:44:HIS:H	1:A:47:HIS:HD1	1.51	0.57
1:B:160:LYS:HE3	5:B:595:HOH:O	2.05	0.56
1:A:117:THR:OG1	2:A:401:EOH:H21	2.05	0.56
1:A:44:HIS:HE1	3:A:405:2DQ:O	1.88	0.56
1:A:47:HIS:NE2	3:A:405:2DQ:NAQ	2.50	0.56
1:B:263:ASN:ND2	1:B:263:ASN:N	2.54	0.55
1:B:254:ASP:HB3	1:B:261:PRO:HD3	1.87	0.55
1:B:278:ARG:NH1	1:B:278:ARG:CG	2.61	0.54
1:A:197:SER:O	1:A:200:ARG:HG2	2.09	0.52
1:A:286:GLU:HG2	1:A:290:PHE:CG	2.43	0.52
1:A:24:SER:OG	1:A:28:ARG:NH2	2.42	0.52
1:A:174:ASP:HB3	4:B:404:EDO:H21	1.92	0.51
1:B:46:GLY:O	3:B:402:2DQ:H7	2.11	0.50
3:B:403:2DQ:H3	3:B:403:2DQ:OAD	2.12	0.50
1:A:160[A]:LYS:HE3	5:A:528:HOH:O	2.11	0.49
1:A:114:LEU:HD22	1:A:117:THR:HG22	1.94	0.49
1:A:286:GLU:HB3	1:A:290:PHE:HB3	1.94	0.49
1:A:27:LEU:HD13	1:A:61:VAL:HG11	1.94	0.48
3:B:403:2DQ:OAD	3:B:403:2DQ:CA	2.61	0.48
1:B:263:ASN:H	1:B:263:ASN:HD22	1.58	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:40:MET:HE2	3:B:402:2DQ:H18	1.95	0.48
1:A:69:ASN:HD22	1:A:72:GLN:HE21	1.61	0.47
1:A:131:PRO:HG2	1:A:132:ARG:NH2	2.29	0.47
1:B:124:ALA:O	1:B:129:GLY:HA3	2.15	0.47
1:B:217:LEU:HD23	1:B:268:LEU:CD2	2.44	0.47
1:A:262:LEU:HD21	1:A:287:ILE:HG13	1.97	0.46
1:A:151:ARG:NH2	1:B:151:ARG:HD2	2.29	0.46
1:B:191:ASP:OD2	1:B:211[B]:VAL:CG2	2.64	0.46
1:A:202:LEU:HD22	1:A:206:GLN:HB3	1.98	0.45
1:B:132:ARG:HB3	1:B:135:HIS:HB2	1.99	0.45
1:A:120:PRO:HB2	1:A:124:ALA:HB2	1.99	0.44
1:B:44:HIS:HE1	3:B:402:2DQ:OXT	2.00	0.44
1:A:268:LEU:HB2	1:A:285:ILE:HG13	2.00	0.43
1:B:191:ASP:OD2	1:B:211[B]:VAL:HG22	2.18	0.43
1:A:45:GLU:HG3	1:A:187:VAL:HG11	2.01	0.43
1:A:290:PHE:HD1	1:A:290:PHE:HA	1.74	0.43
1:A:43:LEU:HA	1:A:47:HIS:HD1	1.84	0.43
1:B:287:ILE:HD13	1:B:287:ILE:HA	1.87	0.43
1:A:221:ALA:HA	1:A:285:ILE:HD13	2.01	0.42
1:B:44:HIS:CE1	1:B:47:HIS:CE1	3.07	0.42
1:A:174:ASP:OD2	1:B:115:ARG:HB3	2.19	0.42
1:B:269:LEU:N	1:B:269:LEU:HD12	2.35	0.42
4:A:406:EDO:H12	5:A:639:HOH:O	2.19	0.41
1:B:148:GLN:OE1	5:B:622:HOH:O	2.22	0.41
1:B:208:ALA:O	1:B:211[A]:VAL:HG12	2.20	0.41
1:A:148:GLN:NE2	5:A:552:HOH:O	2.49	0.41
1:B:37:VAL:HB	1:B:64:VAL:HG22	2.02	0.41
1:A:232:LEU:O	1:A:236:ARG:HG3	2.20	0.41
1:B:151:ARG:NH2	5:B:604:HOH:O	2.52	0.41
1:A:255:ILE:HG23	1:A:266:GLY:HA2	2.03	0.41
1:B:33:ARG:HH11	1:B:33:ARG:HA	1.85	0.40
1:B:44:HIS:CD2	1:B:46:GLY:H	2.26	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	280/300 (93%)	271 (97%)	8 (3%)	1 (0%)	30	28
1	B	270/300 (90%)	264 (98%)	6 (2%)	0	100	100
All	All	550/600 (92%)	535 (97%)	14 (2%)	1 (0%)	43	44

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	261	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	215/223 (96%)	205 (95%)	10 (5%)	23	24
1	B	205/223 (92%)	196 (96%)	9 (4%)	25	26
All	All	420/446 (94%)	401 (96%)	19 (4%)	24	25

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ILE
1	A	30	THR
1	A	114	LEU
1	A	236	ARG
1	A	254	ASP
1	A	260	MET
1	A	263	ASN
1	A	285	ILE
1	A	289	THR
1	A	290	PHE
1	B	32	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	33	ARG
1	B	87	ASP
1	B	88	ASP
1	B	142	VAL
1	B	160	LYS
1	B	263	ASN
1	B	278	ARG
1	B	285	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	44	HIS
1	A	72	GLN
1	A	148	GLN
1	A	222	HIS
1	A	263	ASN
1	B	44	HIS
1	B	148	GLN
1	B	263	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	2DQ	B	403	-	31,31,31	3.44	14 (45%)	44,45,45	2.19	9 (20%)
2	EOH	A	401	-	2,2,2	0.52	0	1,1,1	0.16	0
2	EOH	A	404	-	2,2,2	0.52	0	1,1,1	0.31	0
2	EOH	B	405	-	2,2,2	0.38	0	1,1,1	0.55	0
4	EDO	B	401	-	3,3,3	0.46	0	2,2,2	0.49	0
4	EDO	B	404	-	3,3,3	0.37	0	2,2,2	0.60	0
2	EOH	A	402	-	2,2,2	0.56	0	1,1,1	0.39	0
3	2DQ	B	402	-	31,31,31	2.92	10 (32%)	44,45,45	2.83	16 (36%)
2	EOH	A	403	-	2,2,2	0.56	0	1,1,1	0.27	0
4	EDO	A	406	-	3,3,3	0.44	0	2,2,2	0.50	0
3	2DQ	A	405	-	31,31,31	2.70	13 (41%)	44,45,45	2.77	9 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	2DQ	B	403	-	-	0/23/23/23	0/3/3/3
4	EDO	B	401	-	-	0/1/1/1	-
4	EDO	B	404	-	-	1/1/1/1	-
3	2DQ	B	402	-	-	7/23/23/23	0/3/3/3
3	2DQ	A	405	-	-	6/23/23/23	0/3/3/3
4	EDO	A	406	-	-	0/1/1/1	-

All (37) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	403	2DQ	OAF-SBC	8.59	1.53	1.43
3	B	403	2DQ	OAE-SBC	8.40	1.53	1.43
3	B	402	2DQ	OAE-SBC	8.00	1.52	1.43
3	B	402	2DQ	OAF-SBC	7.94	1.52	1.43
3	B	403	2DQ	CAY-CAU	-6.81	1.35	1.49
3	A	405	2DQ	OAF-SBC	6.73	1.51	1.43
3	B	402	2DQ	CAU-NAQ	-5.81	1.32	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	405	2DQ	CAY-CAU	-5.49	1.38	1.49
3	B	402	2DQ	CAY-CAU	-5.43	1.38	1.49
3	B	403	2DQ	SBC-NAQ	5.41	1.77	1.64
3	B	403	2DQ	CAY-N	-4.86	1.33	1.39
3	A	405	2DQ	CAU-NAQ	-4.73	1.33	1.38
3	A	405	2DQ	CAZ-CBA	-4.65	1.34	1.41
3	A	405	2DQ	CAY-N	-4.55	1.33	1.39
3	B	403	2DQ	CAZ-CBA	-4.42	1.34	1.41
3	A	405	2DQ	OAE-SBC	3.94	1.48	1.43
3	B	403	2DQ	CAU-NAQ	-3.92	1.34	1.38
3	B	403	2DQ	CA-C	3.88	1.59	1.51
3	B	402	2DQ	CA-N	3.75	1.53	1.46
3	B	402	2DQ	CAM-CBA	-3.73	1.33	1.39
3	B	403	2DQ	CAM-CBA	-3.55	1.34	1.39
3	B	402	2DQ	CAN-CAZ	-3.48	1.34	1.40
3	B	403	2DQ	CAN-CAZ	-3.45	1.34	1.40
3	B	403	2DQ	CA-N	3.25	1.52	1.46
3	A	405	2DQ	CBA-N	-3.10	1.33	1.39
3	A	405	2DQ	CAN-CAW	3.09	1.44	1.39
3	B	403	2DQ	CAZ-CAO	-2.97	1.34	1.44
3	B	402	2DQ	CAZ-CBA	-2.90	1.37	1.41
3	A	405	2DQ	CAN-CAZ	-2.76	1.36	1.40
3	B	403	2DQ	CBA-N	-2.76	1.34	1.39
3	A	405	2DQ	CAM-CBA	-2.73	1.35	1.39
3	A	405	2DQ	SBC-NAQ	2.58	1.70	1.64
3	B	402	2DQ	CAZ-CAO	-2.57	1.36	1.44
3	A	405	2DQ	O-C	-2.42	1.22	1.30
3	B	403	2DQ	CAX-SBC	2.30	1.80	1.76
3	B	402	2DQ	CAX-SBC	2.28	1.80	1.76
3	A	405	2DQ	CAZ-CAO	-2.19	1.37	1.44

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	405	2DQ	OAF-SBC-OAE	-13.20	103.49	119.52
3	B	402	2DQ	OAF-SBC-OAE	-11.02	106.13	119.52
3	B	403	2DQ	OAF-SBC-OAE	-8.73	108.92	119.52
3	B	403	2DQ	CAU-NAQ-SBC	-7.58	113.36	123.76
3	B	402	2DQ	OAE-SBC-CAX	6.76	116.51	107.98
3	A	405	2DQ	CA-N-CBA	-5.88	116.19	125.18
3	A	405	2DQ	CAO-CAY-N	-5.81	106.14	109.42
3	B	402	2DQ	OAF-SBC-NAQ	4.86	120.69	106.77

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	402	2DQ	CA-N-CAY	4.41	134.13	127.43
3	B	402	2DQ	CAU-NAQ-SBC	-4.05	118.21	123.76
3	A	405	2DQ	CA-N-CAY	4.01	133.52	127.43
3	A	405	2DQ	OAF-SBC-NAQ	3.96	118.09	106.77
3	A	405	2DQ	OAE-SBC-CAX	3.82	112.79	107.98
3	B	402	2DQ	CA-N-CBA	-3.71	119.49	125.18
3	B	403	2DQ	CAY-CAU-NAQ	3.58	120.15	114.26
3	B	402	2DQ	CAY-CAU-NAQ	3.58	120.15	114.26
3	B	403	2DQ	CA-N-CAY	-3.23	122.53	127.43
3	B	402	2DQ	CBA-N-CAY	-3.16	106.12	108.44
3	B	403	2DQ	CA-N-CBA	3.14	129.99	125.18
3	B	403	2DQ	OAF-SBC-CAX	3.11	111.90	107.98
3	B	402	2DQ	CAK-CAX-SBC	3.03	123.09	119.76
3	B	402	2DQ	OAF-SBC-CAX	-2.96	104.25	107.98
3	B	402	2DQ	OAD-CAU-NAQ	-2.81	117.87	121.94
3	B	402	2DQ	CAA-OAR-CAV	-2.77	111.55	117.50
3	A	405	2DQ	CBA-N-CAY	2.50	110.28	108.44
3	B	403	2DQ	OAE-SBC-NAQ	2.40	113.64	106.77
3	B	402	2DQ	CAL-CAW-CAN	-2.37	117.31	120.50
3	B	402	2DQ	CAZ-CBA-N	2.16	109.19	107.85
3	B	402	2DQ	CAJ-CAH-CAV	2.14	122.18	119.73
3	A	405	2DQ	CAX-SBC-NAQ	2.14	109.47	106.06
3	B	402	2DQ	CAB-OAS-CAW	2.14	122.08	117.50
3	B	403	2DQ	CAK-CAI-CAV	2.12	122.15	119.73
3	A	405	2DQ	CAB-OAS-CAW	-2.09	113.02	117.50
3	B	403	2DQ	CAM-CAL-CAW	-2.05	117.39	119.73

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	405	2DQ	C-CA-N-CBA
3	B	402	2DQ	C-CA-N-CBA
3	B	402	2DQ	CAI-CAV-OAR-CAA
3	B	402	2DQ	CAH-CAV-OAR-CAA
3	A	405	2DQ	C-CA-N-CAY
3	B	402	2DQ	C-CA-N-CAY
3	A	405	2DQ	CAI-CAV-OAR-CAA
3	B	402	2DQ	CAU-NAQ-SBC-OAE
3	A	405	2DQ	CAH-CAV-OAR-CAA
3	A	405	2DQ	CAJ-CAX-SBC-OAE
3	A	405	2DQ	CAK-CAX-SBC-OAE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	B	404	EDO	O1-C1-C2-O2
3	B	402	2DQ	CAJ-CAX-SBC-OAE
3	B	402	2DQ	CAK-CAX-SBC-OAE


There are no ring outliers.

7 monomers are involved in 12 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	403	2DQ	2	0
2	A	401	EOH	2	0
4	B	401	EDO	1	0
4	B	404	EDO	1	0
3	B	402	2DQ	4	0
4	A	406	EDO	1	0
3	A	405	2DQ	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand 2DQ B 403



Ligand 2DQ B 402





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	283/300 (94%)	0.19	15 (5%) 32 34	11, 25, 46, 56	5 (1%)
1	B	272/300 (90%)	0.06	11 (4%) 42 44	7, 23, 43, 59	3 (1%)
All	All	555/600 (92%)	0.12	26 (4%) 36 38	7, 24, 45, 59	8 (1%)

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	258	GLY	5.9
1	A	261	PRO	4.9
1	A	81	ALA	4.4
1	A	290	PHE	4.2
1	B	70	PRO	3.8
1	B	262	LEU	3.3
1	A	79	LEU	3.1
1	B	131	PRO	3.0
1	B	30	THR	2.9
1	A	3	ILE	2.8
1	B	201	TYR	2.8
1	A	71	MET	2.7
1	A	70	PRO	2.7
1	A	260	MET	2.7
1	B	86	PRO	2.7
1	B	69	ASN	2.6
1	A	80	ASP	2.5
1	A	262	LEU	2.5
1	A	95	ALA	2.4
1	B	130	GLY	2.3
1	B	2	ALA	2.3
1	B	42	ALA	2.2
1	A	83	PRO	2.2
1	B	3	ILE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	90	LEU	2.2
1	A	72	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	EOH	A	402	3/3	0.63	0.25	48,48,48,49	0
2	EOH	A	404	3/3	0.78	0.17	45,45,46,47	0
2	EOH	B	405	3/3	0.78	0.18	31,31,32,33	0
3	2DQ	B	403	29/29	0.80	0.14	48,50,55,56	0
2	EOH	A	403	3/3	0.83	0.17	33,33,34,34	0
2	EOH	A	401	3/3	0.84	0.11	23,23,32,33	0
4	EDO	B	401	4/4	0.84	0.13	40,43,45,47	0
3	2DQ	A	405	29/29	0.92	0.08	25,28,34,34	0
3	2DQ	B	402	29/29	0.93	0.09	23,27,30,33	0
4	EDO	A	406	4/4	0.97	0.05	23,26,26,27	0
4	EDO	B	404	4/4	0.98	0.04	20,23,25,25	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 2DQ B 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 2DQ A 405:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 2DQ B 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.